

Influence of ionic liquid on polar organic compounds solubility in dense CO₂ phase

Małgorzata Ewa Zakrzewska^{a,b,*}, Yegor Borovkov^a, Ana Brandão Paninho^a, Zeljko Petrovski^a, Vesna Najdanovic-Visak^c, Ana Vital Morgado Nunes^a

^a LAQV-REQUIMTE, Departamento de Química, Faculdade de Ciências e Tecnologia, Universidade NOVA de Lisboa, Quinta da Torre, 2829-516 Caparica, Portugal


^b Centro de Química Estrutural, Instituto Superior Técnico, Universidade de Lisboa, Av. Rovisco Pais, 1049-001 Lisbon, Portugal

^c Chemical Engineering and Applied Chemistry, Energy & Bioproducts Research Institute, Aston University, Birmingham B4 7ET, United Kingdom

HIGHLIGHTS

- solubility of 1,2-butanediol and butylene carbonate in dense CO₂.
- phase behaviour of ternary and quaternary CO₂-IL biphasic mixtures.
- 1-hexyl-3-methylimidazolium tris(pentafluoroalkyl)trifluorophosphate ionic liquid as solubility modifier.

GRAPHICAL ABSTRACT



ARTICLE INFO

Keywords:

High-pressure carbon dioxide
Cyclic carbonate
Butanediol
Tris(pentafluoroethyl)trifluorophosphate ionic liquid
Multicomponent systems
Phase behaviour

ABSTRACT

Accurate measurement and prediction of the phase behaviour of mixtures involved in a chemical process are crucial for its optimisation. Given the importance of CO₂ conversion technologies and considering possible benefits of CO₂-ionic liquid biphasic systems, i.e., facilitating a product separation, we investigated the high-pressure behaviour of components of interest in a recently developed process of cyclic carbonate synthesis directly from CO₂ and potentially bio-based alcohols. The solubility of 1,2-butanediol and 1,2-butylene carbonate in a dense carbon dioxide phase was determined experimentally at the temperature of 313.2 K and pressures between 6 and 18 MPa. The influence of 1-hexyl-3-methylimidazolium tris(pentafluoroethyl)trifluorophosphate ionic liquid, [hmim][FAP], as a solvent, on the solubility of these compounds in CO₂-rich phase, in ternary (CO₂ + 1,2-butanediol + [hmim][FAP]), CO₂ + butylene carbonate + [hmim][FAP] and quaternary (CO₂ + 1,2-butanediol + butylene carbonate + [hmim][FAP]) mixtures was investigated. The experimental results of the two binary systems were correlated using the density-based Chrastil equation. The knowledge of phase equilibria

* Corresponding author at: LAQV-REQUIMTE, Departamento de Química, Faculdade de Ciências e Tecnologia, Universidade NOVA de Lisboa, Quinta da Torre, 2829-516 Caparica, Portugal.

E-mail address: ma.zakrzewska@campus.fct.unl.pt (M.E. Zakrzewska).

<https://doi.org/10.1016/j.supflu.2023.106089>

Received 9 March 2023; Received in revised form 10 October 2023; Accepted 12 October 2023

Available online 13 October 2023

0896-8446/© 2023 The Authors. Published by Elsevier B.V. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

behaviours reported in this work will be useful for designing chemical conversions of carbon dioxide using [hmim][FAP] ionic liquid as reaction solvents.

1. Introduction

Increasing demand for biodegradable plastic materials revived the interest in aliphatic polycarbonates due to their biocompatibility and low toxicity [1,2]. Virtually ignored in the role of thermoplastics due to their poor thermal stability and high susceptibility to hydrolysis, aliphatic polycarbonates attracted increasing attention in biomedical fields. They can be degraded *in vivo* by surface erosion into neutral diols and carbon dioxide, in contrast to aliphatic polyesters, i.e., poly(lactic acid) – PLA, poly(lactic-co-glycolic acid) – PLGA, or poly(ϵ -caprolactone) – PCL, which upon decomposition give rise to diols and carboxylic acids leading to increased acidity of the surrounding tissue and inflammatory response, or affecting the loaded drug [3].

Aliphatic polycarbonates can be synthesised either by condensation polymerisation of dialkyl or diphenyl carbonate and aliphatic diols, copolymerisation of carbon dioxide with epoxides or by ring-opening polymerisation (ROP) of cyclic carbonates [4,5]. Cyclic carbonates themselves can be produced through a coupling reaction of carbon dioxide and cyclic ethers such as oxiranes or oxetanes, with alternative pathways including, i.e., carboxylation of olefins with carbon dioxide or

transformation of diols: transesterification with dialkyl carbonate/urea, oxidative carbonylation with carbon monoxide, or diols dehydrative condensation with carbon dioxide [6–8]. The last method is particularly relevant as it satisfies the requirements of green chemistry, such as lower production costs (single-step reaction) and eco-friendly properties (hazardless reagents and water as the sole by-product). Moreover, it involves carbon dioxide conversion, which gains further advantages in light of global concerns over greenhouse gas pollution and means to alleviate the shortage in conventional petroleum fuel supplies [9,10].

High-pressure CO₂ combined with an ionic liquid in a biphasic system has been proposed as an efficient and greener medium for many chemical processes [11–13], owing to the peculiar phase behaviour of (scCO₂+IL) mixtures first described by Brennecke and co-workers [14]. While the solubility of carbon dioxide in the ionic liquid-rich phase is significant, the molten salt does not dissolve in the pure carbon dioxide phase, even at high pressures. The discovery of this phenomenon led to a new method to separate ionic liquid from organic solvent [15–17], using CO₂ as a separation switch. As a further advance, high-pressure CO₂ allowed for an attractive integration of chemical reaction and separation unit operations by extracting the desired product from the ionic liquid




Fig. 1. Scheme of the experimental apparatus: 1 – CO₂ tank; 2 – ice bath; 3 – compressor; 4 – manual compressor; 5 – high-pressure cell; 6 – magnetic stirrer; 7 – magnetic stirring bar; 8 – temperature probe; 9 – HPLC valve; 10 – sampling loop; 11 – temperature controller; 12 – heating lamps; 13 – ventilator; 14 – ice bath; 15 – air bath; T – glass trap; F – calibrated expansion volume flask; VP – vacuum pump; P1 – P5 – pressure indicator; S1 and S2 – washing syringes.

phase, leaving behind not only ionic liquid but also the catalyst to reuse, permitting thus a continuous mode of operation.

In this context, in our ongoing project, we explore the possibility of synthesis of cyclic carbonates through a direct coupling of CO₂ and potentially biomass-derived alcohols in a biphasic system comprising one of the most CO₂-philic ionic liquid with physical absorption mechanism – 1-hexyl-3-methylimidazolium tris(pentafluoroethyl)trifluorophosphate, [hmim][FAP] [18,19]. This work discusses the influence of the ionic liquid on the solubility of cyclic carbonate and diol in the CO₂-dense phase. Here, the phase behaviours of binary, ternary, and quaternary mixtures containing dense carbon dioxide, [hmim][FAP], 1,2-butanediol (1,2-BDO) and/or corresponding butylene carbonate (BC) are reported at 313.2 K and pressures between 6 and 18 MPa. High-pressure fluid-phase equilibrium studies of (scCO₂ + IL) systems continue to play a secondary role in the undergoing research on carbon dioxide - ionic liquids interactions, and the data is limited, particularly for multicomponent mixtures [20,21]. To the best of our knowledge, the high-pressure solubility data of these organic compounds in (CO₂ + IL) multicomponent systems has not been reported in the literature.

2. Materials and methods

2.1. Materials

1,2-butanediol and butylene carbonate (4-ethyl-1,3-dioxolan-2-one) (≥ 98.0 % purity) were purchased from Fluka Analytical and Fluorochem, respectively, and dried over molecular sieves (Fluka Analytical, 0.3 nm pore diameter) for at least 24 h prior to their use. The ionic liquid, 1-hexyl-3-methylimidazolium tris(pentafluoroethyl)trifluorophosphate, [hmim][FAP], was purchased from Merck KGaA, Germany, (≥ 99.0 % purity) and used without further purification (water content < 300 ppm as confirmed with Karl Fischer titration method). Carbon dioxide (99,998 % purity) was supplied by Air Liquid and also used as received. Absolute anhydrous ethanol of 99.9 % purity (Carlo Erba, Barcelona, Spain) was used as a solvent to collect the vapour samples and for gas chromatography analysis. The following chemicals: hexadecane (Sigma-Aldrich, ≥ 99.0 % purity), n-nonane (Sigma-Aldrich, ≥ 99.0 % purity) and 1-octanol (Alfa Aesar, ≥ 99.0 % purity) were used as internal standards, depending on the systems analysed.

2.2. Apparatus and experimental procedure for phase behaviour studies

The phase behaviour investigation was carried out in a high-pressure apparatus described for the first time in 1993 [22], with the most current version including all the posterior modifications reported, in detail, in the work of Rebocho et al. [23]. Briefly, the experimental system is built around a 30 cm³ high-pressure cylinder-shaped sapphire cell placed inside a thermostatic air bath with a high-power ventilator for uniform heat distribution (Fig. 1). A transparent window allows for real-time visual surveillance of phase behaviour, i.e., for an inspection of the number of coexisting phases. The temperature is regulated by the Hart Scientific temperature controller (model 2100) coupled with an RTD probe (model 2622) from the same supplier (uncertainty of 0.1 K), and the pressure is measured by the Setra pressure transducer (model 204/C204), with an accuracy of ± 0.007 MPa.

The experiment begins with removing of all gases from the apparatus using a vacuum pump (Edwards, model RV3). Then, a certain amount of liquid sample (approximately 4 cm³ for each system) of known composition (weight was determined using KERN ABT 220-5DNM balance (uncertainty of 0.01 mg) and reported in Table S2 of the Supplementary Material) is loaded into the cell and heated up to the experimental temperature. Overall mole ratios of components investigated were kept constant and equalled to 0.3 for [hmim][FAP]:1,2-butanediol or [hmim][FAP]:butylene carbonate, and 1 for 1,2-

butanediol:butylene carbonate. These ratios were selected to mimic a reaction mixture with a 50 % conversion of diol to carbonate. When the temperature of 313.2 K is reached, the required CO₂ pressure is slowly introduced into the reactor using a CO₂ compressor. To accelerate the equilibration of liquid and vapour phases, the entire content of the cell is mixed with a magnetic stirrer for 1.5 h and then left to settle for 30 min. The sample from the gas phase is taken into the previously evacuated sampling loop (0.1 cm³) and then expanded to a low-pressure calibrated volume (276.5 cm³) using an HPLC valve. The loop is linked to an expansion zone through a glass trap immersed in an ice bath, ensuring precipitation of all components except CO₂. Next, a solvent (ethanol) is pumped through the tubing leading from the HPLC valve to the glass trap in order to collect any solute that might have precipitated along the way to the trap. Due to the limitations of the apparatus regarding highly viscous mixtures (very small inner diameter of tubing), it was impossible to take representative samples from the liquid phase. Each data point reported is the average of three replicate sampling measurements. In order to avoid the disturbance of the equilibrium upon sampling, only a small amount of sample was collected as verified by the pressure drop in the cell always smaller than 0.4 MPa. The validity of the experimental method for the phase behaviour determination was confirmed by carrying out replicas of the assays at random conditions of composition and pressure. The average standard deviation of mole fractions of alcohol and cyclic carbonate in the CO₂-rich phase is ± 0.0002 .

The amount of CO₂ in each sample is calculated by examining the pressure change in the calibrated sampling volume. The average standard deviation of the mole fraction of carbon dioxide is ± 0.0002 . The concentration of organic compounds, 1,2-BDO and BC, was analysed by a gas chromatographer coupled with a flame ionisation detector (GC-FID), using ZB-1HT Inferno capillary column (15 m, 0.25 ID, 0.1 film). Helium served as the carrier gas and 1-octanol (for samples with 1,2-BDO), n-nonane (for samples with BC) and hexadecane (for samples with 1,2-BDO and BC) were used as internal standards.

2.3. Correlation of solubility of polar compounds in dense CO₂

The density-based model proposed by Chrastil [24] was used to correlate experimental equilibrium data of the binary (CO₂ + 1,2-butanediol and CO₂ + butylene carbonate) systems:

$$S = \rho^k \exp\left(\frac{a}{T} + b\right) \quad (1)$$

where S is the solubility of a solute in CO₂ (g L⁻¹), ρ is the density of CO₂ (g L⁻¹) and T is the temperature (K), while k , a and b are constants. For a constant temperature, Eq. (1) can be written as:

$$\ln(S) = k \ln(\rho) + c \quad (2)$$

where c is a constant. The density of carbon dioxide was taken from the literature [25]. Parameters k and c were obtained by linear regression.

Root mean square deviation (RMSD) was used to compare the experimental data and data predicted by the Chrastil equation, according to:

$$RMSD = \sqrt{\frac{\sum_i (S_i^{calc} - S_i^{exp})^2}{N}} \quad (3)$$

where S_i^{exp} and S_i^{calc} are the experimental and calculated values of the solubility of either 1,2-butanediol or butylene carbonate in the vapour phase, respectively, while N is the number of data points.

3. Results and discussion

Table S1 (Supplementary Material) shows the experimental phase equilibrium data for the vapour phase of binary (CO₂ + 1,2-butanediol) and CO₂ + (butylene carbonate), ternary (CO₂ + 1,2-butanediol +




Fig. 2. Solubility data in dense CO₂ for 1,2-butanediol (filled circles – this work, empty circles – data from the literature [26]), butylene carbonate (filled squares–this work) and propylene carbonate (empty squares– data from literature [27]) at 313.2 K, expressed in mole fraction of solute in the gas phase.

[hmim][FAP]) and (CO₂ + butylene carbonate + [hmim][FAP]) and quaternary (CO₂ + 1,2-butanediol + butylene carbonate + [hmim][FAP]) at 313.2 K. These data were used to construct Figs. 2 and 3. Fig. 2 presents a vapour-liquid equilibrium diagram for the two binary systems and their comparison with the previously reported equilibrium data for the (CO₂ + 1,2-butanediol) [26] and (CO₂ + propylene carbonate) [27] systems, showing a relatively good agreement. The observed discrepancies in solubility values could be due to the difference in the methods

used – here, the static sampling method was used, while Inomata et al. [26] employed a flow flow-type apparatus. The authors have noted that their results are characterised by small discrepancies compared to a static method, particularly at lower pressures.

The solubilities of 1,2-butanediol and butylene carbonate in the vapour phase are similar at lower pressures. At elevated pressures, the solubility of butylene carbonate in the CO₂ phase is enhanced, probably due to the π - π interactions present in the carbonyl group, which forms a more stable complex [28]. Additionally, the affinity between CO₂ and carbonate is reflected in a high solubility of CO₂ in the butylene carbonate, leading to a significant expansion of the liquid phase up to the point of creating one phase at pressures exceeding 15 MPa. The strong hydrogen bonds between the hydroxyl groups of diol in the liquid phase as well as weaker CO₂-diol interactions might be responsible for the lower solubility of the alcohol in the CO₂ phase.

The Chrastil equation (Eq. (2)) was used to correlate the experimental solubility data of the two binary systems. Table 1 presents obtained constants (k and c) and root mean square deviation ($RMSD$), obtained from Eqs. (2) and (3). Figs. S1 (a) and S1 (b) presented in Supplementary Material show the fitting results.

Fig. 3(a) shows the solubility of 1,2-butanediol and butylene carbonate in the vapour phase of their ternary mixtures with CO₂ and [hmim][FAP] as a function of pressure at 313.2 K. We did not detect the presence of [hmim][FAP] in the vapour phase in all experiments. Very similar solubilities for the two ternary systems were observed in the whole pressure range. For comparison, Fig. 3(a) also contains data

Table 1

Summary of the parameters of Chrastil Eq. (1) fitted to the experimental solubility data for binary (CO₂ + 1,2-butanediol) and (CO₂ + butylene carbonate) system. The data are provided in Table S1 (Supplementary Material). $RMSD$ stands for the root mean square deviation.

	CO ₂ + 1,2-butanediol	CO ₂ + butylene carbonate
k	0.560	4.412
c	−8.77	−32.62
$RMSD$ (g L ^{−1})	0.0004	0.0075




Fig. 3. (a) Solubility of solutes (1,2-butanediol or butylene carbonate) as a function of pressure at 313.2 K in the vapour phase of the following systems: CO₂ + 1,2-butanediol (filled squares), CO₂ + 1,2-butanediol + [hmim][FAP] (empty squares), CO₂ + butylene carbonate (filled circles) and CO₂ + butylene carbonate + [hmim][FAP] (empty circles), in mole fraction (y_i). Overall mole ratios [hmim][FAP]:1,2-butanediol and [hmim][FAP]:butylene carbonate were kept at 0.3. (b) Solubility of 1,2-butanediol (full circles) and butylene carbonate (empty circles) as a function of pressure at 313.2 K in the vapour phase of the quaternary system composed of CO₂ + 1,2-butanediol + butylene carbonate + [hmim][FAP], in mole fraction (y_i). Overall mole ratios were: [hmim][FAP]:1,2-butanediol = 0.3, [hmim][FAP]:butylene carbonate = 0.3, and 1,2-butanediol:butylene carbonate = 1.

points for the solubilities in the corresponding binary systems. It can be observed that the solubility of 1,2-butanediol in the binary system is higher than in the corresponding ternary system at lower pressures but becomes very similar at higher pressures. A similar trend is observed when comparing the solubilities of butylene carbonate at low pressure. However, at higher pressures, the solubility of butylene carbonate is much lower in its ternary mixture than in binary, demonstrating the effect of the ionic liquid present in the liquid phase.

Finally, we studied the solubilities in the quaternary system composed of carbon dioxide, 1,2-butanediol, butylene carbonate and [hmim][FAP] and these results are presented in Fig. 3(b). In this system, the overall mole ratios [hmim][FAP]:1,2-butanediol and [hmim][FAP]:butylene carbonate were kept at 0.3, the same as in their corresponding ternary systems (Fig. 3(a)) in order to allow comparisons. The solubility of both solutes in the vapour phase was enhanced by increasing the pressure until reaching the maximum at 16 MPa, while a further increase in pressure resulted in lower solubility. The maximum solubility was 0.0043 and 0.0057 for butylene carbonate and 1,2-butanediol, respectively. These values are similar to their maximum solubilities observed in ternary systems (0.0054 and 0.0060). These results highlight the significant effect that the presence of an ionic liquid exerts on solute solubilities in dense CO₂, which, in this case, hinders the utilisation of CO₂ for an effective separation between the diol and the carbonate at the studied conditions.

4. Conclusions

The results show that the solubility of both organic compounds, 1,2-butanediol and butylene carbonate, in dense CO₂, in binary systems, increases with the increasing pressure in the entire measured range, with the effect being more pronounced in the case of cyclic molecule with carbonyl moiety. The strong affinity between CO₂ and butylene carbonate is reflected in a high solubility of CO₂ in the butylene carbonate, leading to a significant expansion of the liquid phase up to the point of creating one phase at pressures exceeding 15 MPa. For both organic compounds, the presence of fluorinated ionic liquid in the ternary systems leads to a decrease of the solubility values to almost to the same level within the whole pressure range studied. When present in the quaternary (CO₂ + 1,2-butanediol + butylene carbonate + [hmim][FAP]) mixture, at conditions of the example given, both organic molecules show increased solubility in the condensed phase, but only to approximately 16 MPa of pressure, after which their solubility starts to synchronically decrease.

Declaration of Competing Interest

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

Data availability

Data will be made available on request.

Acknowledgements

This work was supported by Fundação para a Ciência e Tecnologia FCT/MCTES projects PTDC/EQU-EPQ/31926/2017, UIDB/50006/2020, and MIT-EXPL/CS/0052/2021 of the Associate Laboratory for Green Chemistry – LAQV, and UIDB/00100/2020 of the Centro de Química Estrutural – CQE. GC-FID data were obtained by the Analysis Laboratory LAQV/Requimte - Chemistry Department, FCT NOVA - Portugal.

Appendix A. Supporting information

Supplementary data associated with this article can be found in the online version at doi:10.1016/j.supflu.2023.106089.

References

- [1] J. Xu, E. Feng, J. Song, Renaissance of aliphatic polycarbonates: new techniques and biomedical applications, *J. Appl. Polym. Sci.* 131 (2014) 39822, <https://doi.org/10.1002/app.39822>.
- [2] Y. Dai, X. Zhang, Recent development of functional aliphatic polycarbonates in the construction of amphiphilic polymers, *Polym. Chem.* 8 (2017) 7429–7437, <https://doi.org/10.1039/C7PY01815K>.
- [3] T. Artham, M. Doble, Biodegradation of aliphatic and aromatic polycarbonates, *Macromol. Biosci.* 8 (2008) 14–24, <https://doi.org/10.1002/mabi.200700106>.
- [4] X. Zhang, M. Fevre, G.O. Jones, R.M. Waymouth, Catalysis as an enabling science for sustainable polymers, *Chem. Rev.* 118 (2018) 839–885, <https://doi.org/10.1021/acs.chemrev.7b00329>.
- [5] W. Yu, E. Maynard, V. Chiaradia, M.C. Arno, A.P. Dove, Aliphatic polycarbonates from cyclic carbonate monomers and their application as biomaterials, *Chem. Rev.* 121 (2021) 10865–10907, <https://doi.org/10.1021/acs.chemrev.0c00883>.
- [6] P.P. Pescarmona, Cyclic carbonates synthesised from CO₂: applications, challenges and recent research trends, *Curr. Opin. Green. Sustain. Chem.* 29 (2021), 100457, <https://doi.org/10.1016/j.cogsc.2021.100457>.
- [7] E.J.C. Lopes, A.P.C. Ribeiro, L.M.D.R.S. Martins, New trends in the conversion of CO₂ to cyclic carbonates, *Catalysts* 5 (2020) 479, <https://doi.org/10.3390/catal10050479>.
- [8] K. Tomishige, Y. Gu, Y. Nakagawa, M. Tamura, Reaction of CO₂ with alcohols to linear-, cyclic-, and poly-carbonates using CeO₂-based catalysts, *Front. Energy Res.* 8 (2020) 1–24, <https://doi.org/10.3389/fenrg.2020.00117>.
- [9] A.J. Kamphuis, F. Picchioni, P.P. Pescarmona, CO₂-fixation into cyclic and polymeric carbonates: principles and applications, *Green. Chem.* 21 (2019) 406–448, <https://doi.org/10.1039/C8GC03086C>.
- [10] B. Grignard, S. Gennen, C. Jérôme, A.W. Kleij, C. Detrembleur, Advances in the use of CO₂ as a renewable feedstock for the synthesis of polymers, *Chem. Soc. Rev.* 48 (2019) 4466–4514, <https://doi.org/10.1039/C9CS00047J>.
- [11] Y. Medina-Gonzalez, S. Camy, J.-S. Condoret, ScCO₂/green solvents: biphasic promising systems for cleaner chemicals manufacturing, *ACS Sustain. Chem. Eng.* 2 (2014) 2623–2636, <https://doi.org/10.1021/sc5004314>.
- [12] E. García-Verdugo, B. Altava, M.I. Burguete, P. Lozano, S.V. Luis, Ionic liquids and continuous flow processes: a good marriage to design sustainable process, *Green. Chem.* 17 (2015) 2693–2713, <https://doi.org/10.1039/C4GC02388A>.
- [13] A.B. Paninho, A.L.R. Ventura, L.C. Branco, A.J.L. Pombeiro, M.F.C. Guedes da Silva, M. Nunes da Ponte, K.T. Mahmudov, A.V.M. Nunes, CO₂+ionic liquid biphasic system for reaction/product separation in the synthesis of cyclic carbonates, *J. Supercrit. Fluids* 132 (2018) 71–75, <https://doi.org/10.1016/j.supflu.2017.07.039>.
- [14] L.A. Blanchard, D. Hancu, E.J. Beckman, J.F. Brennecke, Green processing using ionic liquids and CO₂, *Nature* 399 (1999) 28–29, <https://doi.org/10.1038/19887>.
- [15] A.M. Scurto, S.N.V.K. Aki, J.F. Brennecke, CO₂ as a separation switch for ionic liquid/organic mixtures, *J. Am. Chem. Soc.* 124 (2002) 10276–10277, <https://doi.org/10.1021/ja0268682>.
- [16] V. Najdanovic-Visak, L.P. Rebelo, M. Nunes da Ponte, Liquid–liquid behaviour of ionic liquid–1-butanol–water and high pressure CO₂-induced phase changes, *Green. Chem.* 7 (2005) 443–450, <https://doi.org/10.1039/B500854A>.
- [17] V. Najdanovic-Visak, A. Serbanovic, J.M. Esperança, H.J. Guedes, L.P. Rebelo, M. Nunes da Ponte, Supercritical carbon dioxide-induced phase changes in (ionic liquid, water and ethanol mixture) solutions: application to biphasic catalysis, *Chem. Phys. Chem.* 4 (2003) 520–522, <https://doi.org/10.1002/cphc.200300663>.
- [18] M. Althuluth, M.T. Mota-Martinez, M.C. Kroon, C.J. Peters, Solubility of carbon dioxide in the ionic liquid 1-Ethyl-3-methylimidazolium tris(pentafluoroethyl)-trifluorophosphate, *J. Chem. Eng. Data* 57 (2012) 3422–3425, <https://doi.org/10.1021/jc300521y>.
- [19] M. Nunes da Ponte, M.E. Zakrzewska, Volumetric and phase behaviour of mixtures of fluoroalkylphosphate-based ionic liquids with high pressure carbon dioxide, *J. Supercrit. Fluids* 113 (2016) 61–65, <https://doi.org/10.1016/j.supflu.2016.03.017>.
- [20] J.M.S. Fonseca, R. Dohrn, S. Peper, High-pressure fluid-phase equilibria: experimental methods and systems investigated (2005–2008), *Fluid Phase Equilib.* 300 (2011) 1–69, <https://doi.org/10.1016/j.fluid.2010.09.017>.
- [21] S. Pepera, J.M.S. Fonseca, R. Dohrn, High-pressure fluid-phase equilibria: Trends, recent developments, and systems investigated (2009–2012), *Fluid Phase Equilib.* 484 (2019) 126–224, <https://doi.org/10.1016/j.fluid.2018.10.007>.
- [22] P.J. Pereira, M. Gonçalves, B. Coto, E.G. de Azevedo, M. Nunes da Ponte, Phase equilibria of CO₂ plus dl- α -tocopherol at temperatures from 292 K to 333 K and pressures up to 26 MPa, *Fluid Phase Equilib.* 91 (1993) 133–143, [https://doi.org/10.1016/0378-3812\(93\)85084-Y](https://doi.org/10.1016/0378-3812(93)85084-Y).
- [23] S. Rebocho, A.V.M. Nunes, V. Najdanovic-Visak, S. Barreiros, P. Simões, A. Paiva, High pressure vapor–liquid equilibrium for the ternary system ethanol/($\pm$)-menthol/carbon dioxide, *J. Supercrit. Fluids* 92 (2014) 282–287, <https://doi.org/10.1016/j.supflu.2014.06.006>.
- [24] J. Chrastil, Solubility of solids and liquids in supercritical gases, *J. Phys. Chem.* 86 (1982) 3016–3021, <https://doi.org/10.1021/j100212a041>.

- [25] R. Span, W. Wagner, A new equation of state for carbon dioxide covering the fluid region from the triple-point temperature to 1100 K at pressures up to 800 MPa, *J. Phys. Chem. Ref. Data* 25 (1996) 1509–1596, <https://doi.org/10.1063/1.555991>.
- [26] H. Inomata, A. Kondo, H. Kakehashi, Vapor–liquid equilibria for CO₂–fermentation alcohol mixtures: Application of a new group contribution equation of state to isomeric compounds, *Fluid Phase Equilib.* 228–229 (2005) 335–343, <https://doi.org/10.1016/j.fluid.2004.09.033>.
- [27] H. Li, R. Zhu, W. Xu, Y. Li, Y. Su, Y. Tian, Vapor-liquid equilibrium data of the carbon dioxide + ethyl butyrate and carbon dioxide + propylene carbonate systems at pressures from (1.00 to 13.00) MPa and temperatures from (313.0 to 373.0) K, *J. Chem. Eng. Data* 56 (2011) 1148–1157, <https://doi.org/10.1021/JE101086R>.
- [28] P.N. Khanh, N.T. Trung, Understanding Interaction Capacity of CO₂ with Organic Compounds at Molecular Level: A Theoretical Approach, in: I. Karamé, J. Shaya, H. Srou (Eds.), *Carbon Dioxide Chemistry, Capture and Oil Recovery*, IntechOpen, 2018. (<https://www.intechopen.com/chapters/58864>).