



RESEARCH PAPER



Microbial Occupational Exposure in e-waste Recycling: Biological Hazards, Their Sources, and Potential Toxic Effects

Carla Viegas^{1,2} · Bianca Gomes^{1,3} · Marta Dias^{1,2} · Renata Cervantes^{1,2} · Pedro Pena^{1,2} · Sara Gonçalves¹ · Marina Almeida-Silva^{1,4} · Elisabete Carolino¹ · Magdalena Twarużek⁵ · Liliana Aranha Caetano^{1,6} · Susana Viegas^{1,2}

Received: 31 January 2024 / Revised: 3 September 2024 / Accepted: 5 October 2024
© The Author(s) 2024

Abstract

E-waste management is a major environmental concern that also poses serious risks to occupational health. While e-waste workers are exposed to a variety of chemicals, little is known about occupational exposure to microbiological agents. This study aims to fill that gap. For that purpose, two e-waste facilities were assessed: one battery-recycling facility (BRF), and another facility (SRE) dedicated to e-waste sorting and storage before sending it for different processing. Filtering respiratory protection devices (FRPD, 24 samples) and settled dust (13 samples) were collected using standardized sampling approaches, and analyzed by culture-based, chemical and molecular methods. Workers also filled in a questionnaire regarding work activities and the use of FRPD. Microbial contamination, fungal diversity, azole resistance, and prevalence of *Aspergillus* sections *Fumigati* and *Flavi* (as surrogates of harmful fungal contamination) and mycotoxins were determined. To estimate potential health effects related to exposure, the FRPD cytotoxicity was determined in human A549 lung and HepG2 hepatic cells. Microbial contaminants with pathogenicity and toxigenic potential were observed in FRPD and settled dust. *Aspergillus* section *Fumigati* was widespread. The observed presence of *Aspergillus* section *Flavi* relates to the risk of exposure to aflatoxin B1 (a potent hepatocarcinogen). Four different mycotoxins were detected in 8% FRPD and 15% dust samples, including fumonisin B1 and sterigmatocystin (below the limit of quantification). A low to moderate cytotoxic effect of sampled FRPD was observed, with human lung alveolar cells more sensitive than hepatic cells. No relevant azole resistance was observed, nor correlations between fungal reduced azole-susceptibility and fungal growth at 37 °C or cytotoxicity. In summary, it can be concluded that this approach using FRPD and settled dust presents a promising potential for screening occupational exposure to microbiological contaminants in e-waste management. More studies in this occupational context should be prioritized to identify critical biological hazards and to support the implementation of appropriate health risk mitigation strategies.

Highlights

- The FRPD should be used to assess occupational exposure to biologic agents in e-waste units
- Fungi with pathogenic and toxigenic potential were found in FRPD and settled dust
- *Aspergillus* section *Fumigati* was widespread in the analysed e-waste units
- *Aspergillus* section *Flavi* results suggest potential workers exposure to AFB1
- Exposure to mycotoxins might occur through FRPD since a daily replacement is not ensured

Keywords Filtering respiratory protection devices · E-waste · Microbial occupational exposure · Cytotoxicity · *Aspergillus* section *Fumigati*

Extended author information available on the last page of the article

Introduction

E-waste (electronic and electrical waste) is one of the fastest-growing waste flows in the European Union (EU) and is a major environmental concern, with less than 40% of e-waste being recycled (European Parliament 2023). The waste management sector, and more precisely the industries devoted to sorting waste are of utmost importance for the Sustainable Development Goals (SDGs) achievement, since they contribute directly and indirectly to many of the SDGs. The European Commission (EC) adopted the new circular economy action plan (CEAP) in March 2020, one of the main building blocks of the European Green Deal, Europe's new agenda for sustainable growth. The EU's transition to a circular economy will reduce pressure on natural resources and will create sustainable growth and jobs. It is also a prerequisite to achieve the EU's 2050 climate neutrality target and to halt biodiversity loss (EC 2023). To accomplish these goals, the number of units dedicated to waste management and the corresponding workforce is anticipated to increase in all the EU countries and partners. One of the EC's priorities is the reduction of e-waste, although few EU countries recycle more than 50%. In 2016, Portugal reported a recycling rate of 43.5% for e-waste (European Parliament 2023). The same trends in production and recycling are observed in other parts of the world (UNITAR, 2024).

It is well-known that e-waste workers are exposed to a variety of chemicals including metals, particulates, persistent organic compounds, and flame retardants (Gravel et al. 2023; ILO, 2021; Okeme and Arrandale 2019; Scheepers et al. 2021). Regarding exposure routes, workers involved in e-waste dismantling activities may be exposed to inorganic and organic contaminants through dermal contact, inhalation, and ingestion (via hand-mouth contact). Most studies assessing external chemical exposures during e-waste recycling have used air sampling and dust or soil (Akram et al. 2019; Scheepers et al. 2021).

Exposure to microbiological agents during regular waste sorting activities has been reported in different countries (Degois et al. 2017; Eriksen et al. 2022; Lavoie et al. 2006; Madsen et al. 2016; Park et al. 2011; Salamanga et al. 2022; Viegas et al. 2022a, 2022b; Wouters et al. 2005). In Portugal, assessments of occupational exposure in waste management determined the prevalence of microbiological hazards such as toxigenic fungi (Viegas et al. 2022a, 2022b, 2017; Viegas et al. 2015a, b), azole-resistance (Gonçalves et al. 2021; Viegas et al. 2022a) and mycotoxins (Viegas et al. 2015a, b; Viegas et al. 2014, 2018). However, little is known regarding microbial occupational exposure in e-waste recycling.

Since Portuguese legislation comprises the mandatory use of filtering respiratory protection devices (FRPDs) in

waste sorting facilities, a previous study has used samples of used FRPD to assess microbial occupational exposures in this setting (Viegas et al. 2021a, 2021b, 2020a, b, c). It was found that workers in waste-sorting and belt-feeding activities are more exposed to microbial contaminants and that samples of used FRPD led to similar results compared to air sampling, with higher microbial contamination and cytotoxicity levels observed in the interior layers of FRPD (compared to exhalation valves). Of utmost importance, FRPD-recovered azole-resistant *Aspergillus fumigatus* isolates were found to be harboring the TR34/L98H mutation. Settled dust sampling has also been used to evaluate microbiological exposures in the waste sorting sector, offering significant outcomes for risk assessment and management (Salamanga et al. 2022; Scheepers et al. 2021; Viegas et al. 2022a). These findings provide valuable insights into the importance and methodology to follow when conducting microbial exposure assessments in the waste sorting industry, and in e-waste sorting facilities.

The present study aims to address the lack of information on occupational exposure to microbial contamination in e-waste sorting facilities using novel sampling methods, such as samples of used FRPD (besides settled dust), and their analysis by culture-based techniques and molecular methods. Additionally, it intends to determine possible deleterious health effects through used FRPD cytotoxicity analysis.

Materials and Methods

Study Site

This study was developed between September and November of 2021 in 2 units dedicated to e-waste management: one battery recycling facility (BRF) and one dedicated to sorting and storing different e-waste before sending it to further processing facilities (SRE). Both units are located in Portugal. Briefly, the BRF comprises different workstations aiming to separate the different battery components for dedicated recycling processes such as the plastics and heavy metals components, namely: the ovens, logistics, crucibles, and maintenance area. Additionally, a battery-breaking device and crushing and casting machines are also present on the site to support the recycling process. Regarding the SRE unit, manual sorting and storage of different electric and electronic devices are the main activity and comprises several storage areas to facilitate the distribution for other facilities to continue the specific recycling process.

Samples Collection

Thirteen samples of settled dust were collected (3 g from each sampling point) from the two units. The selected sampling sites were the workstations belonging to the different units where workers spend more time. In the BRF, settled dust samples were collected from nine distributed "indoor" and "outdoor" sites (furnaces, casting machine, crucibles, stacker, furnace stacker, plastic, beater, shovel, and maintenance area). In the SRE, four outdoor sites (storage 1,2,3, and storage shovel) were selected. Regarding FRPD, 24 samples identified as FFP3 (based on EN 149 classification on filtering efficiency) were collected from the BRF and the interior layers were analyzed. A non-used FRPD was used as a control. All samples were individually packed in sterilized bags, identified, and transported in refrigerated conditions (0–4 °C).

Questionnaire on FRPD Use

The study enrolled twenty workers who completed a questionnaire about their usage of FRPDs. The questionnaire included questions about the type of FRPD, workstation, hours of use, consecutive hours of use, frequency of replacement, experience of use (in years), FRPD storage location during breaks, and whether they noticed any odours while using FRPD. Four workers in the ovens area, three workers in crucibles, and three workers in the logistics area used the FRPD. One worker used their FRPD only when managing the crushing machine, while another used it only when operating the battery-breaking machine. One worker did not provide information about the corresponding workstation. Additionally, the participants rated on a scale of 1–5, the amount of heat and sweat they felt while wearing the FRPD, and how well it adjusted their face. This study followed the guidelines of the Helsinki Declaration and Oviedo Convention and all data was stored and analyzed in accordance with the Portuguese General Data Protection Regulation (GDPR) law n° 58/2019. Workers were invited to participate voluntarily in the project, and before enrolment, all volunteers filled out a written informed consent.

Microbial Contamination

FRPDs were collected after being used during a work shift and a 2 cm² section of the interior layer was extracted. using 10 ml of PCR grade water, subjected to agitation for 30 min at 2500 rpm on an orbital laboratory shaker as previously reported (Viegas et al. 2020a). The extracted material was spread on selective culture media for fungi, (Malt extract agar (MEA) supplemented with chloramphenicol (0.05%) and dichloran-glycerol agar (DG18) incubated at 27 °C and at 37 °C for potential pathogenic evaluation) and bacteria

(tryptic soy agar (TSA) supplemented with nystatin (0.2%), and Violet Red bile agar (VRBA) incubated at 30 °C and 37 °C, respectively) as well as in azole-supplemented Sabouraud agar media (4.0 mg/L itraconazole (ITZ), 2.0 mg/L voriconazole (VCZ), and 0.5 mg/L posaconazole (PCZ)) as previously reported (Viegas et al. 2020b). Fungal species were identified by an environmental and occupational mycologist using macro and microscopic features (De Hoog 2000).

The targeted *Aspergillus* sections *Fumigati* and *Flavi* were detected using the extracts obtained, following previously published procedures (Viegas et al. 2020a). Fungal DNA was extracted using the ZR Fungal/Bacterial DNA MiniPrep Kit (Zymo Research, Irvine, USA) as per the manufacturer's instructions. Fungal identification was achieved by Real-Time PCR (qPCR) using the CFX-Connect PCR System (Bio-Rad). For qPCR, reactions were prepared with 1 × iQ Supermix (Bio-Rad, Portugal), 0.5 μM of each primer, and 0.375 μM of TaqMan (Table S1– Supplementary material). A three-step reaction including 40 cycles of denaturation at 95 °C for 30 s, annealing at 52 °C for 30 s, and extension at 72 °C for 30 s. For each gene that was amplified, a non-template control and a positive control consisting of DNA obtained from reference strains (kindly provided by the Mycology laboratory of the National Institute of Health Dr. Ricardo Jorge) were used.

Mycotoxins Analysis

The method used to prepare samples and conduct chromatographic analysis of mycotoxins followed the procedure described in Viegas et al. (2020a). Samples of settled dust (0.10 g) and FRPD (0.05 g) were mixed with 3.0 ml and 2.5 ml respectively, of a mixture containing acetonitrile, water, and acetic acid (79/20/1; v/v/v) and shaken for 60 min. After centrifugation (5000 rpm for 5 min), the resulting samples were dried under nitrogen and dissolved in the mobile phase (A: B 7:3). Both mobile phase A (methanol/water/acetic acid, 10/89/1 (v/v/v)) and mobile phase B (methanol/water/acetic acid, 97/2/1 (v/v/v)) contained 5 mmol/L ammonium acetate. Separation was conducted on a Phenomenex Gemini C18 column (150 × 4.6 mm, 5 μm) (Torrance, CA, USA) with a flow rate of 1 ml/min and injection volume of 5 μL. The following gradient was used for mobile phase B: 0% up to 2', increasing to 50% from 2' to 5', increasing to 100% from 5' to 14', maintaining 100% up to 18', and returning to 0% by 22.5'.

Mycotoxins were detected using a Shimadzu HPLC Nexera (Tokyo, Japan) coupled with a Sciex mass spectrometry detector 5500 QTrap (Foster City, USA). The analysis was conducted in the scheduled multiple reaction monitoring (sMRM) mode for both negative and positive polarities within a single chromatographic run. The electrospray

ionization (ESI)-source parameters were set as follows: curtain gas at 30 psi, collision gas at medium level, ion spray voltage set at -4500 V (negative polarity) and 5500 V (positive polarity), ion source temperature maintained at 550 °C, ion source gas1 at 80 psi, and ion source gas2 at 80 psi.

Cytotoxicity Analyses

To assess the biological impact of the contaminants, present in the FRPD after use, an *in vitro* study was conducted using human lung epithelial cells (A549) and human liver carcinoma cells (HepG2). These cells were selected as they are relevant to the primary exposure routes protected by FRPD, namely, inhalation and ingestion. These cells were exposed to various concentrations of FRPD extracts and evaluated their viability using the 3-(4,5-dimethylthiazol-2-yl)-2,5-diphenyltetrazolium bromide (MTT) assay, which has been previously described (Hanelt et al. 1994; Viegas et al. 2023).

Cells were cultured using Eagle's Minimum Essential Medium (MEM) supplemented with 10,000 units of penicillin and 10 mg/mL streptomycin in 0.9% NaCl and fetal bovine serum (Sigma-Aldrich, USA). The cells were detached using 0.25% (w/v) 0.53 mM Trypsin EDTA, and then 100 μ L of cell suspensions (3.0×10^5 cells/mL A549 cells; 2.0×10^5 cells/mL HepG2 cells) were serially diluted to 96-well plates. These plates were incubated with FRPD extracts for 48 h at 5% CO₂, 37 °C in a humidified atmosphere. The cellular viability was determined by MTT assay at 510 nm (using the LEDETECT 96 ELISA and MikroWin2013SC software). The threshold toxicity was considered as the lowest concentration decreasing absorption to less than 50% of cellular metabolic activity (IC50).

Statistical Analysis

The Spearman's correlation coefficient was used to evaluate the relationship among contamination variables (prevalence's of bacteria, fungi, azole resistance, *Aspergillus* sp., cytotoxicity) and information regarding the FRPD use. The Mann–Whitney *U* test was used to compare contamination variables among sampling matrices (FRPD and settled dust), and in settled dust among different units. The Simpson and Shannon indices [given by Shannon Index ($H = -\sum_{i=1}^s p_i \ln(p_i)$) and Simpson Index ($D = \frac{1}{\sum_{i=1}^s p_i^2}$), where p_i is the proportion (n_i/n) of individuals of one particular species found (n_i) divided by the total number of individuals found (n)], were used to assess fungal species diversity. All analyses were conducted in SPSS statistical software for Windows, version 27.0. The results were considered significant at the 5% significance level.

Results

Patterns of FRPD Use

The questionnaire showed that 35% of workers ($n = 7$) use the same FRPD for all workstations and tasks without changing it. Out of the 20 participants surveyed, 18 of them (90%) reported that they used the FRPD for seven hours, or five consecutive hours. Additionally, 8 participants (40%) mentioned they had used the same FRPD for three years. One participant's experience varied depending on their seniority in the plant, with some indicating 30, 32, 33, or 36 years of experience. As for the discomfort caused by the FRPD, 10 participants (50%) reported feeling no more discomfort than usual, while two (10%) felt more discomfort than usual. A small percentage of participants (15%; 3 out of 20) reported that the FRPD caused more sweating than usual. However, all participants agreed that the FRPD was easily adjustable to fit their face.

Microbial Exposure

Total bacterial concentrations (cultivated on TSA agar) were higher in workers' FRPD from the ovens area (6.20×10^4 CFU.m⁻²), and Gram-negative bacterial concentrations (cultivated on VRBA agar) were higher in workers' FRPD from all areas (1.05×10^4 CFU.m⁻²). Bacterial prevalence was higher in FRPD than in settled dust (Fig. 1). Based on settled dust results, the highest total bacteria counts were found near the battery crushing machine from the RBF (TSA: 1.77×10^2 CFU.g⁻¹), while the Gram-negative bacteria highest counts were obtained in the SRE (VRBA: 2.56×10^2 CFU.g⁻¹).

The highest fungal concentrations were found in DG18 (after 5 days incubation at 27 °C), being 1.16×10^5 CFU.



Fig. 1 Bacteria counts from used filter respiratory protective device (FRPD) (log [CFU]. m⁻²) and settled dust samples (log [CFU].g⁻¹) inoculated in TSA and VRBA culture media at 30 °C and 37 °C respectively

m⁻² FRPD (from the battery-breaking machine area) and 2.77 × 10² CFU.g⁻¹ settled dust. Fungal concentrations on MEA were 1 log lower in both used FRPD and settled dust. DG18 plates incubated at 37 °C (to evaluate exposure to potentially pathogenic fungi) reached 4.00 × 10³ CFU.m⁻² FRPD and 2.60 × 10¹ CFU.g⁻¹ settled dust (Fig. 2). The highest fungal counts, among settled dust samples from both units, were obtained in SRE (MEA: 3.9 × 10¹ CFU.g⁻¹; DG18: 2.77 × 10² CFU.g⁻¹).

Regarding fungal variability, *Penicillium* sp. was dominant in settled dust (42.15% samples) and FRPD (74.27% samples) on DG18. *Cladosporium* sp. was dominant in used FRPD (47.52% samples) on MEA. *Aspergillus* sp. was also present in settled dust (MEA: 8.77%) and in used FRPD (DG18: 8.05%) (Table 1; Figure S1–Supplementary material).

Based on FRPD results, the greatest fungal biodiversity on MEA was found in the ovens workstation (Shannon index (H) = 1.11, Simpson Index (D) = 9.39), followed by crucibles (Shannon index (H) = 1.61, Simpson Index (D) = 5.00), whereas on DG18 it was in all the workstations




Fig. 2 Fungal counts obtained from used filter respiratory protective device (FRPD) (log [CFU]. m⁻²) and settled dust samples (log [CFU].g⁻¹) inoculated in MEA and DG18 at 27 °C for environmental purposes and in DG18 at 37 °C for potential pathogenic evaluation

Table 1 Fungal distribution (%) on MEA and DG18 from used FRPD and settled dust samples

Samples	MEA			DG18		
	Species	CFU.m ⁻² /g ⁻¹	%	Species	CFU.m ⁻² /g ⁻¹	%
FRPD	<i>Cladosporium</i> sp.	6.70X10 ⁴	47.52	<i>Penicillium</i> sp.	1.66X10 ⁵	74.27
	<i>Penicillium</i> sp.	6.40X10 ⁴	45.39	<i>Cladosporium</i> sp.	3.85X10 ⁴	17.23
	<i>Aspergillus</i> sp.	4.00X10 ³	2.84	<i>Aspergillus</i> sp.	1.80X10 ⁴	8.05
	Others species	6.00X10 ³	4.26	Others species	1.00X10 ³	0.45
	TOTAL	141X10 ⁵	7.09	TOTAL	2.24X10 ⁵	100.00
Settled dust	<i>Penicillium</i> sp.	2.20X10 ¹	19.30	<i>Penicillium</i> sp.	3.73X10 ²	42.15
	<i>Cladosporium</i> sp.	2.00X10 ¹	17.54	<i>Cladosporium</i> sp.	2.35X10 ²	26.55
	<i>Aspergillus</i> sp.	1.00X10 ¹	8.77	<i>Aspergillus</i> sp.	5.00X10 ¹	0.56
	Others species	6.20X10 ¹	54.39	Others species	2.72X10 ²	30.73
	TOTAL	1.14X10 ²	100.00	TOTAL	8.85X10 ²	100.00

(All) (Shannon index (H) = 1.24, Simpson Index (D) = 3.05). Regarding settled dust, the greatest fungal biodiversity was observed in the sorting plastic area both for MEA (Shannon index (H) = 0.92, Simpson Index (D) = 2.23) and DG18 (Shannon index (H) = 0.95, Simpson Index (D) = 14.42) (Tables S2 and S3 – Supplementary material).

An overview of the distribution of *Aspergillus* sections can be found in Fig. 3. *Aspergillus* diversity was higher in settled dust inoculated on DG18 at 37 °C, with 5 sections identified, namely, *Nigri* (51.43%), *Fumigati* (44.29%), *Nidulantes* (1.43%), *Flavi* (1.43%) and *Aspergilli* (1.43%). At 27 °C, three *Aspergillus* sections were identified in DG18 (40% *Circumdati*; 40% *Nigri*; 20% *Flavi*), and two sections in MEA (80% *Nigri*; 20% *Nidulantes*). Regarding FRPD, four sections were found in MEA (50% *Nidulantes*; 25% *Nigri*; 12.50% *Fumigati*; 12.50% *Circumdati*), and three sections in DG18 (61.11% *Nidulantes*; 30.56% *Circumdati*; 8.33% *Aspergilli*). In FRPD inoculated on DG18 and incubated at 37 °C, *Aspergillus* section *Fumigati* was dominant (52.00%), followed by *Nigri* (16.00%) and *Nidulantes* (12.00%) sections (Fig. 3).

Screening of Antifungal Resistance

No fungal growth on azole-supplemented media was observed for used FRPD samples. Regarding settled dust, in one sample only, *Mucor* sp. (CFU.m⁻²/g⁻¹) with reduced susceptibility to 2.0 mg/L voriconazole and 0.5 mg/L posaconazole was identified in the storage unit for recycling electric and electronic devices (SRE).

Aspergillus Section *Fumigati* and Section *Flavi* Detection

From the two targeted *Aspergillus* sections, *Aspergillus* section *Fumigati* was detected in 9 out of 24 used FRPD samples (37.5%) from the RBF. Concerning *Aspergillus* section *Flavi* it was detected in 9 out of 24 used FRPD (37.5%) and




Fig. 3 *Aspergillus* sections' distribution in MEA, DG18, and DG18 (37 °C) in used FRPD and settled dust samples

in 3 out of 9 settled dust samples from the BRF, whereas from SRE it was detected in 2 out of 4 settled dust samples (50%). In 5 used FRPD both sections were detected. In 15 out of 33 samples (45.5%), section detection was obtained, while the identification was not possible to achieve by culture-based methods (Supplementary material—Table S4).

Mycotoxin Exposure

Two used FRPD (used more than 5 consecutive hours and without daily replacement) were contaminated with mycotoxins: one FRPD with fumonisin B2 below the LOQ (9 ng/g); another FRPD with fumonisin B1 (<LOQ, 12 ng/g), fumonisin B2 (9.5 ng/g), and sterigmatocystin (<LOQ, 4 ng/g). Also, two settled dust samples (from SRE) were contaminated, including one sample with mycophenolic acid (26 ng/g) and sterigmatocystin (< 6 ng/g – LOQ).

FRPD Cytotoxicity

All used FRPD samples with absorbance values less than 50% of the cell division activity were considered toxic. A semi-quantitative scale was used to grade cytotoxicity. A low cytotoxic effect was observed for IC50 values ranging from 31.251 to 7.813 cm²/mL, a moderate cytotoxic effect for values ranging from 3.906 to 0.977 cm²/mL, and a high cytotoxic effect for values ranging from 0.488 to 0.061 cm²/mL. The absence of cytotoxicity was determined when the absorbance value was ≥ 50 for

the first dilution tested (31.5 cm²/mL). Tables 2 and 3 show the results of the MTT cytotoxicity test for FRPD extracts. The analyzed FRPD showed cytotoxicity in 16 cases (66.7%) in A549 cells, with only 8 FRPD showing no cytotoxicity. In HepG2 cells, only 16.7% of the FRPD studied AQ3 exhibited cytotoxicity, all demonstrating low levels of cytotoxicity. Among the 24 tested FRPD, 4

Table 2 Cytotoxicity of used FRPD on A549 cells

Sample	Level of cytotoxicity (low; moderate; high)	IC50	Dilution step
F1	Low	10 mm ² /mL	1
F2	Low	10 mm ² /mL	1
F3	Low	10 mm ² /mL	1
F3.1	Low	10 mm ² /mL	1
F5	Low	10 mm ² /mL	1
F6	Low	10 mm ² /mL	1
F9	Low	10 mm ² /mL	1
F10	Moderate	5 mm ² /mL	2
F11	Low	10 mm ² /mL	1
F12	Low	10 mm ² /mL	1
F14	Low	10 mm ² /mL	1
F15	Low	10 mm ² /mL	1
F16	Low	10 mm ² /mL	1
F17	Low	10 mm ² /mL	1
F19	Moderate	5 mm ² /mL	2
F20	Low	10 mm ² /mL	1




Fig. 4 Comparison of bacterial and fungal contamination, azole resistance and *Aspergillus* sp. counts between BRF and SRE. Mann–Whitney test results

The use of 37 °C as incubation temperature allowed the identification of the fungal species/sections that can grow at that temperature unveiling the pathogenic potential (Anderson et al. 2023). Additionally, it was also possible to overcome the limitation in recovering fungi with clinical relevance, such as *Aspergillus* section *Fumigati* isolates that, in this kind of setting, due to fungi with higher growth rates, such as Mucorales order, *Trichoderma* sp., and *Chrysonilia sitophila* are commonly underestimated. This is the most important drawback of using only culture based-methods to assess exposure to microbiologic agents in highly contaminated occupational environments (Cox et al. 2020; Salambanga et al. 2022; Viegas et al. 2023, 2020a), and by using selective temperatures (37 °C) and/or specific culture media we can better characterize the risk in the setting being assessed.

Microbial Contamination Findings

In this study, lower counts of bacteria were obtained in settled dust samples than in used FRPD extracts. Not only settled dust and FRPD are different sampling/collection methods they also have quite different lab protocols to achieve the relative extracts that can lead to different microbial counts and microbial diversity, as was observed in this study. Furthermore, higher microbial counts, and more specifically bacterial counts (TSA and VRBA), and *Aspergillus* section *Fumigati* isolates, found in used FRPD can be due to the interior layer favourable conditions (Cox 1989; Viegas et al. 2021a, 2021c, 2020a). In fact, during FRPD use, water vapour and sweat are released boosting material humidity,

and the presence of nutritional substances that were blocked during air filtering can provide optimal conditions for microorganism growth (Donlan 2002; Jankowska et al. 2000; Majchrzycka et al. 2016; Viegas et al. 2020a). Additionally, the settled dust from this specific industry is expected to have a high chemical burden (Okeme and Arrandale 2019) and this will impact negatively the microorganism's growth (Roane et al. 2009; Gomes et al. 2020) justifying the lower counts in this matrix when comparing with used FRPD and also with other studies performed in Portugal in similar settings (Viegas et al. 2022a).

In this study, the different culture media enabled a deeper characterization of the microbial counts in both units. Of note, it was possible to observe in settled dust samples that Gram-negative bacteria presented higher counts in SRE, while total bacteria counts were higher in BRF. This trend may have some seasonal influence, since all the workstations from SRE were outdoors, with expected lower temperatures and higher humidity at that time, favouring Gram-negative bacteria proliferation (Won and Ross 1966), besides fungal proliferation and dissemination (Nageen et al. 2023). Bacteria are also more sensitive to the assessment conditions (aerosolization, collection, and culture) impacting more in their viability (Cox 1998) and, consequently, leading to quite different results among settings.

In what concerns the mycobiota, SRE presented the highest fungal counts on the settled dust samples and this can be due to the outdoor environment where the workstations are located. DG18 unveils higher fungal counts in both matrices, being also observed in other different settings (Viegas et al. 2023). DG18 has limited water content, allowing a

more diversified number of species to thrive, by restricting the ones with fast growth rates (Mensah-Attipoe and Taubel 2017). Fungal species in general, and *Aspergillus* species/sections in particular, are sensitive to the occupational environment, sampling methods, and culture media (Chao et al. 2002; Viegas et al. 2021b), which restrains results' comparison among different studies. It has been previously reported that to assess occupational exposure to fungi in the waste sorting industry, used FRPD should be used as collection material and DG18 as the culture media whenever *Aspergillus* section *Fumigati* is targeted (Viegas et al. 2021c). In the present study assessing fungal contamination in e-waste recycling, *Aspergillus*' highest counts were also found in used FRPD using DG18, and the *Fumigati* section was also the most prevalent in used FRPD at 37 °C in the DG18. This trend was confirmed by previous studies that analyzed FRPD used by sorting waste workers (Viegas et al. 2020a, 2020b).

The workstations presenting higher contamination in this study were the ovens for total bacteria, maybe due to the increased temperatures expected in this workstation. The FRPD used in all areas were the ones presenting higher counts of gram-negative bacteria, while for fungi the higher counts were found in the FRPD used in the battery-breaking machine. FRPD used by workers in ovens and the ones used by workers in all the workstations (workers without a fixed workplace) were the ones presenting higher fungal diversity. Previously, it was reported that FRPD interior layers from workers with more waste contact, such as sorting the waste or feeding the belts with waste, showed an increased exposure to microbial contamination (Viegas et al. 2020a). Thus, the used FRPD extracts' analyses can mimic the expected occupational exposure to microbiologic agents and help prioritize interventions for risk management (Viegas et al. 2021c, 2020a, 2020b).

The settled dust samples didn't follow the same trend in the logistics presenting the highest counts in this workstation, and in the plastic area where higher fungal diversity was observed. This findings can be due to lower chemical pollution present in this workplace (Okeme and Arrandale 2019). Thus, the use of different sampling methods was useful in identifying the workstations where workers have an increased exposure to microbial contamination.

The importance of addressing fungal resistance is related to the emerging number of invasive fungal diseases worldwide. As only a few classes of antifungal drugs are available, the emergence of resistance to single and multiple drugs complicates fungal infections' management and patients' prognostics (Fisher et al. 2022; Global Action for Fungal Infections 2022). The recent WHO fungal priority pathogen list identifies 19 fungi with the greatest impact on public health and emerging antifungal resistance risk (WHO 2022). Although no relevant azole resistance was found in this study, *Mucor* sp. from one settled dust sample

presented reduced susceptibility to voriconazole and posaconazole at tested conditions. WHO ranks Mucorales as a high priority based on criteria such as average case fatality, number of new cases annually, complications and sequelae, and antifungal resistance, among other criteria. Mucormycetes are characterized by high-level resistance to most currently available antifungal drugs, particularly to short-tailed azoles such as voriconazole, due to an evolutionarily conserved amino acid substitution of the lanosterol 14 α -demethylase (Caramalho et al. 2017; Danaoui 2017). Standardized antifungal susceptibility testing methods and molecular detection at the species level will be further used to better characterize the antifungal susceptibility of the recovered *Mucor* isolate.

Applying culture-based methods only identifies culturable microorganisms. Furthermore, the majority of microorganisms present in a specific setting, cannot be cultured by routine laboratory assays (Cox et al. 2020; DeLong and Pace 2001), and several microorganisms, such as some pathogenic bacteria, can be in a viable but non-culturable state, but can still cause health outcomes (Li et al. 2014). Additionally, the non-viable, non-cultivable microorganisms can still be harmful to exposed workers, since several health outcomes are related to their toxigenic potential and not linked to viability or infectious potential (Cox et al. 2020). More recently, culture-based methods have gained more attractiveness due to culturomics, which combines culture-based methods with more refined tools (such as sequencing or matrix-assisted laser desorption/ionization-time of flight (MALDI-TOF)) for the identification of unidentified isolates (Cox et al. 2020). Microorganisms' viability is a crucial parameter since only viable microorganisms can lead to infection (Cox et al. 2020), justifying the culture based-methods also as the "gold standard" for occupational exposure assessments, besides clinical purposes (Kung et al. 2018). In this study, several culture media and incubation temperatures were applied together with qPCR detection of two different *Aspergillus* sections (*Fumigati* and *Flavi*). In light of previous studies (Viegas et al. 2020a, 2020b, 2020c, 2018; Viegas et al. 2015a, b), the *Fumigati* section was suggested as a surrogate of harmful fungal contamination in waste management industry (Viegas et al. 2022a) and in the WHO fungal priority list to guide research, development and public health action, listed in the critical priority group due to clinical relevance, antifungal resistance, and health outcomes (WHO, 2022). Thus, it is recommended that occupational exposure to this specific section should be reduced to a minimum (Marchand et al. 2021, 2017; Viegas et al. 2022a). In this study, we can conclude that this section is widespread by analyzing results from culture-dependent and by qPCR, claiming attention for his dominance at 37 °C, in both matrices.

Concerning the *Flavi* section, the concern relies mainly on their ability to produce aflatoxins, although the strains belonging to this section can also cause infections. IARC Monographs reported aflatoxins carcinogenic to humans (Group 1) (IARC, 2002). In this study, section *Flavi* was detected in both matrices, although not identified by culture-based methods in used FRPD. Formerly, in a study intending to assess exposure to aflatoxin B1 in workers sorting waste, through biomonitoring, all the workers participating in the study presented detectable levels of the targeted mycotoxin in blood samples (Viegas et al. 2014). Thus, in light of the results obtained from this section in the assessed units, the exposure to AFB1 should be considered. For all the above reasons, both culture-based and culture-independent methods should be applied to obtain a more accurate image of microbial contamination in occupational settings (Cox et al. 2020; Viegas et al. 2023, 2021a, 2021c, 2020a).

Mycotoxins

In the 24 FRPD samples considered in this study, two presented contamination by mycotoxins, with one of the samples presenting multiple contamination with 3 mycotoxins found in the same sample. To our knowledge, this is the first study developed where used FRPD showed contamination by mycotoxins. Considering that FRPD is positioned in the workers' nose and mouth we can conclude that workers are exposed by inhalation and/or ingestion to mycotoxins, namely fumonisins B1 (FB1) and B2 (FB2) and sterigmatocystin (STE), the ones detected in our study. Human consumption of foods containing FB1 may cause esophageal cancer, and liver and kidney disease (Chen et al. 2021). FB1 is also known to have the potential to cause heart failure in humans due to damage in myocardial contractility and massive blood influx, known as Idiopathic Congestive Cardiopathy (Stoev 2015).

Although no women are working in the production area in the company engaged in this study, fumonisins are also known to be responsible for causing a high risk of bearing a child with a birth defect of the brain or spinal cord in women when exposure to high levels occurs by food consumption during early pregnancy (Missmer et al. 2006; Waes et al. 2005).

Sterigmatocystin is a known biogenic precursor of aflatoxin B1, sharing with it several structural and biological similarities (Zingales et al. 2020). This mycotoxin is hepatotoxic and nephrotoxic in animals and it has been classified as a possible human carcinogen (group 2B) by IARC (IARC, 1987).

As mentioned before, the presence of multiple mycotoxins can result in different types of interactions when co-exposure occurs, as it seems to be the case.

Results regarding mycotoxins were different from a study developed in one waste-sorting industry from Portugal, where it was not found contamination by mycotoxins in the used FRPD, even with a high fungal burden observed (Viegas et al. 2020a). The FRPD used by workers in this company (BRF) was a half mask used consecutive 5 h and without daily replacement. This is a different usage regime than the study developed previously by Viegas C. et al. (2020a), where the FRPD was replaced daily, and this might explain the contamination found in this study.

The mycotoxin contamination found in the settled dust samples (two samples showed contamination; 15.3%) followed the same trend as a previous study developed in a waste collection centre also located in Portugal (Viegas et al. 2022a), where the settled dust samples collected in the waste collection trucks assessed also presented contamination by mycotoxins, however with higher number of samples being contaminated (16 samples; 53.1%). Of note, the chemical contamination can explain the fact that only settled dust samples from SRE presented mycotoxin contamination since lower chemical contamination is expected in SRE than in BRF, allowing the fungal species to thrive (Gomes et al. 2020; Roane et al. 2009). Concerning the detected mycotoxins, the profile reported was the same observed by Viegas and colleagues (2022a), being the mycophenolic acid and sterigmatocystin the mycotoxins detected and also with mycophenolic acid presenting the higher concentrations.

Cytotoxicity

Several mycotoxins, including the ones usually produced by fungi abundant in the used FRPD analyzed in this study (such as *Penicillium* sp. or *Aspergillus* sp.), are greatly related to cytotoxicity in different cells, such as HepG2 cells (Sobral et al. 2018; Zhou et al. 2022; Vu et al. 2022). These secondary metabolites are produced by fungi as a defense mechanism and present toxicity for humans and animals when ingested or inhaled. The described toxicity mechanisms include cell disruption and induction of cellular death (Sobral et al. 2018; Zhou et al. 2022; Vu et al. 2022). In our study, cytotoxicity was observed in most of the used FRPD tested in lung epithelial A549 cells, and to a lower extent, in hepatic HepG2 cells. However, only a small number of used FRPD exhibited mycotoxin contamination. This can be due to the interaction between mycotoxins and other fungal metabolites or with other FRPD components, such as particulate matter or chemicals. The exposure of cells to mixtures of toxic components can lead to a synergistic cytotoxic effect due to exacerbated cellular responses, due to interference with cellular uptake or cellular detoxifying pathways, for instance. Thus, while mycotoxins significantly contribute for cellular damage, other microbial and particulate components

also play a role in cytotoxicity. It should be underscored the importance of assessing the exposure to multiple components and their possible interactions in order to comprehend the possible health risks associated.

Conclusions

The study developed allowed to recognize some specific features in this occupational setting concerning contamination/exposure to microbiologic agents:

- The used FRPD should be used in sampling protocols to assess occupational exposure to microbiologic agents since they suffer less influence from the chemical burden common in this setting;
- Microbial contamination with pathogenic and toxigenic potential was found in both analysed matrices (used FRPD and settled dust);
- *Aspergillus* section *Fumigati* (surrogate of harmful fungal contamination in the waste management industry) was found widespread in the analyzed units;
- *Aspergillus* section *Flavi* results, together with previous studies concerning AFB1 exposure, suggest potential workers exposure to AFB1;
- The common and already reported chemical contamination in this setting might explain the reduced contamination of settled dust samples by microorganisms and mycotoxins;
- Exposure to mycotoxins might occur through the use of FRPD, particularly when FRPD it is not replaced daily.

Further studies focusing on occupational exposure to microbiologic agents in e-waste recycling should be prioritized in light of the results obtained in this study. In addition, future studies should target the co-assessment of chemical contaminants (besides mycotoxins), to explain the importance of cytotoxicity assessment in composite environmental samples and to characterize better the overall workers 'exposure.

This study findings need to be considered when improving the risk management measures to implement in the e-waste recycling industry aiming to control workers' exposure by using FRPD. Other measures should be put in place, such as decreasing exposure and number of workers exposed. Lastly, training and education programs for workers and employers should cover information on the proper use of FRPD and replacement frequency to avoid contamination and, consequently, increased exposure from workers.

Supplementary Information The online version contains supplementary material available at <https://doi.org/10.1007/s41742-024-00677-4>.

Author Contributions All authors contributed to the study's conception and design. Material preparation, data collection and analysis were performed by CV, BG, MD, RC, PP, SG, MA-S, EC, MT, LAC and SV. The first draft of the manuscript was written by Carla Viegas and all authors commented on previous versions of the manuscript. All authors read and approved the final manuscript.

Funding Open access funding provided by FCTIFCCN (b-on). This research was funded by the Polish Minister of Education and Science, under the program "Regional Initiative of Excellence" in 2019–2022 (Grant No. 008/RID/2018/19) and by Instituto Politécnico de Lisboa, Lisbon, Portugal for funding the Projects the Projects IPL/IDI&CA2024/WWTPSValor_ESTeSL and IPL/IDI&CA2024/MycoSOS_ESTeSL. H&TRC authors gratefully acknowledge the FCT/MCTES national support through the UIDB/05608/2020 (<https://doi.org/https://doi.org/10.54499/UIDP/05608/2020>) and UIDP/05608/2020 (<https://doi.org/https://doi.org/10.54499/UIDB/05608/2020>) and to the PhD Grants UI/BD/151431/2021 and UI/BD/153746/2022 and CE3C unit UIDB/00329/2020. We thank the companies and workers for participating in the study.

Declarations

Conflict of Interest The authors have no relevant financial or non-financial interests to disclose.

Ethical Approval This study complied with the Helsinki Declaration and Oviedo Convention and all data were stored and analyzed by the Portuguese General Data Protection Regulation (GDPR) law n° 58/2019.

Consent to Participate Workers were invited to voluntarily participate in the project. Before the enrolment, all volunteers filled out a written informed consent.

Consent to Publish Companies' owners were invited to voluntarily participate in the project and filled out a written informed consent regarding permission for results publication.

Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>.

References

- Akram R, Shah Fahad N, Zaffar Hashmi M, Wahid A, Adnan M, Mubeen M et al. (2019) Trends of electronic waste pollution and its impact on the global environment and ecosystem. *Environ Sci Pollut Res*. 26, 16923–16938. <https://doi.org/10.1007/s11356-019-04998-2>.
- Andersson MA, Vornanen-Winqvist C, Koivisto T, Varga A, Mikkola R, Kredics L, Salonen H (2023) Composition of culturable microorganisms in dusts collected from sport facilities in finland during

- the COVID-19 pandemic. *Pathogens* 12(2):339. <https://doi.org/10.3390/pathogens12020339>
- Caramalho R, Tyndall JDA, Monk BC, Larentis T, Lass-Flörl C, Lackner M (2017) Intrinsic short-tailed azole resistance in mucormycetes is due to an evolutionary conserved amino acid substitution of the lanosterol 14 α -demethylase. *Sci Rep* 21;7(1). <https://doi.org/10.1038/s41598-017-16123-9>.
- Chao HJ, Milton DK, Schwartz J, Burge HA (2002) Dustborne fungi in large office buildings. *Mycopathologia* 154:93–106. <https://doi.org/10.1023/A:1015592224368>
- Chen J, Wen J, Tang Y, Shi J, Mu G, Yan R, Cai J, Long M (2021) Research progress on fumonisin B1 contamination and toxicity: a review. *Molecules*. 29;26(17):5238. <https://doi.org/10.3390/molecules26175238>.
- Cox CS (1998) The microbiology of air. In Topley & Wilson's microbiology and microbial infections (eds L. Collier, A. Balows & M. Sussman), 9th edn. London, UK: Arnold, Oxford University Press. 339–350.
- Cox J, M'bareche H, Lindsley WG, Duchaine C (2020) Field sampling of indoor bioaerosols. *Aerosol science and technology. J Am Assoc Aerosol Res.* 572–584. <https://doi.org/10.1080/02786826.2019.1688759>
- Cruz-Perez P, Buttner MP, Stetzenbach LD (2001) Detection and quantitation of *Aspergillus fumigatus* in pure culture using polymerase chain reaction. *Mol Cell Probes* 15(2):81–88. <https://doi.org/10.1006/mcpr.2000.0343>
- Dannaoui E (2017) Antifungal resistance in mucorales. *Int J Antimicrob Agents* 50(5):617–621. <https://doi.org/10.1016/j.ijantimicag.2017.08.010>
- DeLong EF, Pace NR (2001) Environmental diversity of bacteria and archaea. *Syst Biol* 50(4):470–478
- DeGois J, Clerc F, Simon X, Bontemps C, Leblond P, Duquenne P (2017) First metagenomic survey of the microbial diversity in bioaerosols emitted in waste sorting plants. *Annals of Work Exposures and Health* 61(9):1076–1086. <https://doi.org/10.1093/annweh/wxx075>
- Donlan RM (2002) Biofilms: microbial life on surfaces. *Emerg Infect Dis* 8:881–890. <https://doi.org/10.3201/eid0809.020063>
- Eriksen E, Graff P, Pedersen I, Straumfors A, Afanou AK (2022) Bioaerosol exposure and in vitro activation of toll-like receptors in a norwegian waste sorting plant. *Saf Health Work* 13(1):9–16. <https://doi.org/10.1016/j.shaw.2021.09.002>
- European Commission 2023. Circular economy action plan. https://environment.ec.europa.eu/strategy/circular-economy-action-plan_en. Accessed in 23.07.2023.
- European Parliament. 2023. E-waste in the EU: facts and figures (infographic) <https://www.europarl.europa.eu/news/en/headlines/society/20201208STO93325/e-waste-in-the-eu-facts-and-figures-infographic>. Accessed in 23.7.2023.
- Fisher MC, Alastruey-Izquierdo A, Berman J, Bicanic T, Bignell EM, Bowyer P, Bromley M, Brüggemann R, Garber G, Cornely OA, Gurr SJ, Harrison TS, Kuijper E, Rhodes J, Sheppard DC, Warris A, White PL, Xu J, Zwaan B, Verweij PE (2022) Tackling the emerging threat of antifungal resistance to human health. *Nat Rev Microbiol* 20(9):557–571. <https://doi.org/10.1038/s41579-022-00720-1>
- Global Action for Fungal Infections (2022) Fungal Disease Frequency. <https://gaffi.org/why/fungal-disease-frequency>. Accessed in 30.7.2023.
- Gomes IB, Maillard JY, Simões LC et al. (2020) Emerging contaminants affect the microbiome of water systems—strategies for their mitigation. *npj Clean Water* 3, 39. <https://doi.org/10.1038/s41545-020-00086-y>.
- Gonçalves P, Melo A, Dias M, Almeida B, Caetano LA, Veríssimo C, Viegas C, Sabino R (2021) Azole-resistant *aspergillus fumigatus* harboring the TR34/L98H mutation: first report in portugal in environmental samples. *Microorganisms* 9:57. <https://doi.org/10.3390/microorganisms9010057>
- Gravel B, Roberge B, Calosso M, Gagné S, Lavoie J, Labrèche F (2023) Occupational health and safety, metal exposures and multi-exposures health risk in Canadian electronic waste recycling facilities. *Waste Manage* 165:140–149. <https://doi.org/10.1016/j.wasman.2023.04.026>
- Hanelt M, Gareis M, Kollarczik B (1994) Cytotoxicity of mycotoxins evaluated by the MTT-cell culture assay. *Mycopathologia* 128:167–174. <https://doi.org/10.1007/BF01138479>
- Hoog GS de, Queiroz-Telles F, Haase G, Fernandez-Zeppenfeldt G, Angelis DA, Gerrits van den Ende HG, Matos A, Peltroche-Llacsahuanga T, Pizzirani-Kleiner H, Rainer AA, Richard-Yegres J, Vicente NV, Yegres F (2000) Black fungi: Clinical and pathogenic approaches. *Med Mycol* 38: 243–250
- International agency for research on cancer. 2002. Aflatoxin. <https://monographs.iarc.who.int/wp-content/uploads/2018/06/mono100F-23.pdf>. Accessed in 30.7.2023
- International agency for research on cancer. 1987. Sterigmatocystin. <https://inchem.org/documents/iarc/vol01/sterigmatocystin.html>. Accessed in 30.7.2023
- International labour organization. 2021. Exposure to hazardous chemicals at work and resulting health impacts : a global review. https://ilo.org/wcmsp5/groups/public/---ed_dialogue/---lab_admin/documents/publication/wcms_795460.pdf. Accessed in 29.7.2023
- Jankowska E, Reponen T, Willeke K, Grinshpun SA, Choi KJ (2000) Collection of fungal spores on air filters and spore reentrainment from filters into air. *J Aerosol Sci* 31:969–978. [https://doi.org/10.1016/S0021-8502\(00\)00017-3](https://doi.org/10.1016/S0021-8502(00)00017-3)
- Kung VL, Chernock RD, Burnham CD (2017) Diagnostic accuracy of fungal identification in histopathology and cytopathology specimens. *Eur J Clin Microbiol Infect Dis* 37(1): 157–165. <https://doi.org/10.3390/jof8050505>.
- Lavoie J, Dunkerley CJ, Kosatsky T, Dufresne A (2006) Exposure to aerosolized bacteria and fungi among collectors of commercial, mixed residential, recyclable and compostable waste. *Sci Total Environ* 370:23–28. <https://doi.org/10.1016/j.scitotenv.2006.05.016>
- Li L, Mendis N, Trigui H, Oliver JD, Faucher SP (2014) The importance of the viable but non-culturable state in human bacterial pathogens. *Front Microbiol* 5, 258. <https://doi.org/10.3389/fmicb.2014.00258>
- Madsen AM, Alwan T, Ørberg A, Uhrbrand K, Jørgensen MB (2016) Waste workers' exposure to airborne fungal and bacterial species in the truck cab and during waste collection. *Ann Occup Hyg* 60:651–668. <https://doi.org/10.1093/annhyg/mew021>
- Majchrzycka K, Okrasa M, Skóra J, Gutarowska B (2016) Evaluation of the survivability of microorganisms deposited on filtering respiratory protective devices under varying conditions of humidity. *Int J Environ Res Public Health* 13:98. <https://doi.org/10.3390/ijerph13010098>
- Marchand G (2021) Chapitre 8: microorganismes et leurs sous-produits. In: Hygiène du travail (Ed.), du diagnostic à la maîtrise des facteurs de risques. Montréal: TC Média livres inc. 165–196.
- Marchand G, Bonifait L, Veillette M, Pepin C, Beaudet Y, Legare E Lavoie L., et al., 2017. Évaluation des bioaérosols et des composés gazeux émis lors des compostages de résidus agroalimentaires et résidentiels. IRSST. <http://www.irsst.qc.ca/media/documents/PubIRSST/R-960.pdf>. Accessed in 30.7.2023
- Mensah-Attipoe J, Täubel M (2017) Analysis approaches for fungi in indoor environmental assessments. In: Viegas C, Viegas S, Gomes AQ, Täubel M, Sabino R (Eds) Exposure to Microbiological Agents in Indoor and Occupational Environments. Springer. 109–127. https://doi.org/10.1007/978-3-319-61688-9_6.
- Missmer SA, Suarez L, Felkner M, Wang E, Merrill A, Rothman K, Hendricks KA (2006) Exposure to fumonisins and the occurrence

- of neural tube defects along the Texas-Mexico Border. *Environ Health Perspect* 114:237–241. <https://doi.org/10.1289/ehp.8221>
- Mayer Z, Bagnara A, Färber P, Geisen R (2003) Quantification of the copy number of nor-1, a gene of the aflatoxin biosynthetic pathway by real-time PCR, and its correlation to the cfu of *Aspergillus flavus* in foods. *Int J Food Microbiol* 82(2):143–151. [https://doi.org/10.1016/s0168-1605\(02\)00250-7](https://doi.org/10.1016/s0168-1605(02)00250-7)
- Nageen Y, Wang X, Pecoraro L (2023) Seasonal variation of airborne fungal diversity and community structure in urban outdoor environments in Tianjin. *China Front Microbiol* 9(13):1043224. <https://doi.org/10.3389/fmicb.2022.1043224>
- Okeme, JO., Arrandale, VH. 2019. Electronic Waste Recycling: Occupational Exposures and Work-Related Health Effects. *Curr Environ Health Rep.* 2019 Dec;6(4), 256–268. <https://doi.org/10.1007/s40572-019-00255-3>.
- Park D-U, Ryu S-H, Kim S-B, Yoon C-S (2011) An assessment of dust, endotoxin, and microorganism exposure during waste collection and sorting. *J Air Waste Manag Assoc* 61:461–468. <https://doi.org/10.3155/1047-3289.61.4.461>
- Roane MT, Rensing C, Pepper IL, Raina M, Maier RM (2009) Microorganisms and Metal Pollutants, Ed: Raina M. Maier, Ian L. Pepper, Charles P. Gerba, Environmental Microbiology (Second Edition), Academic Press, 421–441. <https://doi.org/10.1016/B978-0-12-394626-3.00018-1>.
- Salabanga FRD, Wingert L, Valois I, Lacombe N, Gouin F, Trépanier J, Debia M, Soszczyńska E, Twarużek M, Kosicki R, Dias M, Viegas S, Caetano L, Viegas C, Marchand G (2022) Microbial contamination and metabolite exposure assessment during waste and recyclable material collection. *Environ. Res.* 212. <https://doi.org/10.1016/j.envres.2022.113597>
- Scheepers PTJ, Duca RC, Galea KS, Godderis L, Hardy E, Knudsen LE, Leese E, Louro H, Mahiout S, Ndaw S et al (2021) HBM4EU occupational biomonitoring study on e-Waste—study protocol. *Int J Environ Res Public Health* 18:12987. <https://doi.org/10.3390/ijerph182412987>
- Sobral M, Faria MA, Cunha SC, Ferreira IM (2018) Toxicological interactions between mycotoxins from ubiquitous fungi: Impact on hepatic and intestinal human epithelial cells. *Chemosphere* 202:538–548. <https://doi.org/10.1016/j.chemosphere.2018.03.122>
- Stoev SD (2015) Foodborne mycotoxicoses, risk assessment and underestimated hazard of masked mycotoxins and joint mycotoxin effects or interaction. *Environ Toxicol Pharmacol* 39:794–809. <https://doi.org/10.1016/j.etap.2015.01.022>
- United Nations Institute for Training and Research (UNITAR) 2024. <https://ewastemonitor.info/the-global-e-waste-monitor-2024/> Accessed on 8 of July 2024
- Viegas C, Dias M, Almeida B, Aranha Caetano L, Carolino E, Quintal Gomes A, Twarużek M, Kosicki R, Grajewski J, Marchand G, Viegas S (2020a) Are workers from waste sorting industry really protected by wearing Filtering Respiratory Protective Devices? The gap between the myth and reality. *Waste Manage* 102:856–867. <https://doi.org/10.1016/j.wasman.2019.12.001>
- Viegas C, Dias M, Almeida B, Carolino E, Quintal Gomes A, Viegas S (2020b) *Aspergillus* spp. burden on filtering respiratory protective devices. Is there an occupational health concern? *Air Quality, Atmosphere & Health*, 1–10. <https://doi.org/10.1007/s11869-019-00781-x>.
- Viegas C, Dias M, Almeida B, Gomes AQ, Twarużek M, Kosicki R, Grajewski J, Viegas S (2021^a) Mechanic protection gloves used in waste sorting industry: Suitable as passive method to sample for fungal burden. 42nd Mycotoxin Workshop.
- Viegas C, Dias M, Carolino E, Sabino RFP (2021b) Culture media and sampling collection method for *Aspergillus* spp. assessment: Tackling the gap between recommendations and the scientific evidence. *Atmosphere*, 12(1), 23. <https://doi.org/10.3390/atmos12010023>.
- Viegas C, Faria T, de Oliveira AC, Caetano LA, Carolino E, Quintal-Gomes A, Twarużek M, Kosicki R, Soszczyńska E, Viegas S (2017) A new approach to assess occupational exposure to airborne fungal contamination and mycotoxins of forklift drivers in waste sorting facilities. *Mycotoxin Res* 33:285–295. <https://doi.org/10.1007/s12550-017-0288-8>
- Viegas C, Faria T, dos Santos M, Carolino E, Gomes AQ, Sabino R, Viegas S (2015a) Fungal burden in waste industry: an occupational risk to be solved. *Environ Monit Assess* 187:199. <https://doi.org/10.1007/s10661-015-4412-y>
- Viegas C, Gomes B, Cervantes R, Moreira S, Dias M, Pena P et al. (2023) Microbial contamination in grocery stores from Portugal and Spain—The neglected indoor environment to be tackled in the scope of the One Health approach. *Sci.Total Environ.* 875. <https://doi.org/10.1016/j.scitotenv.2023.162602>.
- Viegas C, Pena P, Dias M, Gomes B, Cervantes R, Carolino E, Twarużek M, Soszczyńska E, Kosicki R, Caetano LA, Viegas S (2022a) Microbial contamination in waste collection: Unveiling this Portuguese occupational exposure scenario. *Journal of Environmental Management*, 314. <https://doi.org/10.1016/j.jenvman.2022.115086>.
- Viegas C, Twarużek M, Dias M, Almeida B, Carolino E, Kosicki R, Soszczyńska E, Grajewski J, Aranha Caetano L, Viegas S (2020c) Assessment of the microbial contamination of mechanical protection gloves used on waste sorting industry: a contribution for the risk characterization. *Environ. Res.* 189. <https://doi.org/10.1016/j.envres.2020.109881>.
- Viegas C, Twarużek M, Dias M, Almeida B, Carolino E, Soszczyńska E, Viegas S, Aranha Caetano L (2021c) Cytotoxicity of filtering respiratory protective devices from the waste sorting industry: A comparative study between interior layer and exhalation valve. *Environment International* 155. <https://doi.org/10.1016/j.envint.2021.106603>.
- Viegas C, Twarużek M, Dias M, Carolino E, Soszczyńska E, Aranha Caetano L (2022b) Cytotoxicity of *Aspergillus Section Fumigati* Isolates Recovered from Protection Devices Used on Waste Sorting Industry. *Toxins* 14(2):70. <https://doi.org/10.3390/toxins14020070>
- Viegas S, Veiga L, Figueiredo P, Almeida A, Carolino E, Viegas C (2014) Assessment of Workers' Exposure to Aflatoxin B1 in a Portuguese Waste Industry. *Ann Occup Hyg.* <https://doi.org/10.1093/annhyg/meu082>
- Viegas S, Veiga L, Figueiredo P, Almeida A, Carolino E, Viegas C (2015b) Assessment of Workers' Exposure to Aflatoxin B1 in a Portuguese Waste Industry. *Ann Occup Hyg* 59(2):173–181. <https://doi.org/10.1093/annhyg/meu082>
- Viegas S, Osteresch B, Almeida A, Cramer B, Humpf H-U, Viegas C (2018) Enniatin B and ochratoxin A in the blood serum of workers from the waste management setting. *Mycotoxin Res* 34:85–90. <https://doi.org/10.1007/s12550-017-0302-1>
- Vu THN, Pham NS, Le PC et al. (2022) Distribution, cytotoxicity, and antioxidant activity of fungal endophytes isolated from *Tsuga chinensis* (Franch.) Pritz. in Ha Giang province, Vietnam. *Ann Microbiol* 72, 36. <https://doi.org/10.1186/s13213-022-01693-5>.
- Waes JG-V, Starr L, Maddox J, Aleman F, Voss KA, Wilberding J, Riley RT (2005) Maternal fumonisin exposure and risk for neural tube defects: mechanisms in an In Vivo mouse model. *Birth Defects Res. Part A Clin Mol Teratol* 73:487–497. <https://doi.org/10.1002/bdra.20148>
- Whitby C, Ferguson RMW, Colbeck I, Dumbrell AJA, Nasir ZA, Marczylo E, Kinnersley R, Douglas P, Drew G, Bhui K, Lemon M, Jackson S, Tyrrel S, Coulon F (2022) Chapter Three—Compendium of analytical methods for sampling, characterization and quantification of bioaerosols, Edit(s): David A. Bohan, Alex

- Dumbrell. *Adv Eco Res Academic Press* 67:101–229. <https://doi.org/10.1016/bs.aecr.2022.09.004>
- Won WD, Ross H (1966) Effect of diluent and relative humidity on apparent viability of airborne *Pasteurella pestis*. *Appl Microbiol* 14:742–745. <https://doi.org/10.1128/am.14.5.742-745.1966>
- World Health Organization. 2022. WHO Fungal Priority Pathogens List to Guide Research, Development and Public Health Action. <https://www.who.int/publications/i/item/9789240060241>. Accessed in 18.7.2023.
- Wouters IM, Spaan S, Douwes J, Doekes G, Heederik D (2005) Overview of Personal Occupational Exposure Levels to Inhalable Dust, Endotoxin, $\beta(1\rightarrow3)$ -Glucan and Fungal Extracellular Polysaccharides in the Waste Management Chain. *Ann Occup Hyg* 39–53. <https://doi.org/10.1093/annhyg/mei047>.
- Zhou J, Feng Z, Zhang W et al. (2022) Evaluation of the antimicrobial and cytotoxic potential of endophytic fungi extracts from mangrove plants *Rhizophora stylosa* and *R. mucronata*. *Sci Rep* 12: 2733. <https://doi.org/10.1038/s41598-022-06711-9>.
- Zingales V, Fernández-Franzón M, Ruiz MJ (2020) Sterigmatocystin: Occurrence, toxicity and molecular mechanisms of action—a review. *Food Chem Toxicol* 146. <https://doi.org/10.1016/j.fct.2020>.

Authors and Affiliations

Carla Viegas^{1,2}  · Bianca Gomes^{1,3} · Marta Dias^{1,2} · Renata Cervantes^{1,2} · Pedro Pena^{1,2} · Sara Gonçalves¹ · Marina Almeida-Silva^{1,4} · Elisabete Carolino¹ · Magdalena Twarużek⁵ · Liliana Aranha Caetano^{1,6} · Susana Viegas^{1,2}

✉ Carla Viegas
carla.viegas@estesl.ipl.pt

¹ H&TRC—Health & Technology Research Center, ESTeSL—Escola Superior de Tecnologia e Saúde, Instituto Politécnico de Lisboa, 1990-096 Lisbon, Portugal

² Public Health Research Centre, Comprehensive Health Research Center, CHRC, REAL, CCAL, NOVA National School of Public Health, NOVA University Lisbon, 1099-085 Lisbon, Portugal

³ CE3C—Center for Ecology, Evolution and Environmental Change, Faculdade de Ciências, Universidade de Lisboa, 1749-016 Lisbon, Portugal

⁴ OSEAN—Outermost Regions Sustainable Ecosystem for Entrepreneurship and Innovation, 9000-039 Funchal, Portugal

⁵ Faculty of Biological Sciences, Department of Physiology and Toxicology, Kazimierz Wielki University, Chodkiewiczza 30, 85-064 Bydgoszcz, Poland

⁶ Research Institute for Medicines (iMed.uLisboa), Faculty of Pharmacy, University of Lisbon, Lisbon, Portugal