Working Paper nº 51

Non-Linear Time Series Models

Kamil Feridun Turkman

Working Paper nº 51

ISSN: 0872-895X

Depósito Legal nº: 90631/95

Kamil Feridun Turkman
DEIO e CEAUL
Universidade de Lisboa

Maio 1996

Trabalho apresentado no âmbito do Seminário de Estatística e Econometria -
coordenado pelo Professor Doutor Bento Murteira

NON-LINEAR TIME SERIES MODELS

K.F. Turkman

DEIO and CEAUL, University of Lisbon

Abstract:

The last three decades have seen quite dramatic changes the way we modeled time dependent data. Linear processes have been in the center stage in modeling time series. As far as the second order properties are concerned, the theory and the methodology are very adequate. However, there are more and more evidence that linear models are not sufficiently flexible and rich enough for modeling purposes and that failure to account for non-linearities can be very misleading and have undesired consequences. Together with these evidences, quite a lot of different classes of non-linear models have been suggested, taking into account different aspects of non-linearity. In this paper, we look briefly some of these classes by discussing their basic properties and by suggesting when they might be appropriate.

1 Introduction

Until recently, the general linear model

$$X_t = \sum_{j=0}^{\infty} \psi_j \epsilon_{t-j}, \quad (1)$$

where, $\{\psi_j\}_{j=0}^{\infty}$ are constants such that

$$\sum_{j=0}^{\infty} \psi_j^2 < \infty$$

and $\{\epsilon_j\}$ are uncorrelated random variables with finite variance (usually normally distributed), played the central role in modeling time series. It is known for sometimes that any purely non-deterministic process can be expressed in this linear form and it is also known that, for such time series, the auto correlation function can well be approximated by that of an finite order $ARMA(p, q)$ model. Hence, as far as the second order moments are concerned, the class of linear models are flexible and rich enough for time series analysis. The linear model in (1) has its own characteristics:

- If ϵ_t are normally distributed (that is if X_t is a Gaussian process) then the conditional mean of X_t given the past observations $x_t, x_{t-1}, \dots$ will be a linear function of $x_t, x_{t-1}, \dots$
- the conditional variance of X_t given the past observations will be a finite constant
- the process X_t will be time reversible.
- Linear models have light tails and are not suitable for modeling series with sudden burst of large values.

Recently, we are becoming more and more aware that there are quite a lot of time series which do not fit to these restrictions. For example, there are quite a lot of time series in econometric and environmental sciences which seem to be compatible with heavy tailed marginal distributions (i.e. time series with infinite variances) or time series with finite variances but which change randomly in time. In some cases, the assumption of linear conditional

mean function is incompatible with the observed sample path. For such cases, the class of linear models are not sufficiently flexible and rich for modeling purposes.

It may be tempting to describe these deviations from the linearity by non-gaussian ARMA models, where the innovations ϵ_t are uncorrelated but with a distribution other than normal. Such linear models can be constructed to take into account non-reversibility, non-linearity in the conditional mean and infinite variance cases. Indeed, In practice it is not very easy to differentiate the symptoms of non-normality from that of non-linearity and one may be tempted to use a non-Gaussian linear model. However, we no longer have a guarantee that such a model will indeed be a good approximation and that the second order moments can be used for model identification. In fact, it is shown by Feigin and Resnick(1996) that the use of linear models and the covariances as tool to model a non-linear process would be extremely misleading. We will later look at this point more closely in section 3.

A result by Nisio (1960) says that almost every stationary time series can be approximated in the following polynomial representation (or Volterra expansion)

$$X_t = \sum_p \sum_{i_1, \dots, i_p} g_{i_1 \dots i_p} \prod_{v=1}^p \epsilon_{t+i_v}, \quad (2)$$

where $\{\epsilon_i\}$ are independent rather than uncorrelated, contrary to what is assumed for the linear representation given in (1). For $p = 1$, the representation in (2) is linear and such representation exists only when the process X_t is Gaussian. This result clearly indicates that if we look for general representations in terms of independent innovations rather than uncorrelated ones, then we should consider non-linear models.

In this paper, we will not deal how we test for the presence of non-linearity; we will look at various classes of non-linear models and discuss when they would be appropriate models. We will not touch the difficult problem of model identification, estimation and diagnostics of non-linear models. In most cases, these steps of model building are still in infancy, and a lot of research is yet to be done in these areas.

It is not entirely clear what is meant by a non-linear model due to the fact that it is not very clear what is meant by a linear model. We will assume that a linear model takes the form (1) and that any deviation from (1) will be considered as a non-linear model.

A non-linear model for a time series X_t is a relationship between $\{X_{k+i}, |i| = 0, 1, 2, \dots\}$ which produces a sequence of zero mean, iid random variables. Thus a non-linear model for X_k may be described by an equation of the form

$$h(\dots, X_{k-1}, X_k, X_{k+1}, \dots) = \epsilon_t, \quad (3)$$

where h is a general non-linear function. We assume that the model in (3) is invertible so that it can be written in the equivalent form

$$X_k = f(\dots, \epsilon_{k-1}, \epsilon_k, \epsilon_{k+1}, \dots). \quad (4)$$

Assuming that f can be expanded in a Taylor series about the origin, X_k can be represented in what is called discrete Volterra series. (see for example, Priestley, 1981)

The Basic principle behind the Volterra series expansion is to write a general non-linear process

$$X_t = h(\epsilon_t, \epsilon_{t-1}, \dots)$$

in a Taylor series expansion of the form

$$\begin{aligned} X_t = \mu &+ \sum_{|i| \leq \infty} g_i \epsilon_{t-i} + \sum_{|i| \leq \infty} \sum_{|j| \leq \infty} g_{i,j} \epsilon_{t-i} \epsilon_{t-j} \\ &+ \sum_{|i| \leq \infty} \sum_{|j| \leq \infty} \sum_{|l| \leq \infty} g_{i,j,l} \epsilon_{t-i} \epsilon_{t-j} \epsilon_{t-l} \\ &+ \dots, \end{aligned} \quad (5)$$

$$= \sum_p \sum_{i_1, \dots, i_p} g_{i_1 \dots i_p} \prod_{v=1}^p \epsilon_{t+i_v} \quad (6)$$

where $\{\epsilon\}$ are iid random variables. Note that (6) gives a very suitable representation of X_t in terms of the first order (linear), second, third, etc. order moving averages. The Volterra series representation is very general as well; almost any stationary time series may be approximated by (6), and hence is the non-linear version of the Wold decomposition. However, due to having infinitely many parameters, such representation is not suitable for statistical applications and the study of classes of non-linear time series represented in terms of finite parameter difference equations is important.

We take the readers attention to the relation between general stationary linear model

$$X_t = \sum_{|i| \leq \infty} g_i \epsilon_{t-i}$$

and the class of ARMA models, which is a finite parameter sub-class of the general linear model.

We now look at such various classes.

2 Classes of non-linear models

We can broadly divide non-linear models into two separate groups: parametric and non-parametric models.

2.1 parametric models

2.1.1 Parametric models for the conditional mean

Here, the objective is to model the conditional mean function of the process X_t as a non-linear function of the past observations, assuming that the conditional variance is still a constant. Hence an appropriate general model is given as

$$X_t = f(X_{t-1}, X_{t-2}, \dots, X_{t-p}, \theta) + \epsilon_t, \quad (7)$$

here the function f is known and θ is an unknown vector parameter to be estimated. Several different forms of f give different classes of non-linear models: (For ease in notation, we represent the simpler case when $f(X_{t-1}, X_{t-2}, \dots, X_{t-p}, \theta) = f(X_{t-1}, \theta) = f(x, \theta)$.)

1. *Threshold Autoregressive models:*

$$f(x, \theta) = \theta_1 x 1_{(x \in A^{(1)})} = \theta_2 x 1_{(x \in A^{(2)})}, \quad (8)$$

where, $A^{(1)}$ and $A^{(2)}$ are certain regions, so that the process can be written as

$$X_t = \begin{cases} \theta_1 X_{t-1} + \epsilon_t & \text{if } X_{t-1} \in A^{(1)} \\ \theta_2 X_{t-1} + \epsilon_t & \text{if } X_{t-1} \in A^{(2)} \end{cases}$$

A common form of the region A is given as $A^{(1)} = \{X_{t-1} \geq y\}$, and $A^{(2)} = \{X_{t-1} < y\}$, for some y .

We can generalize this class of models to

$$X_t = \sum_{j=1}^p \theta_{ij} X_{t-j} + \epsilon_t^{(j)} \quad , \quad (X_{t-1}, \dots, X_{t-p} \in A^{(i)} \quad , \quad i = 1, 2, \dots, l) \quad (9)$$

thus taking into account for possible variation in variability in different random regions as well as the different levels of the conditional mean. Such models are known to be very useful, particularly in modeling data which shows random cyclic movements. We exemplify one situation where such models can be very useful. Consider the population of a rat community, where there are no natural predators. Under such circumstances, the rat population would increase explosively (a first order Autoregressive model with positive parameter, probably with a value larger than 1) until the population grows to a certain size when the available food can not sustain the rat population. Then the model inverts and behaves as a first order autoregressive model with a negative parameter. Here the random cyclic nature of the process is evident and is very much tied down the population size threshold which would depend on external factors such as the amount of food stuff available, thus on climatic conditions etc. Note that threshold models are piecewise linear models where the passage from one linear form to another is random.

2. Exponential and logistic autoregressive models:

$$f(x, \theta) = \theta(x)x, \quad (10)$$

so that the process can be written in the form

$$X_t = \theta(X_{t-1})X_{t-1} + \epsilon_t,$$

an AR(1) process with a random coefficient depending on the last observation. We can assume some specific forms for the parameter, such as

$$\theta(x) = \theta_1 + \theta_2 \exp(-\theta_3 x^2)$$

In this case, the model is called the exponential AR(1) model. If

$$\theta(x) = \theta_1 x + \theta_2 x \{ [1 + \exp(-\theta_3(x - \theta_4))]^{-1} - \frac{1}{2} \} \quad , \quad \theta_3 > 0,$$

then the model is called logistic AR(1). There is no reason why the parametric functions should be restricted to these specific forms and much more general forms will be given later in regime models.

These models can be generalized to the form

$$X_t = \mathbf{a}^t \mathbf{X}_{t-1} + \sum_{i=1}^k b_i \phi_i(\gamma_i^t \mathbf{X}_{t-1}) + \epsilon_t, \quad (11)$$

where $\mathbf{X}_{t-1} = (X_{t-1}, X_{t-2}, \dots, X_{t-p})$, and $\mathbf{a}^t = (a_1, a_2, \dots, a_p)$, $\gamma^t = (\gamma_1, \gamma_2, \dots, \gamma_p)$ are p -dimensional parameter vectors and $\phi_i(\cdot)$ are known specific functions. Although such models are used, it is evident that there would be problems of estimation as the parameter space increases.

2.1.2 Parametric models for the conditional variance

The motivation behind these models is to predict not only the mean but also the variance of the process as a function of the past values of the time series. These models have found many applications in econometrics, since the variance of the process can be seen as the risk or the volatility in economic processes.

1. ARCH models

The first order ARCH (autoregressive conditional Heteroskedastic) model given by

$$X_t = (\theta_1 + \theta_2 X_{t-1}^2)^{1/2} \epsilon_t \quad (12)$$

is a good example for this class. Note that

$$Var(X_t / X_{t-1} = x) = (\theta_1 + \theta_2 x^2) \sigma^2.$$

First order ARCH model can be generalized to the form

$$X_t = g(\mathbf{X}_t - 1, \boldsymbol{\theta}) \epsilon_t,$$

where g is a known positive function. It is also possible to introduce an AR part and recursions in the conditional variance structure of the residuals. such extension results in the class of GARCH (generalized ARCH) models.

2.1.3 Mixed models

These models contain both conditional mean and variance components and has the general form

$$X_t = f(X_{t-1}, X_{t-2}, \dots, X_{t-p}, \theta_1) + g(X_{t-1}, X_{t-2}, \dots, X_{t-p}, \theta_2)\epsilon_t, \quad (13)$$

where $g(\cdot)$ is a positive function. Note that

$$E(X_t/X_{t-1} = x_1, X_{t-2} = x_2, \dots, X_{t-p} = x_p) = f(x_1, x_2, \dots, x_p, \theta_1), \quad (14)$$

$$V(X_t/X_{t-1} = x_1, X_{t-2} = x_2, \dots, X_{t-p} = x_p) = g(x_1, x_2, \dots, x_p, \theta_2)^2 V(\epsilon_t). \quad (15)$$

Here, the function f and g can be chosen in accordance with the partial models discussed in sections on models for the conditional mean and variance. However, the most powerful class of non-linear models which falls in this classification of mixed models is the class of bilinear processes.

1. Bilinear models

The process X_t is said to be a bilinear process $BL(p, r, m, k)$ if it satisfies the difference equation

$$\begin{aligned} X_t = & \sum_{j=1}^p a_j X_{t-j} + \sum_{j=0}^r c_j \epsilon_{t-j} \\ & + \sum_{l_1=1}^m \sum_{l_2=1}^k b_{l_1 l_2} X_{t-l_1} \epsilon_{t-l_2}, \end{aligned} \quad (16)$$

The class of bilinear models given in (16) plays an important role in modeling non-linearity for various reasons;

- The class is an obvious generalization of ARMA(p,r) models.
- It is a dense class in the class of non-linear processes and covers virtually all types of non-linearity, provided that the time interval of interest is finite. The reason for this is that any bilinear process can be uniquely written in terms of Volterra series expansion and hence it is a non-linear analog of the linear Wold representation. (Note however that limit cycles, jump phenomenon are pertinent to infinite time problems and hence these types of non-linearity can not be modeled by a bilinear process. Threshold models discussed above are the appropriate models for such time series)

- The class is fairly well studied. Much is known regarding the existence of unique and stationary solutions. Although Identification, estimation and diagnostic techniques are available, still much work on the class such as invertibility conditions and better estimation methods remains to be completed. There is also available software for modeling. (Subba Rao and Gabr, 1984, da Silva, 1994)
- Bilinear models can be written under certain conditions, in terms of a multivariate random coefficient autoregressive process, which we will explain later in this chapter.
- Bilinear processes are capable of producing sudden bursts of large values and hence are very suitable in modeling time series showing burst-like phenomena. (See section 3, for a sample path of a bilinear model)

2. Regime models

This class of models are introduced to take into account in stochastic changes that may exist in the structure of the underlying process. A simple model within this class is the first order AR model

$$X_t = \theta_t X_{t-1} + \epsilon_t,$$

where $\{\theta_t\}$ is a homogeneous markov chain with a finite space and transition probabilities $\{p_{ij}\}$, for example taking values a_1 and a_2 . In this case, The process X_t will alternate between the two processes

$$X_t = a_1 X_{t-1} + \epsilon_t,$$

and

$$X_t = a_2 X_{t-1} + \epsilon_t,$$

according to the transition probabilities $\{p_{ij}\}$. In general a regime model will take the form $\{X_t, S_t\}$, where, $\{S_t\}$ is an unobserved process, possibly a homogeneous markov chain with a finite state space, such that at any time t X_t conditional on $S_t = j$ follows a linear model $ARMA_j(p_j, q_j)$. Hence the process will alternate among various linear models in accordance with the transient behavior of the unobserved process S_t . The estimation, identification and diagnostics for these

models are complicated although not impossible, due to the fact that the process S_t is not observed.

In Figure 1, we have a sample path of the process

$$X_t = C^{(S_t)} + 0.5X_{t-1} + \epsilon_t,$$

where

$$C^{(S_t)} = \begin{cases} 2 & \text{if } S_t = 1 \\ 10 & \text{if } S_t = 10 \end{cases}$$

and S_t is a homogeneous Markov chain with transition probabilities $p_{11} = 0.9$ and $p_{22} = 0.6$.

In Figure 2, we have a sample path from the process

$$X_t = 0.5X_{t-1} + \epsilon_t^{(S_t)},$$

where

$$\epsilon_t^{(S_t)} = \begin{cases} N(0, 1) & \text{if } S_t = 1 \\ N(0, 10) & \text{if } S_t = 2 \end{cases}$$

Figure 1

Figure 2

3. Polynomial models

A regression style model, perhaps the most natural and useful is the class of polynomial models given by

$$X_t = \sum_{i=1}^k a_i X_{t-1}^i + \epsilon_t. \quad (17)$$

More general polynomial models can be devised by introducing terms depending on $X_{t-2}, X_{t-3}, \dots, X_{t-p}$ and cross terms. These models are not very much used due to the feedback of X_t into itself, causing explosive behavior.

2.2 Non-parametric models

In the class of parametric models, the main emphasis was on building parametric models for the conditional mean and variance of the process, either separately or jointly. If the emphasis is on prediction rather than on explaining how these conditional means and variances change in time, then a plausible alternative is to estimate them non-parametrically. This would be quite flexible, since one is not restricted by a specific parametric model. The most common way is to use kernel estimators. For example, For a given time series $X_t, t = 1, \dots, n$, The conditional mean

$$M(x_1, x_2, \dots, x_p) = E(X_t / X_{t-1} = x_1, \dots, X_{t-p} = x_p)$$

can be estimated by

$$\hat{M}(x_1, \dots, x_p) = \frac{(n-p)^{-1} \sum_{t=p+1}^n X_t \prod_{i=1}^p K_h(X_{t-i} - x_i)}{(n-p+1)^{-1} \sum_{t=p+1}^n \prod_{i=1}^p K_h(X_{t-i} - x_i)}, \quad (18)$$

where

$$K_h(x) = \frac{1}{h} K\left(\frac{x}{h}\right),$$

and K is a Kernel function. A similar expression can be obtained for the conditional variance.

The drawback with these models is that one need extremely large sample sizes for a reasonable fit and the sample needs increase drastically with the increase in p , a typical case of the curse of dimension. In general, assuming

that a good parametric model is found for the data, these non-parametric models do not perform as good as the parametric ones.

There are other approaches to non-parametric models, taking care of or at least reducing the curse of dimension. Such models are known as the semi-parametric and restricted non-parametric models.

3 Why non-linear models are so different from linear models

The Wold decomposition says that almost any finite variance stationary process can be written as a linear form. But this representation is correct as far as the second properties of the process is concerned. That is, given a stationary finite variance process, one can find a linear process whose moments up to second order are equal to this process. We can not say much for the higher order moments. They may be totally different. Let us look at a very simple bilinear process

$$X_t = aX_{t-1} + bX_{t-1}\epsilon_{t-1} + \epsilon_t.$$

We can write the same process as

$$X_t = (a + b\epsilon_{t-1})X_{t-1} + \epsilon_t,$$

a AR(1) process with a random coefficient.

Figures 3 and 4 given below show how a simple bilinear model differs from the corresponding AR(1) model without the random perturbation in the coefficient.

As can be seen, the bilinear process has sudden burst of large values, indicating probably that the bilinear process may have infinite variance. Indeed, under quite general conditions X_t has a stationary solution given by

$$X_\infty = \sum_{k=0}^{\infty} \prod_{j=1}^k (a + b\epsilon_j) \epsilon_k, \quad (19)$$

and

$$\lim_{x \rightarrow \infty} P(X_\infty > x) = c_+ x^{-\alpha},$$

$$X_t = -0.1 X_{t-1} + \epsilon_t, \quad \epsilon_t \sim N(0, 1)$$

FIG. 3

A realization of the model $X_t = (-0.1 + 0.9\epsilon_t)X_{t-1} + \epsilon_t$, $\epsilon_t \sim \mathcal{N}(0, 1)$.
(The theoretical standard deviation of X_t is 2.357)

FIG. 4

where $\alpha > 0$ is such that $E|a + b\epsilon_1|^\alpha = 1$. (See Goldie, 1991 and Turkman and Amaral Turkman, 1996)

Hence, even if the residuals are normally distributed, the process X_t has a regularly varying tails and can have infinite variance for $\alpha < 2$. For stationary time series with infinite variance there is no analog of linear process representation or approximation to it. One interesting aspect of these bilinear models is their extremal properties. That is, explaining how this explosive phenomena occurs (See for example Turkman and Amaral Turkman, 1996)

One final caution regarding nonlinearity: We know that if X_t is a finite variance stationary sequence, under quite general conditions, the sample autocorrelation function is consistent and converges the true autocorrelations. On the other hand, it is known that (Davis and Resnick, 1996) if X_t is non-linear then the autocorrelation function converges to a random variable. Thus using the autocorrelations and partial autocorrelations as tools for modeling when there is nonlinearity in the model can be extremely misleading. In fact this result can be used as a quick check if the time series is linear or non-linear: After making sure that the time series is stationary, one can plot the autocorrelation function for different portions of the series and compare. If there is a clear stability and these plot look the same then one can conclude that there are no nonlinearity present. If the plots show erratic behavior for different portions, then this would be a clear sign of nonlinearity.

4 Acknowledgements

I am grateful to Dra Marilia Reis for supplying the simulated series and their graphical representations.

5 References

R.A. Davis and S.I. Resnick (1995) Limit theory for bilinear processes with heavy tailed noise. *Annals of Appl. Probab.* To be published.

Goldie, C. M. (1991) Implicit renewal theory and tails of solutions of random equations. *Ann. Appl. Probab.* 1, 126-166.

Kesten, H. (1973) Random difference equations and renewal theory for products of random matrices. *Acta Math.* 131, 207-248.

Nisio, M (1960) On polynomial approximation for strictly stationary processes. *J.Math. Soc. Jpn.*, 12, 207-226.

Priestley, M.B. (1981) *Spectral Analysis and Time Series*. Academic Press, London.

Subba Rao, T. and M.M. Gabr (1984) *An Introduction to Bispectral Analysis and Bilinear Time Series Models* Springer-Verlag, New York.

Tjøstheim, D. (1994) Non-linear time series: A selective review. *Scand. J. of Statist.* 21, 97-130.

Turkman, K.F. and M.A. Amaral Turkman (1995) Extremes of bilinear time series models. Technical report, *Center of Statistics, University of Lisbon*. (Accepted for publication in *J. Time series Analysis*)