

## Limnic ostracoda from Silveirinha, Portugal (? Late Paleocene - Lowermost Eocene)

Jean-Paul Colin<sup>(1)</sup> & Miguel Telles Antunes<sup>(2,3,4)</sup>

Projecto "POCTI/36531/PAL/2000 – Studies on Portuguese Palaeontology (Post-Paleozoic)", FCT Fundação para a Ciência e a Tecnologia/  
Ministério da Ciência e das Universidades.

1. 3 Impasse des Biroulayres, 33610 Cestas, France. jean-paul.colin@worldonline.fr. 2. Academia das Ciências de Lisboa. 3. European Academy of Arts, Sciences and Humanities, Paris. 4. Centro de Estudos Geológicos, Faculdade de Ciências e Tecnologia (UNL)/ Quinta da Torre 2829-516 Caparica, Portugal. mta@fct.unl.pt

### Abstract

**Key-words:** Ostracoda; ?upper Paleocene/ lowermost Eocene; Silveirinha; Portugal.

Silveirinha, a rich site for mammals and other vertebrates, yielded some molluscs, charophytes and ostracods. The latter are described here. Eight species were recognized. Two of them, *Ilyocypris lusitanicus* n. sp. and *Cypris silveirinhaensis* n. sp. are new. Other, unnamed species may be ascribed to *Cyclocypris?* and *Cypria?*, besides a further four undeterminate Cypridacea. The Ostracod association is typical for limnic environments and is compatible with intertropical conditions. *Cypris silveirinhaensis* suggests an Eocene, and especially a lower Eocene age.

### Resumo

**Palavras-chave:** Ostracodos; ?Paleocénico superior/ Eocénico basal; Silveirinha; Portugal.

A jazida de Silveirinha, rica de mamíferos e outros vertebrados, deu também moluscos, carófitos e ostracodos, descritos neste trabalho. Foram reconhecidas oito espécies, duas novas: *Ilyocypris lusitanicus* n. sp. e *Cypris silveirinhaensis* n. sp. Além destas, ocorrem espécies atribuíveis, possivelmente, aos géneros *Cyclocypris* (uma) e *Cypria* (uma), além de quatro de género(s) indeterminado(s) da superfamília Cypridacea. A associação, tipicamente indicativa de meios límnicos, é compatível com ambientes inter-tropicais. Do ponto de vista cronológico, *Cypris silveirinhaensis* sugere idade do Eocénico, em particular do Eocénico inferior.

### 1. Introduction

Vertebrate remains have been collected by R. Pena dos Reis (University of Coimbra) in a clay pit at Silveirinha, lower Mondego area, West-central Portugal. The concerned lithostratigraphic unit is the "Areias e argilas de Taveiro", whose lower levels are late Cretaceous in age, mainly comprises greenish, clayey beds crossed by fluvial channels (Reis & Soares, 1981). Pelitic sediments are nearly sterile under a palaeontologic viewpoint. On the other hand, the channels' infillings are fossil-rich microconglomerates.

Sedimentation occurred in an alluvial plain, maybe in bogs that often (seasonally?) were flooded through crevasse-splay channels.

The first fossils were sent to M. T. Antunes, who identified some mammals that pointed out to lowermost Eocene (Antunes, 1981). That age was then recognized for the first time in Portugal (ibid.). A fuller account was published soon after (Antunes & Russell, 1981).

A Research Program has been developed since under the direction of M. T. Antunes. Field work and subsequent washing and sieving at the Earth Sciences Department of the Faculdade de Ciências e Tecnologia (Universidade

Nova de Lisboa) of more than 10 thousand kg of sediments allowed us to gather an impressive collection. Silveirinha became one of Western Europe's richest sites for those times. It yielded a rich mammalian fauna (for a synthesis, see Antunes *et al.*, 1997) that was recognized as older than Doormal's (in Belgium, the type site for the MP7 mammal-unit). Later on, a few similar mammalian faunas from the Pyrenean region were shown to lay stratigraphically under marine Paleocene beds (Gheerbrant *et al.*, 1997). An upper Paleocene age cannot therefore be excluded as far as Silveirinha is concerned.

Other groups are represented: birds, including a wader (Harrison, 1983), a small-sized crocodylian, genus *Diplocynodon* (Antunes, under printing), snakes and lizards as well as amphibians (Rage, this volume), Pelomedusid chelonians (De Lapparent de Broin, idem), percomorph fishes (Gaudant & Antunes, idem), scarce (except *Bythinia*) gastropods and exceedingly rare lamellibranchs (Callapez, idem), ostracods, worms or other trace-fossil producers, scarce seeds, charophyte gyrogonites (identified by J. Riveline, Antunes & Colin, this volume), besides coproliths and a few amber fragments.

Most fossils underwent diagenetic silicification and water transportation. They are often abraded.

## 2. The Ostracoda

The limnic ostracode assemblage found in Silverinha is well diversified, with 8 species recognized.

Illustrated material is provisionally deposited in the collections of the Centro de Estudos Geológicos, Faculdade de Ciências e Tecnologia (UNL).

Subclass	O S T R A C O D A Latreille, 1806
Order	PODOCOPIDA Müller, 1894
Suborder	PODOCOPA Sars, 1866
Superfamily	Cypridacea Baird, 1845
Family	Ilyocyprididae Kaufmann, 1900
Genus	<i>Ilyocypris</i> Brady & Norman, 1889

*Ilyocypris lusitanicus* n. sp.  
(Pl. 1, figs. 1-2, 4, 8)

**Derivation of name:** from Portugal.

**Holotype:** 1 carapace (Pl. 1, fig. 2).

**Paratypes:** 3 carapaces.

**Material:** 5 carapaces.

**Diagnosis:** species of *Ilyocypris*, strongly reticulate, with 3 dorsal, a sub-median and a postero-ventral nodes.

**Dimensions:** L = 0.73 mm – 0.75 mm; h = 0.40 mm; e = 0.34 mm.

**Remarks:** Related species are *Ilyocypris essertinesensis* Carbonnel *et al.* 1985 from the Chattian (Oligocene) of Switzerland, which differs by the absence of a postero-ventral tubercle and is slightly larger; and

*Ilyocypris cornae* Mandelstam, 1961 common in the Eocene of China (Hou *et al.*, 1978), which also lacks the postero-ventral node.

Family	Candonidae Kaufmann, 1900
Subfamily	Cyclocypridinae Kaufmann, 1900
Genus	<i>Cyclocypris</i> Brady & Norman, 1889

*Cyclocypris* ? sp.  
(Pl. 2, fig. 8)

**Dimensions:** L = 0.51 mm; h = 0.46 mm.

**Remarks:** This species, tentatively assigned to the genus *Cyclocypris*, is morphologically close to *Porpocypris sphaeroidalis* Guan 1978 and *Cypris globra* Chen, 1982 (in Hou *et al.*) from the uppermost Cretaceous of China (Zhang, 1992).

Genus *Cypria* Zenker, 1854

*Cypria* ? sp.  
(Pl. 2, figs. 7, 9, 10)

**Dimensions:** L = 0.43 mm; h = 0.29 mm.

**Remarks:** Rather small species, strongly compressed. A related species, *Cypris dorsalta* Malz & Moyadepour, 1973 from the Miocene and Oligocene (Carbonnel *et al.*, 1985) of respectively Germany and Switzerland, is higher.

Family	Cyprididae Baird, 1845
Subfamily	Cypridinae Baird, 1845
Genus	<i>Cypris</i> Müller, 1898

*Cypris silveirinhaensis* n. sp.  
(Pl. 1, figs.3, 5-7)

**Derivation of name:** from the type locality Silveirinha.

**Holotype:** Carapace (Pl. 1, fig. 5).

**Paratypes:** 3 carapaces.

**Material:** 12 specimens.

**Type locality and horizon:** Silveirinha, Portugal, Lower Eocene.

**Diagnosis:** Species of the genus *Cypris* with the following characters: moderate size carapace, finely reticulated, weak anterior lip.

**Dimensions:** L = 1.22 – 1.39 mm; h = 0.76-0.90 mm; e = 0.89 mm.

**Remarks:** *Cypris silveirinhaensis* is closely related to the following species:

- *Cypris* sp. Tambareau *et al.* 1991, from the Ilerdian of the Montagne-Noire (Southern France), is smooth or very finely punctated, higher, with the greatest height in a median position.

- *Cypris pseudodecaryi* Guernet 1981 from the Sparnacian of the Paris Basin has a more compressed anterior margin, its dorsal margin is more inclined and the ornamentation much finer.

- *Cypris decaryi* Gauthier, 1933, a recent intertropical species, is larger, higher and finely punctate (Neale, 1976b).

- *Cypris henanensis* Guan & Sun 1978 (= *Cypris* cf. *decaryi* auct.) from the Eocene of China (Guan, 1984; Guan & Sun, 1978; Zheng, 1983; Hou *et al.* 1978, 1982; Hao, 1988; Hao *et al.*, 1988; He & Zhang, 1982; He *et al.*, 1988), has a higher and more symmetrical carapace and a very finely punctate carapace.

- *Cypris favosa* Ye (in Hou *et al.*, 1978) from the Eocene of China has a similar micro-reticulate ornamentation but the carapace is narrower and more elongated.

- *Cypris subglobosa* Sowerby, 1840 *sensu* Neale, 1976a, a recent species from India, Sri Lanka and Eastern Europe, is more elongated and the reticulation meshes are larger.

Indet. Cypridacea

Indet. Cypridacea sp. 1  
(Pl. 2, fig. 5)

**Dimensions:** L = 1.45 ; h = 0.93 mm.

**Remarks:** outline comparable to *Hebeina ovata* Boje, 1978 from the Oligocene of China.

Indet. Cypridacea sp. 2  
(Pl. 2, fig. 6)

**Dimensions:** L = 1.23 mm; h = 0.92 mm.

Indet. Cypridacea sp. 3  
(Pl. 1, figs. 8, 10; Pl. 2, figs. 1, 2)

**Dimensions:** L = 0.78- 0.81 mm; h = 0.46-0.50 mm.

**Remarks:** morphologically related forms from the Cenozoic of China have been referred to the genus *Eucypris* by many Chinese authors (Hou *et al.*, 1978, 1982).

Indet. Cypridacea sp. 4  
(Pl. 2, figs. 3, 4)

**Dimensions:** L = 0.96 – 0.98.

**Remarks:** the same as for Cypridacea sp. 3.

### 3. Conclusions

1) The ostracod association from Silveirinha comprises the following species (absence of loose valves hampers generic attribution of some):

*Ilyocypris lusitanicus* n. sp.  
*Cycloocypris* ? sp.  
*Cypris* ? sp.  
*Cypris silveirinhaensis* n. sp.  
Indet. Cypridacea sp. 1  
Indet. Cypridacea sp. 2  
Indet. Cypridacea sp. 3  
Indet. Cypridacea sp. 4.

2) The most diagnostic species are *Ilyocypris lusitanicus* n. sp. and *Cypris silveirinhaensis* n. sp.

3) The presence of *Cypris silveirinhaensis* n. sp. suggests an Eocene age as this species is very closely related to: *Cypris* sp., described by Tambareau *et al.*, 1991 from the Ilerdian of Southern France; *Cypris pseudodecaryi* Guernet, 1981 from the Sparnacian of the Paris Basin; and *Cypris henanensis* Guan & Sun, 1978, from the Eocene of China.

4) The Silveirinha's association of ostracods clearly points out to limnic environments.

5) The same association seems compatible with intertropical conditions.

### Acknowledgements

We are pleased to express our sincere acknowledgements to Dr. P. Carbonel (Talence, France) and C. Guernet (Paris, France) for the SEM photographs, Dr. E. Kempf (Kiel, Germany) for bibliographic help, and Dr. A. Nascimento (Monte de Caparica, Portugal) for preliminary examination of the Silveirinha ostracodes.

## References

- Antunes, M. Telles (1981) – As areias e argilas de Silveirinha II – Aspectos paleontológicos. *Memórias e Notícias, Publ. Mus. Lab. Mineral. Geol., Univ. Coimbra*, 91-92: 255-267.
- Antunes, M. Telles; Casanovas, M.L.; Cuesta, M.A.; Checa, L.I.; Santafé, J.V. & Agustí, J. (1997) – Eocene mammals from the Iberian Peninsula. *Actes du Congrès Biochrom'97*, J.-P. Aguilar, S. Legendre & J. Michaux (Eds.). *Mém. Trav. E.P.H.E., Inst. Montpellier*, 21: 337-352, 2 fig.
- Antunes, M. Telles & Russell, D. E. (1981) – Le gisement de Silveirinha (Bas Mondego, Portugal): la plus ancienne faune de Vertébrés éocènes connue en Europe. *C. R. Acad. Sci. Paris*, 293 (21 Décembre 1981), Série II: 1099-1102.
- Boije, (1978) - *Early Tertiary ostracod fauna from the coastal region of Bohai*. Nanjing: 1-242.
- Carbonnel, G.; Weidmann, M. & Berger, J.-P. (1985) - Les ostracodes lacustres et saumâtres de la Molasse de Suisse occidentale. *Revue de Paléobiologie*, Genève, 4, 2: 215-251.
- Gheerbrant, E.; Abrial, C. & Cuenca, G. (1997) – Nouveau Mammifère condylarthre du Paléocène supérieur de Campo (Pyrénées aragonaises, Espagne). *C. R. Acad. Sci. Paris*, 324, Série IIa: 599-606, 3 fig., 1 pl.
- Guan Shao-Zeng (1984) - Early Tertiary ostracods from Liguanqiao basin in the border area of Henan and Hubei, with a note on the ontogeny of *Cypris henaensis*. *Acta Micropaleontol. Sinica*, Beijing, 1, 1: 90-98.
- Guan Shao-Zeng (1988) - Palaeocene non-marine ostracods in China. In: Hanai, T., Ikeya, N. & Ishizaki, K. eds. *Evolutionary Biology of Ostracoda. Its fundamentals and applications*. Elsevier-Kodansha, Amsterdam-Tokyo: 1147-1152
- Guernet, C. (1981) - Ostracodes sparnaciens du Bassin de Paris (France). *Revue de Micropaléontologie*, Paris, 24, 1: 51-66.
- Haoyi-Chun (1988) - Cretaceous and Palaeogene ostracod biostratigraphy in Xining and Minhe basins of China. In: Hanai, T., Ikeya, N. & Ishizaki, K. eds. *Evolutionary Biology of Ostracoda. Its fundamentals and applications*. Elsevier-Kodansha, Amsterdam-Tokyo: 1163-1171.
- Hao Yichun; Zhou Xiugao; Song Qushan; Yang Guodong; Cheng Sguwei & Wei Zhenxin (1988) - Middle Jurassic-Tertiary deposits and ostracod-charophyta fossil assemblages of Xining and Minhe basins. *Journal Wuhan College Geology*, 23: 1-210.
- Harrison, C. J. O. (1983) – A new wader (Charadriiformes) from the early Eocene of Portugal. *Ciências da Terra (UNL)*, Lisboa, 7: 9-16, 2 fig.
- He Jun-de & Zhang Xiao-Jun (1982) - Cretaceous-Tertiary ostracods from the Yangtze-Han river basin in Hubei. *Bulletin of Nanjing Institute Geology Palaeontology*, Academia Sinica, Nanjing, 5: 177-208.
- He Jun-de; Van Nieuvanuise, D.S. & Swain, F.M. (1988) - Biostratigraphy of Paleogene non-marine ostracoda from East China. In: Hanai, T.; Ikeya, N. & Ishizaki, K. eds. *Evolutionary Biology of Ostracoda. Its fundamentals and applications*. Elsevier-Kodansha, Amsterdam-Tokyo: 1153-1161.
- Hou You-Tang; Chen Te-Chiuung; Yang Heng-Ren; Ho Jun-de; Zhiu Quan-Chun & Tian Mu-Qu (1982) - Cretaceous-Quaternary ostracod fauna from Jiangsu. *Geol. Publ. House*, Beijing: 1-386.
- Hou You-Tang; Ho Jun-de & Ye Chun-Hui (1978) - The Cretaceous-Tertiary ostracods from the marginal region of the Yangtze-Han river plain in Central Hubei. *Mem. Nanjing Inst. Geol. Palaeontol.*, Academia Sinica, 9: 129-206.
- Keen, M. C. (1975) - The palaeobiology of some upper Palaeogene fresh-water ostracodes. In Swain, F.M., Kornicker, L.S. & Lundin, R.F. eds., *Biology and Palaeobiology of Ostracoda*. *Bull. Amer. Palaeontol.*, 65, 282: 271-283.
- Malz, H. (1976) - *Heterocypris* vel *Cyprinotus*. *Senckenbergiana Lethaea*, Frankfurt-am-Main, 57, 2/3: 185-199.
- Malz, H. (1977) - Cypridopsine Ostracoden aus dem Tertiär des mainzer Beckens. *Senckenbergiana Lethaea*, Frankfurt-am-Main, 58, 4/5: 219-261.
- Malz, H. & Moyaedpour, E. (1973) - Miozäne Süßwasser-Ostracoden aus der Rhön. *Senckenbergiana Lethaea*, Frankfurt-am-Main, 54, 2/4: 281-309.
- Neale J.W. (1976a) - On *Cypris subglobosa* J. de C. Sowerby. *Stereo-Atlas of Ostracod Shells*, London, 3, 2: 125-132.
- Neale, J.W. (1976b) - On *Cypris decaryi* Gauthier. *Stereo-Atlas of Ostracod Shells*, London, 3, 2: 133-140.
- Reis, R. Pena dos & Soares, A. Ferreira (1981) – As areias e argilas de Silveirinha. I – Aspectos sedimentológicos. *Memórias Notícias, Publ. Mus. Lab. Mineral. Geol., Univ. Coimbra*, n.º 91-92: 245-255, 8 fig.
- Tambareau, Y. (1984) - Les ostracodes du "Montien continental" de Hainin, Hainault, Belgique. *Revue Micropaléontologie*, Paris, 27, 2: 144-156.
- Tambareau, Y.; Gruas-Cavagnetto, C.; Feist, M. & Villatte, J. (1991) - Flores et faunes continentales ildardiennes du versant Sud de la Montagne Noire et de la Montagne d'Alaris. *Revue Micropaléontologie*, Paris, 34, 1: 69-89.

- Zhang Xian-Qiu (1985) - Early Tertiary ostracods from Dongguan basin, Guangdong. *Acta Palaeontologica Sinica*, Beijing, 24, 4: 413-423.
- Zhang Xian-Qiu (1991) - Discussion on the Cretaceous-Tertiary boundary in Nanxiong basin in the light of ostracod fauna. *Guangdong geol.*, 6, 3: 61-72.
- Zhang Xian-Qiu (1992) - A study on ostracod fauna of Shanghu Formation and Cretaceous-Tertiary boundary in Nanxiong basin, Guangdong. *Acta Palaeontologica Sinica*, Beijing, 31, 6: 678-702.
- Zheng Shu-Ying (1983) - Early Tertiary non-marine ostracodes from Changhe of Cixi, Zhejiang. *Acta Palaeontologica Sinica*, Beijing, 22, 5: 577-581.

**Plate 1**

1, 2, 4, 8 – *Ilyocypris lusitanicus* n. sp.

1. Carapace, left view, x 80.
2. Carapace, right view, holotype, x 80.
4. Carapace, dorsal view, x 80.
8. Carapace, dorsal view, slightly tilted, x 80.


3, 5-7 – *Cypris silveirinhaensis* n. sp.

3. Carapace, ventral view, x 40.
5. Carapace, right view, holotype, x 45.
6. Carapace, left view, x 40.
7. Carapace, right view, x 45.

9, 10 – Indet. Cypridacea sp. 1

9. Carapace, male, right view, x 70.
10. Carapace, female, right view, x 70.

PLATE 1



**Plate 2**

1, 2 – Indet. Cyprideacea sp. 1

1. Carapace, female left view, x 80.
2. Carapace, female, right view, x 80.

3-4 – Indet. Cypridacea sp. 2

3. Carapace, male, right view, x 60.
4. Carapace, female, right view, x 80.

5 – Indet. Cypridacea sp. 3.

Carapace, right view, x 40.

6 – Indet. Cypridacea sp. 4

Carapace, left view, x 40.

7, 9, 10 – *Cypria* sp.

7. Carapace, right view, x 115.
9. Carapace, left view, x 115.
10. Carapace, right view, x 130.

8 – *Cyclocypris* ? sp.

Carapace, right view, x 80.

PLATE 2

