

## Rationale and design of Long-term Outcomes and Vascular Evaluation after Successful Coarctation of the Aorta Treatment study

[Jose D Martins](#), [Justin Zachariah](#),<sup>1</sup> [Elif Seda Selamet Tierney](#),<sup>2</sup> [Uyen Truong](#),<sup>3</sup> [Shaine A Morris](#),<sup>1</sup> [Shelby Kutty](#),<sup>4</sup> [Sarah D de Ferranti](#),<sup>5</sup> [Jonathan Rhodes](#),<sup>5</sup> [Marta Antonio](#),<sup>5</sup> [Maria Guarino](#),<sup>6</sup> [Boban Thomas](#),<sup>7</sup> [Diana Oliveira](#),<sup>8</sup> [Kimberlee Gauvreau](#),<sup>5</sup> [Nuno Jalles](#),<sup>7</sup> [Tal Geva](#),<sup>5</sup> [Miguel Carmo](#),<sup>6</sup> and [Ashwin Prakash](#)<sup>5</sup>

Department of Pediatric Cardiology, Hospital de Santa Marta, Centro Hospitalar de Lisboa Central, Lisbon, Portugal

<sup>1</sup>Division of Pediatric Cardiology, Texas Children's Hospital, Baylor College of Medicine, Houston, USA

<sup>2</sup>Division of Pediatric Cardiology, Department of Pediatrics, Lucile Packard Children's Hospital, Stanford University, Palo Alto, USA

<sup>3</sup>Division of Pediatric Cardiology, Children's Hospital Colorado, Aurora, USA

<sup>4</sup>Joint Division of Pediatric Cardiology, Children's Hospital and Medical Center University of Nebraska College of Medicine, Omaha, USA

<sup>5</sup>Department of Cardiology, Boston Children's Hospital, Harvard Medical School, Boston, USA

<sup>6</sup>CEDOC Chronic Diseases, Nova Medical School, Lisbon, Portugal

<sup>7</sup>Caselas, Ressonância Magnética, S. A. Lisbon, Portugal

<sup>8</sup>Biomedical Engineering Department, Instituto Superior Técnico, Lisbon, Portugal

**Address for correspondence:** Dr. José D Martins, Department of Pediatric Cardiology, Centro Hospitalar de Lisboa Central, Hospital de Santa Marta, Rua de Santa Marta, 50, 1150-024, Lisbon, Portugal. E-mail: [jdferreiramartins@gmail.com](mailto:jdferreiramartins@gmail.com)

**Copyright** : © 2018 Annals of Pediatric Cardiology

This is an open access journal, and articles are distributed under the terms of the Creative Commons Attribution-NonCommercial-ShareAlike 4.0 License, which allows others to remix, tweak, and build upon the work non-commercially, as long as appropriate credit is given and the new creations are licensed under the identical terms.

### Abstract

[Go to:](#)

### Background:

[Go to:](#)

Coarctation of the aorta (CoA) can be treated using surgery, balloon angioplasty, or stent implantation. Although short-term results are excellent with all three treatment modalities, long-term cardiovascular (CV) morbidity and mortality remain high, likely due to persistently abnormal vascular function. The effects of treatment modality on long-term vascular function remain uncharacterized. The goal of this study is to assess vascular function in this patient population for comparison among the treatment modalities.

### Methods:

[Go to:](#)

We will prospectively assess vascular Afunction in large and small arteries fusing multiple noninvasive modalities and compare the results among the three groups of CoA patients previously treated using surgery, balloon angioplasty, or stent implantation after frequency matching for confounding variables. A comprehensive vascular function assessment protocol has been created to be used in 7 centers. Our primary outcome is arterial stiffness measured by arterial tonometry. Inclusion and exclusion criteria have been carefully established after consideration of several potential confounders. Sample size has been calculated for the primary outcome variable.

### Conclusion:

[Go to:](#)

Treatment modalities for CoA may have distinct impact on large and small arterial vascular function. The results of this study will help identify the treatment modality that is associated with the most optimal level of vascular function, which, in the long term, may reduce CV risk.

**Keywords:** Arterial stiffness, cardiac magnetic resonance imaging, coarctation of the aorta, long-term outcomes, pulse wave velocity, vascular function

## INTRODUCTION

[Go to:](#)

Current treatment techniques are equally effective at eliminating the stenosis in CoA patients.[1] However, a good anatomical result does not preclude late systemic hypertension in office visits (12%–65%), [1,2,3,4,5,6] at peak exercise (10%–47%),[3,4,5,7,8] or during ambulatory blood pressure (BP) monitoring (30%–59%).[7,8,9,10] Furthermore, treated patients have reduced life expectancy [Figure 1],[2] mostly due to cardiovascular (CV) complications[2,11,12,13,14] and stroke.[15]

Successfully treated CoA patients have stiffer large arteries[16,17,18,19,20,21] and compromised vascular reactivity in small arteries,[8,10,22,23,24,25,26] their arterial pressure waveform is altered,[9,10,23,27,28] have imbalances in vascular function biomarkers,[24,25,27,29,30] and increased left ventricular (LV) mass.[8,9,19,20,21,26,31,32] Vascular dysfunction is associated with older age at treatment, [2,19,22,29,31,33] but early treatment does not guarantee normal vascular function.[16,22]

Different treatment modalities may have varying effects on the stiffness of the repaired arterial segment:[34] Surgical repair results in a focal scar in the anastomosis; stenting creates a short, rigid segment; and balloon dilation (BD) produces a controlled tear of the intima and part of the media. Although it is possible that these differences translate into differences in vascular dysfunction, this has not been systematically compared. The largest, albeit observational and nonrandomized, comparison between the three modalities showed a lower BP in patients treated with BD versus those treated with stenting or surgery.[1] A small retrospective study showed less frequent exercise-induced hypertension in BD patients compared with other treatment types.[33] Conclusions drawn from these prior studies are hampered by methodological limitations and limited focus. In the general population, arterial stiffness is associated with major CV events.[35] Thus, choosing the CoA treatment option that optimizes vascular function is crucial for long-term outcomes in CoA.

### Aim and hypothesis

The Long-term Outcomes and Vascular Evaluation after Successful Coarctation of the Aorta Treatment study aims to determine whether surgery, BD, and stenting are associated with differences in arterial stiffness in optimally treated patients. Our hypothesis is that patients who underwent successful BD will have better vascular function than patients who underwent successful surgical repair or stenting since this modality may least likely damage the biomechanical properties of the aortic wall.

## METHODS

[Go to:](#)

### Study overview

This study is a cross-sectional prospective observational study of patients with CoA previously treated using one of three treatment modalities. Patients will be recruited at seven large pediatric cardiac centers from Europe and the United States of America [Appendix 1]. The study procedures will occur in a 1- or 2-day visit [Figure 2].

### Recruitment

Selection criteria are depicted in Table 1. The study protocol was approved by Institutional Review Boards. Recruitment occurred between June 2013 and December 2017. The study data are collected and managed using REDCap software, hosted at Children's Hospital Boston.[36]

### Study procedures

A list with the main clinical and study tests variables are depicted in Tables 2 and 3. The comprehensive list of study variables is in Appendix 7

### Arterial stiffness

CoA treatments alter the biomechanics of the isthmus and may increase arterial stiffness. The velocity of the pulse wave velocity (PWV) travel in the arterial tree increases with arterial stiffness. Carotid–femoral

PWV (cfPWV) is extensively validated in large studies a marker of aortic stiffness, and an independent predictor of CV events.[37] We will measure cfPWV with applanation tonometry, using either the NIHem (CV Engineering, Inc., Norwood, MA USA) or the SphygmoCor (AtCor Medical, West Ryde, NSW, Australia) devices.[37] This technique assumes a homogenous stiffness across the aorta and may potentially not accurately estimate the true carotid-to-femoral artery length. Cardiovascular magnetic resonance (CMR) measurements of PWV, on the other way, enables the detection of more subtle changes in segmental aortic PWV, above versus below the CoA site, and uses real aortic travel paths.[38] We will also use CMR to measure aortic area change during the cardiac cycle, paired with BP measurements, to quantify local arterial strain, compliance, distensibility, and the  $\beta$ -stiffness index [Appendix 2a and b].

### Endothelial function

In CoA, the loss of central aortic pulsatility, which buffers systole, generates chronic shear stress downstream in smaller arteries, creating endothelial dysfunction, which is associated with CV events.[39] We will measure endothelial function with the reactive hyperemia index using endothelial pulse amplitude tonometry (endo-PAT), a novel noninvasive and reproducible technique that measures changes in pulsatile arterial volume with a fingertip probe.[40] Analysis of the pulse waveform allows for automated calculation of endothelial function in one arm, while the contralateral serves as control, making this a patient standardized method [Appendix 3].

### Pulse waveform analysis

In CoA, the stiff aorta and repaired isthmus may be important reflecting sites that impact the pulse waveform. Its analysis is an important clinical tool for monitoring of vascular function and predicting CV events.[37] We will measure three variables that express pulse waveform: central aortic pressure (CAP), pulse pressure (PP), and augmentation index (AIx; ratio of the amplitude of the reflected wave in the ascending aorta and the PP).[37] CAP, PP, and AIx can be measured noninvasively using applanation tonometry (and Endo-PAT for AIx), calibrated by the peripheral diastolic and mean arterial pressure.[37] There is a lack of consensus regarding the optimal method to estimate the CAP with tonometry. The NIHem system assumes that carotid artery pulse waveform accurately reflects the central aortic waveform and the pulsed wave analysis is automatically calculated from the carotid waveform. The SphygmoCor device uses a generalized transform function to generate a central aortic PP curve from the radial or carotid pressure tracings, which has not been validated in children. Considering our largely pediatric group and need to maintain consistency between data acquired on each device, we use the nonprocessed, signal-averaged SphygmoCor carotid tracing as the central aortic tracing which will be then digitized to calculate the CAP, following previously published approach [Appendixes 2a and 3].[41]

### Blood pressure phenotype

BP phenotype is abnormal despite successful treatment of CoA. Office hypertension is a known risk factor for CV disease and the BP response during the ET is predictive of future development of resting hypertension in the general population.[42] Ambulatory blood pressure monitoring (ABPM) is superior to the office measurement in its ability to distinguish patients at the highest risk for target-organ damage.[43] We will assess BP phenotype with the manual auscultation technique to measure the right arm office BP; supine four extremity oscillometric BP measurement to assess for residual coarctation; ABPM to measure the circadian BP profile; and ET to assess the BP response to exercise and exercise-induced arm to leg BP gradient. Based on the office BP and ABPM results, we will classify our patients according to [Table 4](#) [Appendix 4].

### Biomarkers

We will measure asymmetric dimethylarginine (ADMA; NO's inhibitor),[44] and nitrite and nitrate (NO<sub>x</sub>, stable by-product of NO), biomarkers of endothelial function. Arterial stiffness is associated with increased systemic inflammation markers, which we will quantify with high-sensitivity C-reactive protein (hs-CRP) and local inflammatory cytokines of vascular wall function vascular adhesion molecule 1 (VCAM-1) and interleukin-1 beta (IL-1 $\beta$ ).[25,45] We will finally assess the molecular mechanisms of aortic wall response

to vascular dysfunction, with matrix metalloproteases (MMP-2 and MMP-9),[46] and transforming growth factor beta-1 (TGF- $\beta$ 1, a smooth cell growth-modulating factor involved in the arterial wall response to hypertension).[30] NO<sub>x</sub> will be determined by chemiluminescence (Sievers NOAnalyzer 280i) and all remaining measurements will be performed with enzyme-linked immunosorbent assay kits: ADMA (Sunred Biological Technology, Shanghai, China); hs-CRP (BoosterBio, Pleasanton, USA); VCAM-1; IL-1 $\beta$ ; matrix metalloproteases (MMP)-9; MMP-2; and TGF $\beta$ -1 (RayBiotech, Inc. Norcross, USA) [Appendix 5].

### Left ventricular mass

The altered BP phenotype that persists after CoA treatment represents an increase in afterload that leads to LV hypertrophy, strongly related to high BP and carrying a grave prognosis for cardiac events.[47] We will quantify LV mass by CMR, a well-established method for its calculation [Appendix 2b].

### Cardiovascular health assessment

Patients with CoA experience increased CV disease compared to the general population. Literature in the general population has demonstrated that risk of cardiometabolic disease and accelerated atherosclerosis is mitigated by ideal CV health (ICVH),[48] defined as having optimal levels of health factors (BP, total cholesterol, and plasma glucose) and behaviors (smoking, body mass index, physical activity, and diet). We will implement a questionnaire to assess family history of CV disease and ICVH according to the guidelines of the American Heart Association [Appendix 6].[48]

### Statistical considerations

**Adjustment for confounders** We will adjust our treatment groups for three main documented confounders: (a) age at treatment; (b) current age; and (c) bicuspid aortic valve (associated with impaired aortic elasticity).[49] During recruitment, we will attempt to frequency match the three treatment groups. During analysis, the treatment groups will be compared for each of these three confounding variables and adjustments will be made using multivariable modeling with linear and logistic regression models.

**Analytic plan** Our primary outcome variable will be cfPWV assessed by tonometry. Differences across groups will be explored using one-way analysis of variance. If differences in matching variables are detected among the groups, adjustment will be made using analysis of covariance. *Post hoc* analyses will be performed as necessary. Sample size estimates were obtained based on prior data that show that arch PWV measured by CMR is  $3.3 \pm 0.6$  m/s in normal patients and  $4.7 \pm 1.1$  m/sec after CoA surgery.[20,50] Sample size estimates for comparison of PVW between three equal-sized treatment groups (assuming overall significance level = 0.05 and power = 0.8) are shown in Table 5. We plan on recruiting 24–30 patients in each group for a total sample size of 72–90.

## DISCUSSION

Go to:

### Methodological considerations

We chose a multicenter design to overcome recruitment challenges secondary to restrictive enrollment criteria (particularly the lower treatment age limit of 1 year, which excludes a majority of CoA patients that present in infancy, mostly managed by surgery) and need for matching treatment groups for confounders.

cfPWV is our primary outcome variable because it is validated as an accurate and reproducible measure of arterial stiffness with proven association to hard CV outcomes that can be reliably measured by applanation tonometry and CMR. We chose other parameters to complete a complementary and comprehensive assessment of vascular function in small and large arteries.

### Importance of knowledge to be gained

This work will be the first systematic and comprehensive comparison of vascular function between three different treatment modalities in CoA patients. We postulate that the integrity of the arterial wall is best preserved with balloon dilatation, compared to stenting or surgery. We are aware that our population is

highly selected, but believe that this is the only way to compare the three treatment types. The results of our selected population may be relevant when several modalities are applicable to one patient. Currently, the preservation of vascular function is not considered when choosing between treatment modalities. Ultimately, the results of our study may help clinicians choose treatment modalities based not only on relief of anatomic stenosis but also on their ability to preserve long-term vascular health.

### **Study limitations**

Our results will reflect vascular function in a selected group of optimally treated CoA patients and may not be generalizable to all CoA patients. We will compare vascular function after treatment but not before the treatment. Variation in antihypertensive medication protocols between different institutions may affect vascular parameters.

### **CONCLUSION**

[Go to:](#)

There is ample evidence to suggest that CoA is a systemic arterial disease and not merely a focal stenosis of the aortic isthmus. However, the current management paradigm continues to focus on alleviating the anatomic stenosis. Our study aims to refine this treatment paradigm by adding the preservation of vascular function to the goals of successful treatment. The strengths of this study include its multicenter design and the use of multiple noninvasive modalities to perform a comprehensive and prospective assessment of vascular function and CV health.

### **Financial support and sponsorship**

The study was supported by two grants, from the Millennium Foundation BCP (awarded on December 18, 2012) and the Fundação Luso-Americana para o Desenvolvimento (Proj A-1 287/2012).

### **Conflicts of interest**

There are no conflicts of interest.

**APPENDICES: Appendix 1: Core laboratories, clinical sites, investigators** [Go to:](#)  
**for the LOVE-COARCT atudy, funding Study Principal Investigators**

---

Institution	Name
Department of Pediatric Cardiology Hospital de Santa Marta Centro Hospitalar de Lisboa Central Lisbon, PORTUGAL	José D. Martins, MD
Department of Cardiology Boston Children's Hospital Harvard Medical School, Boston, MA USA	Ashwin Prakash, MD
<b>Site Investigators of the LOVE Pediatric Consortium</b>	
Institution	Name
Department of Pediatric Cardiology Hospital de Santa Marta Centro Hospitalar de Lisboa Central Lisbon, PORTUGAL	José D. Martins, MD (Study PI, Site PI) Fátima F. Pinto, MD, PhD (Study Advisor) Marta António, MD Lurdes Medroa Teresa Mateus Mafalda Selas, RN
Department of Cardiology Boston Children's Hospital Harvard Medical School Boston, USA	Ashwin Prakash, MD (Study PI, Site PI) Sarah de Ferranti, MD Kimberlee Gauvreau, ScD Tal Geva, MD Jonathan Rhodes, MD Cara Hass, BS (Study Coordinator) Jeffrey Reichman (Study Coordinator) James E. Lock, MD (Study Advisor) Jane Newburger, MD, MPH (Study Advisor) Shelby Kutty, MD (Site PI)
Joint Division of Pediatric Cardiology Children's Hospital and Medical Center University of Nebraska College of Medicine Omaha, USA Division of Pediatric Cardiology Department of Pediatrics Lucile Packard Children's Hospital Stanford University Palo Alto, USA	Elif Seda Selamet Tierney, MD (Site PI) Angela Chen
Division of Pediatric Cardiology Children's Hospital Colorado Aurora, USA	Uyen Truong (Site PI)
Serviço de Cardiologia Pediátrica Hospital Pediátrico de Coimbra Coimbra, PORTUGAL	António Marinho, MD (Site PI) Eduardo Castela, MD
Division of Pediatric Cardiology Texas Children's Hospital Baylor College of Medicine Houston, USA	Shaine Morris, MD (Site PI) Justin Zachariah, MD
CEDOC Chronic Diseases Nova Medical School Lisbon PORTUGAL Ressonância Magnética, S.A. Lisbon, Portugal	Miguel Mota Carmo, MD PhD (Director) Maria Guarino, MD PhD  Nuno Jalles Tavares (Director) Marta António, MD Boban Thomas, MD Diana Cruz Oliveira, MPH
Biomedical Engineering Department Instituto Superior Técnico Lisbon, Portugal	
<b>Core Laboratories</b>	
Core Lab	Institution
Cardiac Magnetic Resonance (Ashwin Prakash) Preventive Cardiology (Sarah de Ferranti) Biostatistics (Kimberlee Gauvreau)	Department of Cardiology Boston Children's Hospital Harvard Medical School Boston, USA
<b>Core Laboratories</b>	
Institution	Name
Tonometry and BP Assessment (Justin Zachariah)	Division of Pediatric Cardiology Texas Children's Hospital Baylor College of Medicine Houston, USA
Biomarkers (Maria Guarino)	CEDOC Chronic Diseases Nova Medical School Lisbon PORTUGAL
Endothelial Function (Elif Seda Selamet Tierney)	Division of Pediatric Cardiology Department of Pediatrics Lucile Packard Children's Hospital Stanford University Palo Alto, USA
<b>Data Coordination</b>	
Data Coordination (Jose Martins, Ashwin Prakash, Cara Hass, Jeffrey Reichman)	Department of Pediatric Cardiology Hospital de Santa Marta Centro Hospitalar de Lisboa Central Lisbon, PORTUGAL and Department of Cardiology Children's Hospital Boston Harvard Medical School Boston, USA

## Funding

This study was made possible with grants from the Millennium Foundation BCP (signed Dec 18, 2012) and the Fundação Luso-Americana para o Desenvolvimento (Proj A-1 287/2012).

## Appendix 2a: Applanation tonometry manual of operations

[Go to:](#)

For applanation tonometry, some centers will use the NIHem system (Cardiovascular Engineering, Inc.,

Norwood, MA USA) and others the SphygmoCor device (AtCor Medical, West Ryde, NSW, Australia). The technology is similar and the results comparable.

The patient demographics and brachial blood pressure (BP) are entered into the system. First, the tonometer is placed over the right carotid artery, just lateral to the thyroid cartilage. The location is adjusted and pressure applied as needed to optimize waveform. After ensuring that the tracings are optimal, the tracing is recorded. The carotid site is marked. Then, the tonometer is placed over the right femoral artery and the same process for obtaining an optimal curve recording is followed. The femoral site is marked. Finally, in the centers that use the SphygmoCor device, a third recording of the radial artery is performed, in the same fashion. A caliper is used to measure the distance from the suprasternal notch to the carotid site and from the suprasternal notch to the femoral site. Both distances are entered in the system.

For pulse wave velocity and augmentation index calculation, both systems analyze the curves and supply the data with the proprietary software package, without any input from the examiner.

For pulse wave analysis (central aortic pressure and pulse pressure), the analysis procedure differed slightly between systems. The analysis from the NIHEM system is done by the system's software. In the centers that used SphygmoCor, the signal averaged carotid pulse wave is digitalized and calibrated according to a published approach:[41,51] The brachial diastolic and mean pressures are used and the same diastolic and mean pressures are assigned to the averaged carotid pulse. Moreover, the radial pressure waveform is used to retrieve the correspondent time instants of diastolic and mean pressures. Given the two pressure values and the correspondent time instants, it is possible to calibrate each averaged carotid pressure waveform. This process allows a quantitative analysis of the pulse waveform.

## **Appendix 2b: Cardiac magnetic resonance imaging manual of operations**

Cardiac magnetic resonance (CMR) will be performed using commercially available whole-body 1.5 T scanners (Achieva; Philips Healthcare, Best, the Netherlands; Signa 1.5T or GE Medical Systems, Milwaukee, WI, USA). Electrocardiography (ECG)-gated steady-state free precession (SSFP) localizers will be used in sagittal, coronal, and axial planes during free breathing. Ventricular function will be assessed from short-axis stack to cover ventricles from base to apex, acquired using the following imaging parameters: slice thickness 5–8 mm, slice gap 0–1 mm, slice number 12–14, cardiac phases 30, retrospective gating with breath-holding. In patients unable to breath-hold 3 signal averages during free-breathing will be used. SSFP cine imaging will also be performed in two orthogonal long-axis planes of the left ventricular outflow tract (during breath-hold), short axis of the ascending aorta (AAO), and in the long axis of the aortic arch (free-breathing, used as reference for pulse wave velocity measurements), proximal descending aorta (DAO, 2–3 cm distal to the isthmus, sufficiently distal to dephasing jets), mid-DAO (diaphragmatic level), and distal DAO (just above iliac bifurcation). ECG-gated through-plane phase-contrast flow measurements will be performed at the AAO (5 mm distal to the sinotubular junction) and in proximal-, mid-, and distal-DAO segments (matched to location of the cine SSFP acquisitions) using the following imaging parameters: signal averages = 2, cardiac phases 100 (TFE factor/views per segment/ = 1 [to maximize temporal resolution]), and velocity encoding 200–250 cm/s (higher if needed to avoid aliasing). ECG and respiratory navigator-gated three-dimensional SSFP MRA of the aortic arch will be performed in the sagittal plane.

The patient's right arm BP while on scanner table and length of time since last meal and content of last meal will be recorded. Images will be analyzed by a single observer (A.P.) in the CMR core lab using a commercial computer workstation (Extended Workstation; Philips Healthcare) and using commercially available analysis software (QMass and QFlow, Medis, The Netherlands). Ventricular function and mass will be calculated using standard techniques. Cross-sectional areas of the AAO and proximal, mid, and distal DAO will be directly planimeted at peak systole and mid-diastolic frames to calculate parameters of segmental aortic stiffness as previously described.[52] Pulse wave velocity will be measured using the transit-time method.[20] Pulse wave velocity will be calculated for the entire aorta (AAO to distal DAO), as well as in the following segments: AAO to proximal DAO, proximal DAO to mid-DAO, and mid-DAO to distal DAO. Aortic arch shape will be classified and the aortic arch index calculated as previously described.[53]

### **Appendix 3: Endothelial pulse amplitude testing manual of operations** [Go to:](#)

---

The testing room will be arranged to provide a quiet, restful environment with a comfortable temperature of 22°C to 23°C. Before testing, patients will be asked to fast overnight for 12 h, except for the consumption of water. Unless the patients are taking a daily vitamin, they will be asked to refrain from taking vitamin pills and over-the-counter medications; in the case that an over-the-counter medication is used, it will be documented.

The Endo-PAT (Itamar Medical Ltd, Caesarea, Israel) testing protocol[40] will be performed in the morning (starting time between 8 and 11 am) and fasting. Any restrictive clothing that could interfere with blood flow to the arms or fingers will be removed, including heavy coats or clothes with thick sleeves, watches or rings or other jewelry on the hands and fingers, and long fingernails shortened with a fingernail clipper.

Noninvasive pneumatic probes will be placed on the index fingers of both hands. The pulse wave amplitude will be recorded continuously from both index fingers. Reactive hyperemia will be performed by achieved by occlusion of the brachial artery of one arm with a BP cuff for 5 min (to 200–220 mmHg). The tracing in the nonoccluded arm will serve as a control for changes in overall physiologic state. The Endo-PAT data will be analyzed with the proprietary software package, without any input from the examiner. The Endo-PAT index is defined as the ratio of the average pulse amplitude during the 1 minute period beginning after exactly 90 s of reactive hyperemia compared with the average pulse amplitude during the 210-s preocclusion baseline period.

### **Appendix 4a: Right arm, auscultatory blood pressures measurement manual of operations** [Go to:](#)

---

The patient will be seated with the feet flat on the floor, with the knees at 90° and the back supported. After 5 min of resting quietly, with no conversation or television, the auscultatory BP will be obtained in the right arm. For cuff choice, the length of the bladder encircled no <80% and no more than 100%, of the bicep and the width of bladder encircled no <40% and no more than 50%, of the circumference of patient's arm circumference, measured at the widest area of bicep, midway between the tip of the patient's shoulder and the tip of the patient's elbow. The patient's right arm will be placed at heart level, supported at the level of the nipple by resting arm on a table or chair arm or propped on a pillow.

The stethoscope's bell will be placed over patient's brachial pulse. The cuff will be inflated up to 140 mmHg and deflated slowly while listening for the Korotkoff sounds, systole being number when the sound is first heard consistently and diastole when the last pulsation is heard or when it muffles. If pulsations are immediately audible, the cuff will be deflated entirely and the patient allowed to sit quietly for 1 minute. Then, the cuff will be again inflated to 160 mmHg (or higher) and the steps above will be followed. This procedure will be repeated until the blood pressure (BP) is not immediately audible.

Three BPs will be obtained, allowing 1 min between deflation and reinflation of cuff for each measurement. The average of the 2<sup>nd</sup> and 3<sup>rd</sup> measurements will be considered the final right arm BP and interpreted according to the published guidelines for children[54] and adults.[55]

### **Appendix 4b: Four extremity, automated blood pressures (Dinamap) measurement manual of operations**

While the patient is supine, two sets of four extremity blood pressure (BP) will be measured, with the automated BP monitor (Dinamap).

The BP pressure gradient will be registered, between the second systolic right arm measurement and the highest of the two legs systolic second measurements. In the presence of an aberrant right subclavian artery that originates distal to the Coarctation of the aorta site, seen by cardiovascular magnetic resonance, we will use the second left systolic arm measurement for the residual gradient.

### **Appendix 4c: Ambulatory blood pressure (BP) monitor methods of operations**

The patient data will be recorded. The choice of the cuff will follow the same guidelines described for manual auscultation of right arm BP. Cuff inflation will be programmed for 15–20-min intervals. During

nighttime, intervals are wider, but not fewer than one per hour and preferably more. The patient will record the sleep time, wake time, and any periods of vigorous exercise. The patient will be instructed to avoid direct contact of the monitor with water and participation in activities that could damage it.

The study will be considered adequate if there is a record of at least 1 reading per hour, i.e., no more than 1 h between consecutive readings for a full 24-h study. If less than 12 h are recorded, the ambulatory blood pressure monitoring data will be considered inadequate. Diurnal pattern will be determined by the patient diary. Vigorous exercise periods will be excluded.

The data on 24-h systolic BP load, 24-h diastolic BP load, diurnal systolic dipping, diastolic dipping and 24 h, daytime and nighttime mean systolic BP, and mean diastolic BP will be recorded. Patients will be staged as having ambulatory hypertension, masked hypertension, white coat hypertension or normotensive, according to the age-based normative tables based on statements for children and adolescents[43] and adults.[56] Patients currently on antihypertensive medication are also classified into the hypertensive group [Table 4].

#### **Appendix 4d: Exercise test: Manual of operations**

The patient information will be entered per equipment specification and the study identifier on the datasheet and the date of the test. For patient safety issues, medical history, medications, activity level, and symptoms will be reviewed and the exercise stress test protocol will be explained. Antihypertensive medications will be continued the day of testing.

The patient will be asked to lay supine, and a right arm and right or left leg blood pressure measured using a commercial oscillometric and appropriate sized cuff bladders and recorded as preexercise blood pressure (BP) values and gradient. The patient then will step onto the treadmill and instructed to hold the handlebar throughout the test. We will use the standard Bruce treadmill protocol and, when available, a Met Cart. As the patient exercises, their symptoms and electrocardiography (ECG) will be continuously monitored. At 2-min of each stage, a BP will be taken in the right arm by having the patient take their hand off the treadmill and hold onto the arm of the person performing the test. The test will be terminated when the patient can no longer continue the exercise, reaches a systolic BP higher than 240 mmHg, has clinically relevant symptoms or ECG changes. Immediately after the exercise ended, BP in the right arm and the left leg will be recorded in a supine position. For the recovery period, the patient will sit upright in a chair, and right arm BP will be recorded at 1, 3, 5, and 7 min of recovery, at which time the test is ended.

The data on exercise duration, baseline and exercise right arm BP, pre- and post-exercise systolic BP gradient, patient symptoms, ECG changes and, when available, cardiorespiratory physiological data will be documented. We will label exercise-induced hypertension when the systolic BP is  $\geq 220$  mmHg.

#### **Appendix 5: Biomarkers manual of operations**

[Go to:](#)

The patients will follow a low-nitrate diet for 3 days before the blood sample collection, which avoids of a list of foods with a high content in nitrites that influence nitric oxide determination, including bacon, beets, broccoli, canned food, cauliflower, celery, Chinese cabbage, corned beef, ham, hot dogs, lettuce, old cheese, radish, salami, sausages, smoked fish, spinach, and turnip. After an overnight fast (for 12 h), samples will be collected by venipuncture from catheters maintained with saline only, since heparin interferes with accuracy of the biomarkers assessed. The first 5–10 mL of blood will be discarded and 2.7 ml of venous blood will be collected into 3.2% sodium citrate (light-blue) tubes (BD Vacutainer<sup>®</sup>), and into plastic microtubes (Safe-Lock Eppendorf). Within 3 h of collection, samples will be centrifuged for 20 minutes at 3000g (4°C). Aliquots of 250  $\mu$ l of the supernatant will be collected into 14 labeled microtubes of 1.5 ml and immediately stored at  $-80^{\circ}\text{C}$  until shipping to the Biomarkers Core Laboratory.

Aliquots for NO<sub>x</sub> analysis will be deproteinized using cold ethanol precipitation methodology. Ethanol will be refrigerated to 0°C and added to the plasma sample in a 1:3 proportion. After letting it stand at 0°C for 30 min, the sample will be centrifuged at 14,000 rpm for 10 min. The supernatant will be then removed for analysis. The quantification of plasma NO levels will be carried out using a nitric oxide analyzer, the Sievers Instruments NOA 280i<sup>™</sup>, a high sensitivity detector of that allows determination of NO based on a

chemiluminescence reaction between NO and ozone.

Plasma asymmetric dimethylarginine (ADMA); vascular cell adhesion molecule 1 (VCAM-1); high-sensitivity C-reactive protein (hs-CRP) interleukin-1-beta (IL-1 $\beta$ ); MMP-2 and MMP-9 will be quantified using the following double-antibody sandwich enzyme-linked immunosorbent assay ELISA kits: Human asymmetrical dimethylarginine, ADMA (Sunred Biological Technology, Shanghai, China); hs-CRP (BoosterBio, Pleasanton, USA); VCAM-1; IL-1 $\beta$ ; MMP-9/Gelatinase A; MMP-2/Gelatinase B; and transforming growth factor beta (RayBiotech, Inc. Norcross, USA).

## **Appendix 6: Cardiovascular health assessment manual of operations**

[Go to:](#)

The following questionnaires will be used.

Lifestyle questionnaire:

- On an average weekday, how many hours do you watch TV?
- On an average weekday, how many hours do you play video/computer games or use a computer for something that is not school/work related?
- In the past week, how many days were you/was your child physically active for a total of at least 30 min/day?
- In the past week, how many days did you/your child eat breakfast? In the past week, how many days did you/your child eat food from a fast food restaurant?
- In the past week, how many days did all or most of your family sit down and eat dinner at home?
- On an average weekday, how many hours of sleep do you get a night?
- Have you smoked one or more cigarettes in the past month? If yes, please quantify.
- Were you previously a smoker?
- Do you live in a household with a smoker?

Family history questionnaire:

For all the following questions, the possible answers will be “no,” “parents/siblings,” “grandparents/aunts/uncles,” and “both”

- Biological relatives of you/your child with overweight/obesity
- Biological relatives of you/your child with type 2 diabetes
- Biological relatives of you/your child with high blood pressure
- Biological relatives of you/your child with high cholesterol
- Biological relatives of you/your child with heart disease/stroke
- All answers had the following options: Parents/siblings/grandparents/aunts/uncles.

## **Appendix 7. Comprehensive List of Study Variables**

[Go to:](#)

	Variables	Comments or Definitions
Medical History	BSA at Initial Echocardiogram	using Haycock's Formula; m <sup>2</sup>
	Minimum Transverse Arch Diameter Z-score on Initial Echo	Calculated with Boston z-scores
	Isthmus z score on Initial Echo	Calculated with Boston z-scores
	Initial Doppler coarctation gradient	mmHg
	Bicuspid/Bicommissural Aortic Valve?	Yes/No
	Initial arm-leg systolic BP gradient	mmHg
	Type of Initial Treatment	Balloon/Stent/Surgery
Local blood results	Currently daily medications?	Yes/No. If yes, please specify.
	Total Cholesterol	mg/dL
	LDL	mg/dL
	HDL	mg/dL
	Triglycerides	mg/dL
	Plasma Glucose	mg/dL
Applanation tonometry	Insulin	uIU/mL
	Hemoglobin A1C	%
	Central Systolic Blood Pressure	mmHg
	Central Pulse Pressure	mmHg
	Heart Rate	bpm
	Carotid Femoral PWV	meters/second
	Augmentation Index (%)	%
CMR	Augmentation Index at HR75	%
	LV End-Diastolic Volume indexed to BSA	ml/m <sup>2</sup>
	LV End-Systolic Volume indexed to BSA	ml/m <sup>2</sup>
	LV Ejection Fraction	%
	LV Mass indexed to BSA	g/m <sup>2</sup>
	Ascending Ao - Descending Ao PWV (Ascending Ao to proximal, mid and distal descending Ao)	$\frac{\text{Distance (Asc Ao to Desc Ao)}}{\text{Time Delay (Asc Ao to Desc Ao)}}$
	Type of arch	Romanesque; Gothic; Crenel
	Aortic diameter (Ascending, Proximal, Mid and Distal Ao)	mm/mm <sup>2</sup>
	Aortic strain (Ascending, Proximal, Mid and Distal Ao)	$\frac{\text{Sistolic Area} - \text{Diastolic Area}}{\text{Diastolic Area}}$
	Aortic compliance (Ascending, Proximal, Mid and Distal Ao)	$\frac{\text{Ao Area Sist} - \text{Ao Area Diast}}{\text{SBP} - \text{DBP}}$ cm <sup>2</sup> /mmHg
	Aortic Distensibility (Ascending, Proximal, Mid and Distal Ao)	$\frac{\text{Ao strain}}{\text{SBP} - \text{DBP}}$ mmHg <sup>-1</sup>
	Aortic stiffness β index (Ascending, Proximal, Mid and Distal Ao)	$\frac{\ln(\text{SBP} / \text{DBP})}{\text{Strain}}$
	Loss of pulse amplitude	$100 \times \frac{\text{flow (AscAo - DescAo)}}{\text{flow (AscAo)}}$
	Aorta Young's modulus (Ascending, Proximal, Mid and Distal Ao)	$\frac{(\text{SBP} - \text{DBP})\text{Ao diameter diastole}}{\text{Ao diameter (systole - diastole)}\text{Ao wall thickness}}$
	Arterial elastance (Ea)	$\frac{\text{End Systolic Pressure}}{\text{Stroke Volume}}$ mmHg/ml
LV end-systolic elastance (Ees)	$\frac{\text{End Systolic Pressure}}{\text{LV end Systolic Volume}}$ mmHg/ml	
Endo-PAT	Reactive Hyperemia Index (RHI)	%
	Augmentation Index	%
	Augmentation Index at 75 bpm	%
BP during the patient's visit	Residual SBP gradient (between right arm and highest of the legs)	Supine. Automated. Two sets of measurements; mmHg
	Right arm BP	Seated. Manual. Three sets of measurements; mmHg
ABPM	24 Average Systolic BP	mmHg
	24 Hour Average Diastolic BP	mmHg
	Daytime Average Systolic BP	mmHg
	Daytime Average Diastolic BP	mmHg
	Nighttime Average Systolic BP	mmHg
	Nighttime Average Diastolic BP	mmHg
	24h systolic load	%
	24h diastolic load	%
	Diurnal Systolic Variation	%
	Diurnal Diastolic Variation	%
Exercise test	Exercise Duration	Minutes
	Pre-exercise right arm BP	mmHg
	Pre-exercise leg BP	mmHg
	Pre-Exercise SBP gradient	mmHg
	Post-exercise right arm BP	mmHg
	Post-exercise leg BP	mmHg
	Pre-Exercise SBP gradient	mmHg
Biomarkers	Peak exercise BP	mmHg
	NOx	ug/ml
	ADMA	ng/L
	High Sensitivity CRP	mg/L
	VCAM-1	ng/ml
	IL-1beta	pg/ml
	TFG-Beta	ng/ml
MMP-2/Gelatinase A	ng/ml	
MMP-9/Gelatinase B	ng/ml	

Legend: ADMA = Asymmetric Dimethylarginine; Ao = Aorta; BP = blood pressure; BP = Blood Pressure; BSA= Body Surface Area; GMR = Cardiac magnetic resonance; DBP = Diastolic Blood Pressure; HDL = High-density lipoprotein cholesterol; Hs-CRP = High sensitivity C-Reactive Protein; IL-1β = Interleukin 1 beta; LDL = Low-density lipoprotein cholesterol; MMP-2 = Matrix Metalloproteinase-2; MMP-9 = Matrix Metalloproteinase-9; NOx = Nitric Oxide; PWV = Pulse Wave Velocity; LV = Left Ventricle; SBP = Systolic Blood Pressure; TFG-β = Transforming Growth Factor beta; VCAM-1 = Vascular Cell Adhesion Molecule 1; † using previously published normative data.<sup>67†</sup>

**REFERENCES**[Go to:](#)


1. Forbes TJ, Kim DW, Du W, Turner DR, Holzer R, Amin Z, et al. Comparison of surgical, stent, and balloon angioplasty treatment of native coarctation of the aorta: An observational study by the CCISC (Congenital Cardiovascular Interventional Study Consortium) *J Am Coll Cardiol*. 2011;58:2664–74. [PubMed: 22152954]
2. Brown ML, Burkhart HM, Connolly HM, Dearani JA, Cetta F, Li Z, et al. Coarctation of the aorta: Lifelong surveillance is mandatory following surgical repair. *J Am Coll Cardiol*. 2013;62:1020–5. [PubMed: 23850909]
3. Martins JD, Thomas B, Jalles N, Pinto FF. Aortic arch geometry in repaired coarctation: Systematic review of a consecutive series by magnetic resonance imaging. *Rev Port Cardiol*. 2012;31:403–4. [PubMed: 22494835]
4. Peres A, Martins JD, Paramés F, Gil R, Matias C, Franco J, et al. Isolated aortic coarctation: Experience in 100 consecutive patients. *Rev Port Cardiol*. 2010;29:23–35. [PubMed: 20391897]
5. Hager A, Kanz S, Kaemmerer H, Schreiber C, Hess J. Coarctation long-term assessment (COALA): Significance of arterial hypertension in a cohort of 404 patients up to 27 years after surgical repair of isolated coarctation of the aorta, even in the absence of restenosis and prosthetic material. *J Thorac Cardiovasc Surg*. 2007;134:738–45. [PubMed: 17723827]
6. Quennelle S, Powell AJ, Geva T, Prakash A. Persistent aortic arch hypoplasia after coarctation treatment is associated with late systemic hypertension. *J Am Heart Assoc*. 2015;4:pii: e001978. [PMCID: PMC4608082]
7. Bocelli A, Favilli S, Pollini I, Bini RM, Ballo P, Chiappa E, et al. Prevalence and long-term predictors of left ventricular hypertrophy, late hypertension, and hypertensive response to exercise after successful aortic coarctation repair. *Pediatr Cardiol*. 2013;34:620–9. [PubMed: 23052661]
8. Meyer AA, Joharchi MS, Kundt G, Schuff-Werner P, Steinhoff G, Kienast W, et al. Predicting the risk of early atherosclerotic disease development in children after repair of aortic coarctation. *Eur Heart J*. 2005;26:617–22. [PubMed: 15618050]
9. Pedersen TA, Pedersen EB, Munk K, Hjortdal VE, Emmertsen K, Andersen NH, et al. High pulse pressure is not associated with abnormal activation of the renin-angiotensin-aldosterone system in repaired aortic coarctation. *J Hum Hypertens*. 2015;29:268–73. [PubMed: 25355011]
10. Chen SS, Donald AE, Storry C, Halcox JP, Bonhoeffer P, Deanfield JE, et al. Impact of aortic stenting on peripheral vascular function and daytime systolic blood pressure in adult coarctation. *Heart*. 2008;94:919–24. [PubMed: 17686804]
11. Toro-Salazar OH, Steinberger J, Thomas W, Rocchini AP, Carpenter B, Moller JH, et al. Long-term follow-up of patients after coarctation of the aorta repair. *Am J Cardiol*. 2002;89:541–7. [PubMed: 11867038]
12. Koller M, Rothlin M, Senning A. Coarctation of the aorta: Review of 362 operated patients. Long-term follow-up and assessment of prognostic variables. *Eur Heart J*. 1987;8:670–9. [PubMed: 3653118]
13. Choudhary P, Canniffe C, Jackson DJ, Tanous D, Walsh K, Celermajer DS, et al. Late outcomes in adults with coarctation of the aorta. *Heart*. 2015;101:1190–5. [PubMed: 25810155]
14. Hoimyr H, Christensen TD, Emmertsen K, Johnsen SP, Riis A, Hansen OK, et al. Surgical repair of coarctation of the aorta: Up to 40 years of follow-up. *Eur J Cardiothorac Surg*. 2006;30:910–6. [PubMed: 17056267]
15. Mandalenakis Z, Rosengren A, Lappas G, Eriksson P, Hansson PO, Dellborg M, et al. Ischemic stroke in children and young adults with congenital heart disease. *J Am Heart Assoc*. 2016;5:pii: e003071. [PMCID: PMC4802444]

16. Vogt M, Kühn A, Baumgartner D, Baumgartner C, Busch R, Kostolny M, et al. Impaired elastic properties of the ascending aorta in newborns before and early after successful coarctation repair: Proof of a systemic vascular disease of the prestenotic arteries? *Circulation*. 2005;111:3269–73. [PubMed: 15956120]
17. Li VW, Cheung YF. Arterial-left ventricular-left atrial coupling late after repair of aortic coarctation and interruption. *Eur Heart J Cardiovasc Imaging*. 2015;16:771–80. [PubMed: 25588795]
18. Agnoletti G, Bonnet C, Bonnet D, Sidi D, Aggoun Y. Mid-term effects of implanting stents for relief of aortic recoarctation on systemic hypertension, carotid mechanical properties, intimal medial thickness and reflection of the pulse wave. *Cardiol Young*. 2005;15:245–50. [PubMed: 15865825]
19. Lam YY, Mullen MJ, Kaya MG, Gatzoulis MA, Li W, Henein MY, et al. Left ventricular long axis dysfunction in adults with “corrected” aortic coarctation is related to an older age at intervention and increased aortic stiffness. *Heart*. 2009;95:733–9. [PubMed: 19095712]
20. Ou P, Celermajer DS, Jolivet O, Buyens F, Herment A, Sidi D, et al. Increased central aortic stiffness and left ventricular mass in normotensive young subjects after successful coarctation repair. *Am Heart J*. 2008;155:187–93. [PubMed: 18082512]
21. Shang Q, Sarikouch S, Patel S, Schuster A, Steinmetz M, Ou P, et al. Assessment of ventriculo-vascular properties in repaired coarctation using cardiac magnetic resonance-derived aortic, left atrial and left ventricular strain. *Eur Radiol*. 2017;27:167–77. [PubMed: 27215582]
22. de Divitiis M, Pilla C, Kattenhorn M, Zadinello M, Donald A, Leeson P, et al. Vascular dysfunction after repair of coarctation of the aorta: Impact of early surgery. *Circulation*. 2001;104:1165–70. [PubMed: 11568050]
23. Trojnarska O, Szczepaniak-Chicheł L, Mizia-Stec K, Gabriel M, Bartczak A, Grajek S, et al. Vascular remodeling in adults after coarctation repair: Impact of descending aorta stenosis and age at surgery. *Clin Res Cardiol*. 2011;100:447–55. [PMCID: PMC3079825] [PubMed: 21161708]
24. Mizia-Stec K, Trojnarska O, Szczepaniak-Chicheł L, Gabriel M, Bartczak A, Cieplucha A, et al. Asymmetric dimethylarginine and vascular indices of atherosclerosis in patients after coarctation of aorta repair. *Int J Cardiol*. 2012;158:364–9. [PubMed: 21334083]
25. Brili S, Tousoulis D, Antoniadis C, Aggeli C, Roubelakis A, Papathanasiu S, et al. Evidence of vascular dysfunction in young patients with successfully repaired coarctation of aorta. *Atherosclerosis*. 2005;182:97–103. [PubMed: 16115479]
26. Jesus CA, Assef JE, Pedra SR, Ferreira WP, Davoglio TA, Petisco AC, et al. Serial assessment of arterial structure and function in patients with coarctation of the aorta undergoing stenting. *Int J Cardiovasc Imaging*. 2016;32:729–39. [PubMed: 26723574]
27. Moutafi AC, Alissafi T, Chamakou A, Chryssanthopoulos S, Thanopoulos V, Dellos C, et al. Neurohormonal activity and vascular properties late after aortic coarctation repair. *Int J Cardiol*. 2012;159:211–6. [PubMed: 21429604]
28. Swan L, Kraidly M, Vonder Muhll I, Collins P, Gatzoulis MA. Surveillance of cardiovascular risk in the normotensive patient with repaired aortic coarctation. *Int J Cardiol*. 2010;139:283–8. [PMCID: PMC2849010] [PubMed: 19059656]
29. Brili S, Antonopoulos AS, Oikonomou E, Kalampogias A, Papamikroulis GA, Chrysochoou C, et al. Impairment of arterial elastic properties and elevated circulating levels of transforming growth factor-beta in subjects with repaired coarctation of aorta. *Int J Cardiol*. 2016;207:282–3. [PubMed: 26812641]
30. Xu C, Lee S, Singh TM, Sho E, Li X, Sho M, et al. Molecular mechanisms of aortic wall remodeling in response to hypertension. *J Vasc Surg*. 2001;33:570–8. [PubMed: 11241129]
31. Sarkola T, Redington AN, Slorach C, Hui W, Bradley T, Jaeggi E, et al. Assessment of vascular phenotype using a novel very-high-resolution ultrasound technique in adolescents after aortic coarctation

- repair and/or stent implantation: Relationship to central haemodynamics and left ventricular mass. *Heart*. 2011;97:1788–93. [PubMed: 21795301]
32. Kutty S, Rangamani S, Venkataraman J, Li L, Schuster A, Fletcher SE, et al. Reduced global longitudinal and radial strain with normal left ventricular ejection fraction late after effective repair of aortic coarctation: A CMR feature tracking study. *Int J Cardiovasc Imaging*. 2013;29:141–50. [PubMed: 22581073]
33. Martins JD, Franco J, Freitas I, Trigo C, Pinto FF, Kaku S. Blood pressure response to exercise: A new paradigm of success of coarctation of the aorta treatment? *Rev Port Cardiol*. 2008;27:122.
34. Egan M, Holzer RJ. Comparing balloon angioplasty, stenting and surgery in the treatment of aortic coarctation. *Expert Rev Cardiovasc Ther*. 2009;7:1401–12. [PubMed: 19900023]
35. Vlachopoulos C, Aznaouridis K, Stefanadis C. Prediction of cardiovascular events and all-cause mortality with arterial stiffness: A systematic review and meta-analysis. *J Am Coll Cardiol*. 2010;55:1318–27. [PubMed: 20338492]
36. Harris PA, Taylor R, Thielke R, Payne J, Gonzalez N, Conde JG, et al. Research electronic data capture (REDCap) – A metadata-driven methodology and workflow process for providing translational research informatics support. *J Biomed Inform*. 2009;42:377–81. [PMCID: PMC2700030] [PubMed: 18929686]
37. Townsend RR, Wilkinson IB, Schiffrin EL, Avolio AP, Chirinos JA, Cockcroft JR, et al. Recommendations for improving and standardizing vascular research on arterial stiffness: A scientific statement from the American Heart Association. *Hypertension*. 2015;66:698–722. [PMCID: PMC4587661] [PubMed: 26160955]
38. Grotenhuis HB, Westenberg JJ, Steendijk P, van der Geest RJ, Ottenkamp J, Bax JJ, et al. Validation and reproducibility of aortic pulse wave velocity as assessed with velocity-encoded MRI. *J Magn Reson Imaging*. 2009;30:521–6. [PubMed: 19711407]
39. Lerman A, Zeiher AM. Endothelial function: Cardiac events. *Circulation*. 2005;111:363–8. [PubMed: 15668353]
40. Selamet Tierney ES, Newburger JW, Gauvreau K, Geva J, Coogan E, Colan SD, et al. Endothelial pulse amplitude testing: Feasibility and reproducibility in adolescents. *J Pediatr*. 2009;154:901–5. [PubMed: 19217124]
41. Kelly R, Fitchett D. Noninvasive determination of aortic input impedance and external left ventricular power output: A validation and repeatability study of a new technique. *J Am Coll Cardiol*. 1992;20:952–63. [PubMed: 1527307]
42. Singh JP, Larson MG, Manolio TA, O'Donnell CJ, Lauer M, Evans JC, et al. Blood pressure response during treadmill testing as a risk factor for new-onset hypertension. The framingham heart study. *Circulation*. 1999;99:1831–6. [PubMed: 10199879]
43. Flynn JT, Daniels SR, Hayman LL, Maahs DM, McCrindle BW, Mitsnefes M, et al. Update: Ambulatory blood pressure monitoring in children and adolescents: A scientific statement from the American Heart Association. *Hypertension*. 2014;63:1116–35. [PMCID: PMC4146525] [PubMed: 24591341]
44. Cooke JP. Asymmetrical dimethylarginine: The uber marker? *Circulation*. 2004;109:1813–8. [PubMed: 15096461]
45. Barbaro NR, Fontana V, Modolo R, De Faria AP, Sabbatini AR, Fonseca FH, et al. Increased arterial stiffness in resistant hypertension is associated with inflammatory biomarkers. *Blood Press*. 2015;24:7–13. [PubMed: 25061978]
46. Takagi H, Manabe H, Kawai N, Goto SN, Umemoto T. Circulating matrix metalloproteinase-9 concentrations and abdominal aortic aneurysm presence: A meta-analysis. *Interact Cardiovasc Thorac Surg*. 2009;9:437–40. [PubMed: 19525292]


47. Kannel WB, Gordon T, Offutt D. Left ventricular hypertrophy by electrocardiogram. Prevalence, incidence, and mortality in the framingham study. *Ann Intern Med.* 1969;71:89–105. [PubMed: 4239887]
48. Lloyd-Jones DM, Hong Y, Labarthe D, Mozaffarian D, Appel LJ, Van Horn L, et al. Defining and setting national goals for cardiovascular health promotion and disease reduction: The American Heart Association's strategic impact goal through 2020 and beyond. *Circulation.* 2010;121:586–613. [PubMed: 20089546]
49. Nistri S, Grande-Allen J, Noale M, Basso C, Siviero P, Maggi S, et al. Aortic elasticity and size in bicuspid aortic valve syndrome. *Eur Heart J.* 2008;29:472–9. [PubMed: 18096569]
50. Walhout RJ. Universiteit Amsterdam; 2009. Assessment of proximal and distal aortic properties with magnetic resonance after successful coarctation management in adults. *Advances in the Management and Surveillance of Patients with Aortic Coarctation* 26 June, 2009; pp. 122–30.
51. Quaal SJ. Interpreting the arterial pressure waveform in the intra-aortic balloon-pumped patient. *Prog Cardiovasc Nurs.* 2001;16:116–8. [PubMed: 11464433]
52. Prakash A, Adlakha H, Rabideau N, Hass CJ, Morris SA, Geva T, et al. Segmental Aortic Stiffness in Children and Young Adults With Connective Tissue Disorders: Relationships With Age, Aortic Size, Rate of Dilation, and Surgical Root Replacement. *Circulation.* 2015;132:595–602. [PubMed: 26115544]
53. Ou P, Bonnet D, Auriacombe L, Pedroni E, Balleux F, Sidi D, Mousseaux E. Late systemic hypertension and aortic arch geometry after successful repair of coarctation of the aorta. *Eur Heart J.* 2004;25:1853–9. [PubMed: 15474701]
54. National High Blood Pressure Education Program Working Group on High Blood Pressure in C and Adolescents. The fourth report on the diagnosis, evaluation, and treatment of high blood pressure in children and adolescents. *Pediatrics.* 2004;114:555–76. [PubMed: 15286277]
55. Chobanian AV, Bakris GL, Black HR, Cushman WC, Green LA, Izzo JL, et al. Joint National Committee on Prevention DE, Treatment of High Blood Pressure. National Heart L, Blood I and National High Blood Pressure Education Program Coordinating C. Seventh report of the Joint National Committee on Prevention, Detection, Evaluation, and Treatment of High Blood Pressure. *Hypertension.* 2003;42:1206–52. [PubMed: 14656957]
56. Pickering TG, Hall JE, Appel LJ, Falkner BE, Graves J, Hill MN, et al. Recommendations for blood pressure measurement in humans and experimental animals: part 1: blood pressure measurement in humans: a statement for professionals from the Subcommittee of Professional and Public Education of the American Heart Association Council on High Blood Pressure Research. *Circulation.* 2005;111:697–716. [PubMed: 15699287]
57. Colan SD. Normal echocardiographic values for cardiovascular structures. In: Lai W. W, Cohen M. S, Geva T, Mertens L, editors. *Echocardiography in Pediatric and Congenital Heart Disease.* West Sussex, UK: Wiley-Blackwell; 2009. pp. 765–785. Appendix 1.
58. Mitchell GF, Wang N, Palmisano JN, Larson MG, Hamburg NM, Vita JA, Levy D, Benjamin EJ, Vasan RS. Hemodynamic correlates of blood pressure across the adult age spectrum: noninvasive evaluation in the Framingham Heart Study. *Circulation.* 2010;122:1379–86. [PMCID: PMC2981604] [PubMed: 20855656]
59. Pickering TG, Hall JE, Appel LJ, Falkner BE, Graves J, Hill MN, et al. Subcommittee of P and Public Education of the American Heart Association Council on High Blood Pressure R. Recommendations for blood pressure measurement in humans and experimental animals: Part 1: blood pressure measurement in humans: a statement for professionals from the Subcommittee of Professional and Public Education of the American Heart Association Council on High Blood Pressure Research. *Hypertension.* 2005;45:142–61. [PubMed: 15611362]

**Figure 1**



Survival after treatment of coarctation of the aorta. Survival curves of 819 surgical patients for over 60 years (reprinted with permission from Elsevier, license number 4131890880395)

**Figure 2**



Long-term Outcomes and Vascular Evaluation after Successful Coarctation of the Aorta Treatment study workflow. ABPM: Ambulatory blood pressure monitoring, Aix: Augmentation index, BP: Blood pressure, CMR: Cardiac magnetic resonance imaging, PWA: Pulse wave analysis, PWV: Pulse wave velocity, RHI: Reactive hyperemia index

**Table 1**

## Inclusion and exclusion criteria

Criteria	Definitions and comments
Inclusion criteria	
Coarctation of the aorta	
Current age 8-35 years	Lower age to allow facilitate the completion of the study tests and higher age to avoid overlap with aging-related vascular dysfunction <sup>184</sup>
Treatment for CoA after 1994	Date after which all three modalities were in clinical use.
Exclusion criteria	
Residual CoA	Systolic upper-to-lower extremity BP gradient >20 mmHg. * Residual gradient is a confounder since it impacts vascular function. †
Atypical CoA	Mid-thoracic or abdominal coarctation.
Severe transverse aortic arch hypoplasia	Transverse arch diameter z-score at initial echocardiogram <-4 <sup>†</sup>
Treatment of CoA at age <1y	A more severe disease subset, essentially amenable to surgery
Clinically significant associated cardiac defects that may affect independently vascular function	Mitral stenosis (echocardiographic mean inflow Doppler gradient >6 mmHg) aortic stenosis (echocardiographic mean Doppler gradient >20 mmHg); ventricular septal defect (>3 mm in diameter); atrial septal defect (required surgical or percutaneous closure other than a patent foramen ovale); other cardiac lesions that required medical, surgical or interventional treatment
Use of two treatment modalities for CoA	This does not include balloon dilation and subsequent stent placement at the same catheterization procedure
History of known vasculopathy with vascular dysfunction	Examples: Kawasaki disease, Takayasu's arteritis, Raynaud's disease
Genetic syndromes with diffuse arteriopathy	Examples: Williams syndrome, juvenile rheumatoid arthritis
Known traditional cardiovascular risk factors	Severe obesity (body mass index >95% for age and sex in children and >40 Kg/m <sup>2</sup> for adults); diabetes (fasting plasma glucose ≥ 126 mg/dl or random (non-fasting) glucose ≥ 200 mg/dl); hyperlipidemia (triglycerides ≥ 250 mg/dl; fasting LDL ≥ 190 mg/dl; HDL ≤ 30 mg/dl, currently taking statins or first degree relatives with familial hypercholesterolemia); smoking

Legend: y=years; BP=blood pressure; CoA=coarctation of the aorta; LDL=low-density lipoprotein cholesterol; HDL=high-density lipoprotein cholesterol; \*using highest lower extremity systolic blood pressure; †using previously published normative values<sup>187</sup>

**Table 2**

List of main clinical variables

<b>Variables</b>	<b>Comments or definitions</b>
Medical history	
Minimum transverse arch diameter	Using published normative values*
Z-score on initial echo	
Isthmus z score on initial echo	Using published normative values*
Initial Doppler coarctation gradient	mmHg
Bicuspid/Bicommissural Aortic Valve?	Yes/No
Initial arm-leg systolic BP gradient	mmHg
Visit BP	
Residual systolic BP gradient	Supine and automated mmHg
Right arm BP	Seated and manual mmHg

Legend: BP=blood pressure; \*using previously published normative values<sup>[57]</sup>

**Table 3**

List of main study test variables

<b>Variables</b>	<b>Comments or definitions</b>
<b>Applanation tonometry</b>	
Central systolic blood pressure	mmHg
Central pulse pressure	mmHg
Carotid-femoral PWV	meters/second
Augmentation index at HR75	%
<b>CMR</b>	
Left ventricular mass indexed to BSA	g/m <sup>2</sup>
Ascending Ao - Descending Ao	Meters/second
PWV (Ascending Ao to proximal, mid and distal descending Ao)	
Type of arch	Romanesque; Gothic; Crenel
<b>Aortic strain</b>	
(Ascending, Proximal, Mid and Distal Ao)	
Aortic Distensibility	mmHg-1
(Ascending, Proximal, Mid and Distal Ao)	
<b>Endo-PAT</b>	
Reactive hyperemia index (RHI)	
Augmentation index at 75 bpm	%
<b>ABPM</b>	
24 h Average systolic and diastolic BP	mmHg
24 h systolic and diastolic load	%
<b>Exercise test</b>	
Pre-Exercise SBP gradient	mmHg
Peak exercise BP	mmHg
<b>Biomarkers</b>	
NOx	ug/ml
ADMA	ng/L
High Sensitivity CRP	mg/L
VCAM-1	ng/ml
IL-1 $\beta$	pg/ml
TFG- $\beta$	
MMP-2/Gelatinase A	ng/ml
MMP-9/Gelatinase B	ng/ml

Legend: ADMA=Asymmetric Dimethylarginine; Ao=Aorta; BP=blood pressure; BP=Blood Pressure; BSA=Body Surface Area; CMR=Cardiac magnetic resonance; DBP=Diastolic Blood Pressure; HDL=High-density lipoprotein cholesterol; Hs-CRP=High sensitivity C-Reactive Protein; IL-1 $\beta$  = Interleukin 1 beta; LDL=Low-density lipoprotein cholesterol; MMP-2=Matrix Metalloproteinase-2; MMP-9=Matrix Metalloproteinase-9; NOx=Nitric Oxide; PWV=Pulse Wave Velocity; LV=Left Ventricle; SBP=Systolic Blood Pressure; TFG- $\beta$  = Transforming Growth Factor beta; VCAM-1=Vascular Cell Adhesion Molecule 1; \* using previously published normative values<sup>[57]</sup>

[Open in a separate window](#)

**Table 4**

Classification of BP Phenotype by ABPM

<b>Classification</b>	<b>Office BP SBP or DBP*</b>	<b>24h Mean ABPM SBP or DBP†</b>
Non-hypertensive	Pediatric: <95 <sup>th</sup> %tile Adults: <140/90 mmHg	Pediatric: <95 <sup>th</sup> %tile Adults: <135/85 mmHg
White Coat Hypertension	Pediatric: ≥95 <sup>th</sup> %tile Adults: >140/90 mmHg	Pediatric: <95 <sup>th</sup> %tile Adults: <135/85 mmHg
Masked Hypertension	Pediatric: <95 <sup>th</sup> %tile Adults: <140/90 mmHg	Pediatric: >95 <sup>th</sup> %tile Adults: >135/85 mmHg
Ambulatory Hypertension	Pediatric: >95 <sup>th</sup> %tile Adults: >140/90 mmHg	Pediatric: >95 <sup>th</sup> %tile Adults: >135/85 mmHg

Legend: AMBP=Ambulatory blood pressure monitoring; BP=Blood Pressure; ABPM=Ambulatory Blood Pressure Monitoring; Pediatric patients have age <18yo and adult patients age ≥18yo; %tile=percentile; BP=blood pressure; DBP=diastolic blood pressure; and SBP=systolic blood pressure. \*For pediatric patients, based on the National High Blood Pressure Education Program Task Force normative data<sup>[54]</sup>; for adult patients, based on the Joint National Committee on the Prevention, Detection, Evaluation, and Treatment of High Blood Pressure report.<sup>[55]</sup> †For pediatric patients, based on normative pediatric ABPM values from the American Heart Association Atherosclerosis, Hypertension and Obesity in Youth Committee of the Council on Cardiovascular Disease in the Young;<sup>[43]</sup> for adult patients, based on the Subcommittee of Professional and Public Education of the American Heart Association Council on High Blood Pressure Research report<sup>[59]</sup>

**Table 5**

Sample size estimation

<b>Smallest mean PWV (m/s) among groups</b>	<b>Largest mean PWV (m/s) among groups</b>	<b>Standard deviation</b>	<b>Sample size for each group</b>	<b>Total sample size</b>
4.0	4.8	1.0	30	90
4.0	4.8	1.1	36	108
4.0	4.8	1.2	43	129
4.4	5.3	1.0	24	72
4.4	5.3	1.1	29	87
4.4	5.3	1.2	34	102

Legend: PWV=Pulse wave velocity

Articles from Annals of Pediatric Cardiology are provided here courtesy of **Wolters Kluwer -- Medknow Publications**