



Research article

Histological and biochemical evidence of Cr₂O₃ and Al₂O₃ nanoparticles toxicity in the marine gastropod *Stramonita haemastoma*: A preliminary application of integrated biomarker response (IBR)

Fateh Sedrati^a, Hana Bouzahouane^{b,c,*}, Mohcèn Menaï^b, Fadila Khaldi^{a,b}, Tayeb Bouarroudj^d, Lassaad Gzara^e, Mounira Bensalem^f, Omar Laouar^{g,h}, Noomene Sleimiⁱ, Hichem Nasri^j, Carla O. Silva^k, Kheireddine Ouali^c

^a Laboratory of Sciences and Technology of Water and Environment, Mohamed Cherif Messaadia University BP1553, 41000 Souk Ahras, Algeria

^b Department of Biology, Faculty of Nature and Life Sciences, Mohamed Cherif Messaadia University Souk Ahras 41000, Algeria

^c Laboratory of Environmental Biosurveillance, Department of Biology, Faculty of Sciences, Badji Mokhtar University BP12, El Hadjar, Annaba 23000, Algeria

^d Center for Scientific and Technical Research in Physico-Chemical Analyses (CRAPC), BP384, Bou-Ismaïl, Tipaza, RP42004, Algeria

^e Center of Excellence in Desalination Technology, King Abdulaziz University P.O. Box: 80200, Jeddah 21589, Saudi Arabia

^f University August 20, 1955, Skikda, BP26, El Hadjak, Skikda, Algeria

^g Central laboratory of pathology and molecular biology, CHU, Annaba, Algeria

^h Faculty of Medicine, Badji Mokhtar University BP12, El Hadjar, Annaba 23000, Algeria

ⁱ RME-Laboratory of Resources, Materials, and Ecosystems, Faculty of Sciences of Bizerte, University of Carthage, Bizerte 7021, Tunisia

^j Laboratory of Biodiversity and Ecosystems Pollution, Faculty of Life and Nature Sciences, University of Chadli Bendjedid, El Taref, Algeria

^k MARE - Marine and Environmental Sciences Centre, ARNET - Aquatic Research Network Associate Laboratory, NOVA School of Science and Technology, NOVA University Lisbon, Caparica, Portugal

ARTICLE INFO

Edited by Martin Grosell

Keywords:

Stramonita haemastoma

Al₂O₃ NPs

Cr₂O₃ NPs

Integrated biomarkers response (IBR)

Oxidative stress

Histopathology

ABSTRACT

Nanoparticles (NPs) have actively contributed to nanotechnologies advancement over the last years, due to the unique properties they possess compared to their pristine counterparts. Consequently, NPs found wide applications in various fields such as the medical, biomedical, chemical, agro-food industries, and cosmetology. NP's extensive uses could lead to their release into the environment, especially in the marine ecosystems, considered as NPs sink, resulting in harmful effects on organisms. Concerns regarding NPs' toxicity in aquatic organisms have emerged, however, several points remain unexplored. In the present study, the toxicity of chromium oxide (Cr₂O₃ = 42 nm) and aluminum oxide (Al₂O₃ = 38 nm) NPs (1 mg/L, 2.5 mg/L, and 5 mg/L) in the gills of the marine gastropod *Stramonita haemastoma* was assessed through time (7, 14, and 28 days) by a multi-biomarker, Integrated biomarkers response (IBR), and Histological analysis. Both NPs induced varied changes in the antioxidant system, suggesting the onset of oxidative stress marked by superoxide dismutase (SOD), catalase (CAT), acetylcholinesterase (AChE), malonaldehyde (MDA), and glutathione (GSH) levels imbalance. Varied histological alterations in the gills of *S. haemastoma* were also observed including inflammation, hypertrophy, and lamellar fusion, IBR proved to be a promising tool for assessing NPs toxicity in gastropods. In this study results indicated the co-response of reduced glutathione (GSH), glutathione S-transferase (GST), glutathione peroxidase (GPx), CAT, SOD, and MT after 28 days of exposure. *S. haemastoma* showed sensitivity to all exposure concentrations of NPs thus validating this species as a suitable indicator of NPs contamination and toxicity.

* Corresponding author at: Department of Biology, Faculty of Nature and Life Sciences, Mohamed Cherif Messaadia University, Souk Ahras 41000, Algeria.

E-mail address: h.bouzahouane@univ-soukahras.dz (H. Bouzahouane).

<https://doi.org/10.1016/j.cbpc.2025.110159>

Received 14 October 2024; Received in revised form 29 January 2025; Accepted 20 February 2025

1532-0456/© 20XX

1. Introduction

The advances in nanotechnologies over the last decade have contributed to extending the interest given to Nanoparticles (NPs) in a wide range of application areas. NPs are typically defined as particles with at least one dimension varying from 1 to 100 nm, allowing them to exhibit specific functional characteristics superior to their bulk elements (Abdel-Latif et al., 2020; Abramenko et al., 2021). Consequently, NPs have been utilized in various sectors including medical, biomedical, chemical industries, environmental applications, and cosmetics (Sánchez et al., 2011; Corsi et al., 2020). According to economic data, the global market size of metallic nanoparticles (Me-NPs) is predicted to reach 4.2 billion USD with an annual growth rate of nearly 11.5 % (focus on catalysis, 2023).

The increasing use of NPs in various applications raises concerns about their release into the environment, especially in the aquatic ecosystems, which are considered their major sink (Baker et al., 2014; Bour et al., 2015). For instance, processes such as atmospheric deposition, environmental dumping, effluents, and others could lead to the introduction of NPs in the marine ecosystem as their final destination (Matranga and Corsi, 2012; Xia et al., 2017). Once in seawater, NPs can form agglomerations and settle in marine sediments, leading to their accumulation over time (Canesi et al., 2015; Peng et al., 2017). Studies have shown that NPs can reach concentrations in marine sediments in the range of mg/kg (Nam et al., 2014; Fan et al., 2018; Tou et al., 2021). These NPs are considered pollutants that can potentially harm effects to marine species and compromise the integrity of the aquatic ecosystem (Canesi et al., 2017). Consequently, concerns have emerged regarding NPs' potential toxicity (Blasco et al., 2015). Despite the recent expansion of ecotoxicological research, previous investigations primarily focused on freshwater species and specific NPs types such as TiO_2 ,

CuO , and ZnO_2 . Likewise, these studies were mainly limited to bivalves regarding marine species (Minetto et al., 2014 and 2016).

The characteristics that make NPs profitable, including their small size, high reaction surface, shape, and optical ones are crucial factors in their potential toxicity to aquatic organisms (khan et al., 2019; Turan et al., 2019). Several studies reported their ability to interact with biological barriers, affecting their bioavailability and causing harmful effects, particularly by inducing oxidative stress, which is considered the main toxicity mechanism of NPs (De Felice and Parolini, 2020; Hichem et al., 2022). Exposure to NPs can alter cellular antioxidant defenses, leading to various effects including the modification of cellular components such as proteins, lipids, and DNA, mainly through indirect oxidation pathways due to ions released in aquatic organisms (Handy et al., 2008; Deng et al., 2016; Hou et al., 2019).

Chromium oxide (Cr_2O_3) and Aluminum oxide (Al_2O_3) are among the most widely used NPs. Due to their high functional properties, both NPs found application in numerous sectors, namely as green pigments, catalysts, semiconductors, chemical and agro-industrial industries, biomedical, medical, cosmetics, antibacterial agents, and others (Becker et al., 2016; Mui et al., 2016; Sharma et al., 2022; Adnan and Mohammed, 2024). Several studies have reported their toxicity to various biological models. Cr_2O_3 was reported to induce cytotoxicity in the green alga *Chlamydomonas reinhardtii* and L929 cells (Alarifi et al., 2016; da Costa et al., 2015), genotoxicity in rats (Singh et al., 2016), high acute toxicity in *Daphnia magna* (Puerari et al., 2016), and oxidative stress induction in the freshwater fish *Labeo rohita* (Kanwal et al., 2019). Although Al_2O_3 NPs have been considered less toxic than most common NPs, numerous studies reported also toxic effects induced by Al_2O_3 (García-Saucedo et al., 2011). For instance, cytotoxicity in mouse neuroblastoma cells (Nogueira et al., 2019), oxidative stress induction in the mussel *Unio*




Fig. 1. XRD of Cr_2O_3 (A) and Al_2O_3 (B) nanoparticles, SEM of spherical shape Cr_2O_3 (C) and $\alpha\text{-Al}_2\text{O}_3$ (D) NPs.




Fig. 2. zeta potential And The Zeta-average diameter (nm) of Cr_2O_3 and Al_2O_3 Nps as a function of pH.

tigridis (Canli and Canli, 2021), histological alterations in the fish *Oreochromis mossambicus* (Murali et al., 2018), neurotoxicity on dopaminergic neurons in rats (Liu et al., 2020), and accumulation in *Artemia salina* larvae (Ates et al., 2015). Despite the extensive research on NPs toxicity assessment, the existing literature concerning both NPs remains limited, especially on aquatic organisms (Załęska-Radziwiłł et al., 2020). For instance, to the best of our knowledge, no prior study has explored the toxicity of Cr_2O_3 NPs on marine organisms, with limited existing studies on freshwater species.

The increasing population density and economic activities are putting pressure on coastal ecosystems. Therefore, it is becoming increasingly important to evaluate NPs fate and toxicity on benthic organisms (Mouneyrac et al., 2014). Mollusks have been widely used as bioindicators for assessing NPs aquatic contamination and toxicity by several relevant studies. For instance, marine bivalves have been recognized as particularly affected by NPs contamination, leading to a focus on bivalves in many studies, while research focused only on other species is still lacking (Canesi et al., 2010a; Canesi et al., 2012; Rocha et al., 2015).

Stramonita haemastoma is a coastal marine gastropod, with a wide geographic distribution. These marine snails are principal predators of bivalves and other small gastropods. They have been used as indicators of Tributyltin (TBT) pollution and imposex induction; besides owning economic importance in purple dye extraction and human consumption (Bouzahouane et al., 2018;

Madeira et al., 2018; El Ayari et al., 2018). Although *S. haemastoma* has received little attention as a bioindicator, recent limited *in situ* studies have judged this marine snail as a promising indicator of marine metallic contamination (Di Bella et al., 2018; Bouzahouane et al., 2024). However, to the extent of our knowledge, this marine gastropod has not been used as an indicator of NPs contamination and toxicity. Considering the earlier key points, *S. haemastoma* could be a new promising bioindicator of NPs toxicity assessment, since gastropods are usually suitable biological models as they offer several practical advantages namely their abundance, breeding resistance, and sensitivity to pollutants exposure (Caixeta et al., 2020). To our best knowledge, No prior study apart from ours regarding the use of *S. haemastoma* as a bioindicator of NPs toxicity same as *in vivo in situ* studies was done. Key biochemical and histological biomarkers linked to critical physiological functions were selected for this study. Biochemical indicators include the activities of antioxidant enzymes such as superoxide dismutase (SOD), catalase (CAT), reduced glutathione (GSH), and glutathione peroxidase (GPx), all essential for defense against reactive oxygen species (ROS) generated by xenobiotic exposure. Oxidative damage was assessed by measuring malondialdehyde (MDA) levels. Additionally, Metallothionein (MT) levels were evaluated as a biomarker for metal detoxification. The activity of glutathione-S-transferase (GST), a phase II biotransformation enzyme involved in xenobiotic detoxification, and acetylcholinesterase (AChE) activity, associated with cholinergic neurotransmission,




Fig. 3. GSH, and GST vr ates in the Gills of *Stramo nita haem astoma* treated with Cr_2O_3 (A, A') and Al_2O_3 (B,B'). The values are expressed as mean $\pm$ S.d. $n = 3$ (* $p < 0.05$).

was also evaluated as biomarkers for neuromuscular toxicity. Additionally, a detailed histological study was conducted to observe structural alterations in the gill tissues, providing insights into tissue-level damage caused by NPs exposure.

Integrated biomarker response (IBR) is a useful tool that allows the visualization of multivariate data of numerous biomarker response trends (Beliaeff and Burgeot, 2002). This method has been widely used, especially for *in situ* investigations, including ecotoxicological studies to better understand the correlations among biomarkers and contamination levels in field studies (Devin et al., 2014). However, the application of IBR in *in vivo* and *in vitro* NPs toxicological studies is still limited. For example, in marine species, the use of IBR is mainly limited to bivalves in some studies (Xia et al., 2013; Devin et al., 2017; Xia et al., 2017; Li et al., 2021).

This study aims to investigate the potential *in vivo* toxicity of Cr_2O_3 and Al_2O_3 NPs, with increasing exposure concentrations through time in the gills of *S. haemastoma*. This will be accomplished using a multi-biomarkers approach and the first application of IBR to a range of biomarkers over time, along with histological analysis.

2. Materials and methods

2.1. Nanoparticle synthesis

The solvothermal method was used to prepare both Al_2O_3 and Cr_2O_3 NPs, by dissolving 0.4 mol of $\text{Al}(\text{NO}_3)_3 \cdot 9\text{H}_2\text{O}$ in 40 mL of deionized water (Dw) and $\text{Cr}(\text{NO}_3)_3 \cdot 9\text{H}_2\text{O}$ in 60 mL of Dw under stirring, respectively. For Al_2O_3 preparation, the main mixture was centrifuged then the precipitate was dispersed in distilled water,

heated at 400°C and washed and finally dried for 8 h at 90°C . The Cr_2O_3 NPs mixture was heated at 140°C for 24 h, the resulting precipitates were washed with Dw, dried, and calcined at 500°C to obtain green Cr_2O_3 NPs.

2.2. Nanoparticle characterization

The structural characterization of both samples was performed using X-ray diffraction (XRD) with a Siemens D-5000 diffractometer in $20^\circ - 90^\circ$ 2θ range with $\text{Cu-K}\alpha$ radiation ($\lambda = 1.5418 \text{ \AA}$). UV-Vis spectrophotometer was used to investigate UV-Vis absorption spectra for both NPs and Morphological analysis using Scanning Electron Microscopy (SEM: Thermo Fisher Scientific) was also performed (Bouarroudj et al., 2023).

Malvern Zetasizer Nano-ZS (Malvern Panalytical) equipped with DTS 1070 disposable folded capillary cells and MPT-2 Titrator, was used to assess the zeta potential and the zeta average diameter of both NPs in function of pH using NaOH (0.03 M) and HCl (0.03 M) solutions, by dynamic light scattering (DLS) and calculation of 20 measurements points. Energy-dispersive spectroscopy (EDS) was also used to investigate the purity of both further prepared Al_2O_3 and Cr_2O_3 NPs and the optical properties of both NPs were also evaluated by calculating the optical band gap (Eg) from UV-Visible spectra for Cr_2O_3 and Al_2O_3 NPs (supplementary).

2.3. Animal sampling and conservation

Stramonita haemastoma sampling (4 to 10 cm) followed previous work by Sedrati et al. (2024). Briefly, after collection from




Fig. 4. GPx, and CAT rates in the Gills of *Stramonita haemastoma* treated with Cr_2O_3 (A, A') and Al_2O_3 (B,B'). The values are expressed as mean $\pm$ S.d. n = 3 (* p < 0.05).

Annaba city, Algeria in the Cap de Garde (36° 96' N, 7° 78' E) by SCUBA. This area is an isolated natural site, characterized by the absence of industrial activity or human habitation (Bouzahouane et al., 2024). Natural seawater (NSW) was collected from the same sampling site, filtered at 50 μm , and stored in the dark. The preliminary average water parameters were monitored by a multiparameter (®Horiba) and were maintained during all experimental periods (salinity: 34.2 ppt, Temperature: 23 °C, pH: 8.2, dissolved oxygen: 9.22 mg/L). The collected snails were acclimated for 7 days and kept in laboratory conditions with a photoperiod of 12:12 h dark/light in NSW-contained aquariums with constant aeration, water was renewed daily during the acclimation period, and then every 48 h in exposure periods.

2.4. Experimental design

S. haemastoma snails were randomly divided into groups of 20 individuals each, and each group was placed in identically sized boxes containing 3 L of NSW. The snails were maintained in constant aeration in bubbling conditions, mimicking their natural conditions. Marine snails were exposed for a total of 28 days, with three assessment endpoints at 7, 14, and 28 days to either Cr_2O_3 or Al_2O_3 NPs. Nano-powders of Cr_2O_3 and Al_2O_3 were added to MilliQ water with stirring to prepare stock solutions (1 g/L) of both NPs using a bath sonicator (50 kHz) for 30 min. Exposure was made by diluting equal volumes of stock solutions into each box. The snail groups were set and exposed to nominal concentrations as follows: group 1 as control, group 2 exposed to 1 mg/L Al_2O_3 NPs, group 3 exposed to 2.5 mg/L Al_2O_3 NPs, group 4 exposed to 5 mg/L Al_2O_3 NPs, group

5 exposed to 1 mg/L Cr_2O_3 NPs, group 6 exposed to 2.5 mg/L Cr_2O_3 NPs, and group 7 exposed to 5 mg/L Cr_2O_3 NPs. Gastropods were not fed during the exposure term, and experiments were conducted with 3 independent replicates. NPs nominal concentrations in this study were selected following the previously applied concentrations in prior studies, and the predicted concentrations in sediments (Canesi et al., 2010a, 2010b; Hao and Chen, 2012; Gornati et al., 2016; Xia et al., 2017; Caixeta et al., 2020). Exposure duration was selected according to some criteria as follows: - The average exposure periods reported in related previous studies and reviews, - periods were selected to assess the potential effects evolution in a sub-acute term (Canesi et al., 2012; Abdel-Latif et al., 2020; Roma et al., 2020).

2.5. Biochemical analysis

2.5.1. Preparation of tissue homogenates

At the end of each exposure duration point (7, 14, and 28 days), snails from the control and the exposed groups were euthanized by instant freezing and carefully dissected. The gills were obtained for oxidative stress biomarkers analysis, and 03 individuals were used for analysis at each period from different replicates. Immediately, aliquot samples were homogenized using an ultrasonicator bath and a vortex, using 3 buffers, each specific to biomarkers. Briefly, aliquots samples were homogenized in Tris-HCl buffer (10 mM, pH: 7.3, 1:3 w/v) for reduced glutathione (GSH), glutathione peroxidase (GPx), glutathione-S-transferase (GST), superoxide dismutase (SOD), catalase (CAT), and malondialdehyde (MDA) assays (Zaidi et al., 2021). Aliquot samples were homogenized in phosphate buffer




Fig. 5. SOD, and MDA rates in the Gills of *Stramonita haemastoma* treated with Cr_2O_3 (A, A') and Al_2O_3 (B, B'). The values are expressed as mean $\pm$ S.d. n = 3 (* p < 0.05).

(100 mM, pH, 7, 1:3 v/w) for acetylcholinesterase (AChE) analysis. Homogenates were then centrifuged at $11250 \times g$ for 15 min (4°C) and supernatants were immediately stored at -80°C .

Tris-HCl (20 mM), saccharose (0.5 M), and leupeptin (0.006 mM) were used as a buffer for the metallothionein (MT) assay, homogenates were centrifuged at $20,000 \times g$ for 30 min (4°C) and instantly conserved at -80°C (Bouzahouane et al., 2018).

2.5.2. Protein quantification

The total protein rates were determined according to the Bradford (1976) assay at 595 nm using Bovine serum Albumin (BSA) as a standard (Sigma®).

2.5.3. Oxidative stress biomarkers

The Weckbecker and Cory (1988) method was used to monitor GSH levels, based on the 5,5'-dithio-bis-2-nitrobenzoic acid (DTNB) reduction by thiol group (-SH) of glutathione absorbance assessment at 412 nm. The GSH content was expressed as nanomole per milligram of protein (nmol mg^{-1} protein).

The GST activity was assessed using 1-chloro-2,4-dinitrobenzene (CDNB) as a substrate for reduced glutathione resulting in the formation of molecules from the conjugation reaction then monitored the absorbance changes for 2 min at 340 nm according to Habig et al. (1974) method, GST activity was expressed in units (U) of GST per milligram of protein (U GST mg^{-1} protein).

The GPx activity was measured following the Chiu et al. (1976) method, by absorbance assessment at 412 nm and GPx activity was defined as the amount of enzyme oxidizing 1 μmol of GSH. GPx was

expressed as nanomoles of GSH oxidized mg^{-1} protein (Thirupurasundari et al., 2009; Wu et al., 2011).

The Marklund and Marklund (1974) method based on the pyrogallol oxidation inhibition by SOD in the presence of EDTA assay was employed to assess SOD enzymatic activity by absorbance monitoring for 5 min at 420 nm, SOD concentrations were represented in (U) SOD mg^{-1} protein.

CAT enzymatic activity was measured using the Hydrogen peroxide (H_2O_2) rates decrease and monitored optical densities variation at 240 nm according to Beer and Size (1952) assay, one unit (U) of CAT is equivalent to the amount of enzyme catalyzing the degradation of 1 μmol of H_2O_2 per min and was expressed as $\text{U min}^{-1} \text{mg}^{-1}$ protein.

2.5.4. Lipid peroxidation biomarker (LPO)

The evaluation of MDA rates in the gills of the marine snails was measured as an indicator of Lipid peroxidation (LPO), MDA levels were assessed by the thiobarbituric acid (TBA) reaction assay at 535 nm in a final mixture composed of 1 ml of the supernatant, 1 ml TBA (0.67 %), and 1 ml trichloroacetic acid (TCA, 5 %) according to the initial protocol reported by Fatima et al. (2000) with some modifications by Zhao et al. (2009). An extinction coefficient of $1.56 \times 10^5 \text{ M}^{-1} \text{cm}^{-1}$ was used to calculate the MDA rates which was expressed as nmol MDA mg^{-1} protein. More detailed procedures of the biomarkers analysis were previously reported by Zaidi et al. (2021).




Fig. 6. MT, and AChE rates in the Gills of *Stramonita haemastoma* treated with Cr₂O₃ (A, A') and Al₂O₃ (B, B'). The values are expressed as mean $\pm$ S.d. n = 3 (* p < 0.05).

2.5.5. Neurotoxicity biomarker

The AChE enzymatic activity was determined following the Ellman et al. (1961) assay, based on acetylcholine (ACh) as a substrate for (AChE) and the formation of yellow color molecules due to thiocholine (Sch) reaction with DTNB, which can be monitored at 412 for 5 min for the estimation of enzymatic activity, AChE activity was expressed in nmol min⁻¹ mg⁻¹ protein.

2.5.6. Metallothionein (MT) analysis

Metallothionein (MT) content was evaluated according to the method proposed by Viarengo et al. (1997) specifically adapted for marine invertebrates and based on the monitoring of absorbance at 412 nm with Ellman's reagent (DTNB) for the determination of the thiols (-SH) after obtaining a partially purified MT fraction from the cytosolic content using an ethanol/chloroform treatment. The MT content was measured by 1 mol Mt. = 20 mol GSH as an equation and was expressed as nmol g⁻¹ tissue.

2.6. Histological study

The gills samples from each group (control, exposed to Cr₂O₃ and Al₂O₃ NPs) were dissected immediately for histological observations after 28 days of exposure, then preserved in 10 % formaldehyde. 2 individuals were pulled from different replicates for all groups and then histological samples were dehydrated with a succession of graduated ethanol and cleared using xylene, 3 to 5 μ m sections were obtained by a microtome after embedding gills samples in prepared paraffin wax (@Sigma). Tissue cuts were then directly mounted in microscopic slides and colored using Eosin (e) and

Hematoxylin (H). Finally, the prepared histological slides of gills samples were observed for alteration determination in a Leica DM1000 LED optical microscope, and photographic shots of the histological slides were recorded (Hould, 1984; Zaidi et al., 2021).

2.7. IBRv2 analysis

IBR is a method used to integrate multiple biomarker responses and experimental factors into one single index. This helps to better frame the key impacts and reactions of biomarkers. The method has been successfully used to evaluate the effects of NPs on oxidative stress biomarkers in bivalves (Beliaeff and Burgeot, 2002; Marigómez et al., 2013). The term "v2" in IBRv2 typically stands for "version 2," which indicates an updated version of the original Integrated Biomarker Response (IBR) index and is referred to as IBR in this study. This version incorporates adjustments to improve its applicability, accuracy, or sensitivity in environmental biomonitoring. In this study, IBRv2 was applied to various biomarkers and exposure concentrations for each exposure period to identify the potential impacts of NPs. The IBRv2 calculations are based according to Sanchez et al. (2013) modifications to adapt IBR in oxidative stress biomarkers as follows:

Individual biomarkers data (X_i) is compared to average biomarkers references (X_0) to obtain (Y_i), and log transformation is applied to reduce variations.

$$Y_i = \log (X_i/X_0)$$

Standardized biomarkers response (Z_i) is obtained by estimating the global mean (μ) and standard deviation (σ) of Y_i .




Fig. 7. IBR v2 Star plots of GST, GPx, CAT, SOD, and AChE activities (enzymatic biomarkers) and GSH, MT, MDA Contents (Non enzymatic biomarkers) in gills of *Stramonita haemastoma* treated with Cr_2O_3 . For 7, 14, and 28 exposure days.

$$Z_i = (Y_i - \mu) / \sigma$$

A deviation index (A_i) is calculated to establish a basal line of 0 as a center in which biomarkers' response changes are represented, by applying the average reference biomarker data (Z_0) and (Z_i).

$$A_i = (Z_i - Z_0)$$

Finally, the values of A_i parameters for each biomarker are calculated to obtain the IBR values.

$$\text{IBR} = \sum |A_i|$$

2.8. Statistical analyses

All statistical data analyses were conducted using R software (R Team, 2023; Core Team, 2023), and a two-way ANOVA permutation was performed to compare groups, NPs concentrations, and duration. Exposure was compared by applying the *Post hoc* test of Dunnett for Multiple Comparisons of Means; followed by the Dunn multiple comparison test with *p*-values adjusted with the Bonferroni method. Descriptive statistics are expressed as mean and standard deviation and statistical significance was determined at a significance level of $p < 0.05$ for all analyses. The percentage variation rates of biomarker activity in the gills of snails were calculated to quantify the relative changes between the control and exposed groups for each time duration. The formula used to calculate the percentage increase or decrease is as follows:

$$\text{Percentage Change (\%)} = ((X_{\text{exposed}} - X_{\text{control}}) \times 100) / X_{\text{control}}$$

Where:

X_{exposed} represents the biomarker activity measured in the gills of snails exposed to nanoparticles (NPs) at a given concentration and time.

X_{control} represents the biomarker activity measured in the gills of snails from the control group (not exposed to NPs).

The resulting percentage values indicate an increase (positive percentage) or decrease (negative percentage) in biomarker activity relative to the control.

3. Results

3.1. Cr_2O_3 and Al_2O_3 NPs characterization

3.1.1. XRD analysis

The crystal structures and purity of both NPs were assessed by X-ray diffraction (XRD) analysis. The diffraction peaks observed at 2θ values correspond to the lattice plane (012), (104), (110), (006), (113), (202), (024), (116), (122), (214), (300), (119), (220), (306), (312), (0210) and (134), indicating the rhombohedra phase for Cr_2O_3 (Fig. 1A) (space group, R-3c). For $\alpha\text{-Al}_2\text{O}_3$, the peaks correspond to the lattice plane (012), (104), (110), (006), (024), (116), (211), (018), (214), (300), (119), (220), (131), (128) and (0 2 10) (Fig. 1B) (space group, R-3c). Both phase observations agree with the literature values (Kafi-Ahmadi et al., 2022).

The average grain size (D) of Al_2O_3 NPs was determined to be 38 nm, while for Cr_2O_3 NPs, it was calculated to be 42 nm, (D) using the Scherrer formula (Messai et al., 2018).

3.1.2. SEM analysis

SEM analysis was performed to investigate the morphology and structure of both Al_2O_3 NPs and Cr_2O_3 NPs. Both NPs samples exhibited remarkably consistent sizes indicating a successful synthesis




Fig. 8. IBR v2 Star plots of GST, GPx, CAT, SOD, and AChE activities (enzymatic biomarkers) and GSH, MT, MDA Contents (Non enzymatic biomarkers) in gills of *Stramonita haemastoma* treated with Al_2O_3 NPs. For 7, 14, and 28 exposure days.

process. The SEM images revealed that Cr_2O_3 NPs exhibit a spherical shape (Fig. 1C), while Al_2O_3 NPs showed plate-like $\alpha\text{-Al}_2\text{O}_3$ (Fig. 1D).

3.1.3. EDS analysis

The presence of Cr_2O_3 and Al_2O_3 NPs in each sample was confirmed along with high purity by the characteristic peaks corresponding to Cr, O, and Al, O elements in the EDS analysis (supplementary).

3.1.4. DLS analysis

The zeta potential and the zeta average diameter of the NPs in this study showed high variations depending on pH. The isoelectric points (IEP) of Al_2O_3 and Cr_2O_3 were 2.33 and 2.95, respectively, indicating an acidic character. Suspensions of Al_2O_3 and Cr_2O_3 NPs are stable only at pH higher than 6.5, and optimal stability was noticed in a pH range of 7 to 7.6 (Fig. 2). Additionally, the optical properties of both NPs were also evaluated and presented in the supplementary materials.

3.2. Biochemical biomarkers

The percentage variation rates among the control and the groups exposed to both NPs in the gills of snails were calculated at each time duration. The percentage variation rates of biomarker activity in the gills of snails were calculated to quantify the relative changes between the control and exposed groups for each time duration, the multi-biomarkers approach results will be presented by exposure periods. The variations trends of all the studied biomarkers for the control groups and the exposed groups (Cr_2O_3 or Al_2O_3 NPs) are represented in Fig. 11, in the gills of *S. haemastoma* the levels and activities of GSH, CAT, AChE, MT, and MDA showed significant

difference ($p < 0.05$) among exposure periods (7, 14, and 28 days). GST and SOD activities showed significant differences ($p < 0.05$) among concentrations and control groups for both NPs.

3.2.1. Following 7 days

The changes in GSH, GST, GPx, and CAT activity in the gills of snails are represented in Fig. 3 and Fig. 4 for both NPs. After 7 days, exposure to 1 mg/L ($p < 0.05$) and 2.5 mg/L of Al_2O_3 NPs led to 55.9 % and 42.6 % increase in GSH, while 1 mg/L of Cr_2O_3 caused 34.08 % GSH depletion. GST showed an increase of 145 % and 290 % ($p < 0.05$) in groups exposed to 1 mg/L and 5 mg/L of Al_2O_3 and approximately 133.01 % in the group exposed to 5 mg/L Cr_2O_3 . GPx activity tended to increase in the gills of all exposed groups, ranging from 105 to 170 % for Al_2O_3 NPs and 117 to 385 % for groups treated with Cr_2O_3 . CAT activity was reduced by 70.9 % following exposure to 1 mg/L of Cr_2O_3 , while groups exposed to 2.5 mg/L and 5 mg/L of Al_2O_3 showed higher ($p < 0.05$) CAT activity, ranging from 37.4 % to 65.4 %. SOD, MDA, MT, and AChE variations in the gills of snails for both NPs are represented in Fig. 5 and Fig. 6. A significant decrease ($p < 0.05$) ranging from 50 % to 66 % in SOD activity was observed in all groups treated with Al_2O_3 , with similar reduction ($p < 0.05$) of 71 % in the group exposed to 2.5 mg/L of Cr_2O_3 . MDA in all exposed groups showed higher levels ranging from 156 % to 300 % for Al_2O_3 and significant depletion ($p < 0.05$) in MT rates was observed in the gills of all groups exposed to Cr_2O_3 ranging from 45 % to 49 %. Exposure to 2.5 and 5 mg/L of Al_2O_3 induced significantly lower MT levels ($p < 0.05$), ranging from 44 % to 109 %. AChE activity in the gills of *S. haemastoma* was significantly ($p < 0.05$) decreased by 263.28 % due to 2.5 mg/L exposure to Cr_2O_3 , while a 215 % significant increase and a 75 % decrease at 2.5 mg/L and 1 mg/L was observed, respectively, for Al_2O_3 .




Fig. 9. Histological alterations observed in the Gills of *Stramonita haemastoma* exposed to 28 days of Cr_2O_3 and Al_2O_3 . A,B: Control gills filaments, C: Control ospharidium, D (Cr_2O_3 exposure), E (Al_2O_3 exposure): Goblet cells proliferations, F (Cr_2O_3 exposure), H (both NPs), I (Al_2O_3 exposure): central cavity hypertrophy, lamella fusion G (Al_2O_3 exposure): filaments thickening (observed in both NPs exposure), I (Cr_2O_3 exposure): filaments alteration. coloration H&E X100, X200, X400 low power.

3.2.2. Following 14 days

After 14 days, the levels of GSH significantly increased ($p < 0.05$) by 65 % at 1 mg/L and 2.5 mg/L of Al_2O_3 , while GSH depletion was observed in all groups exposed to Cr_2O_3 with a significant ($p < 0.05$) decrease of 46 % observed at 2.5 mg/L. GST activity showed significantly decreased concentrations in all groups, ranging from 33 % to 74 % for Al_2O_3 and, 6 % to 45 % for Cr_2O_3 . GPx activity decreased by 24 % (1 mg/L) ($p < 0.05$) and 35 % (5 mg/L) due to Cr_2O_3 exposure. On the other hand, GPx activity increased by 15.5 % (2.5 mg/L) and 24 % (5 mg/L) in gills exposed to Al_2O_3 . CAT activity increased by 125 % at 1 mg/L and a significant decrease ($p < 0.05$) by 64 % at 5 mg/L was noted in groups exposed to Cr_2O_3 with decreases ranging from 46 % to 56 % at 2.5 mg/L and 5 mg/L of Al_2O_3 . After 14 days, SOD was highly affected by Al_2O_3 ($p < 0.05$) NPs, with a significant decrease ranging from 45 % to 55 % in all treated groups. In the same way, MDA levels were significantly ($p < 0.05$) higher (95 %) at 5 mg/L of Al_2O_3 . The exposure to 1 mg/L and 2.5 mg/L of Cr_2O_3 NPs caused significant ($p < 0.05$) depletion of MT levels (76 % and 83 %, respectively), while 1 mg/L of Al_2O_3 induced reduced (183 %) MT rates. There was a significant inhibition (60 % to 81 %) of AChE activity in all groups treated with Cr_2O_3 , and similarly, all groups exposed to Al_2O_3 presented decreased enzymatic activity (36 % to 63 %).

3.2.3. Following 28 days

At the end of exposure duration (28 days), both NPs increased the levels of GSH in all groups, with significant ($p < 0.05$) induction at 5 mg/L of Cr_2O_3 (239 %) and Al_2O_3 (268 %). There was a pronounced induction in GST activity in all treated groups, with the most significant increase ($p < 0.05$) at 2.5 mg/L of Cr_2O_3 (310.1 %) and 5 mg/L of Al_2O_3 (351 %). Similarly, all groups for both NPs showed increased GPx activity with the most significant ($p < 0.05$) variations at 2.5 mg/L of Cr_2O_3 (382 %) and 5 mg/L of Al_2O_3 (272 %). Likewise, CAT activity increased for all groups at day 28, with significant ($p < 0.05$) induction at 1 mg/L of Cr_2O_3 (65 %) and 5 mg/L of Al_2O_3 (132 %). In contrast, SOD activity was reduced in all exposed groups for both NPs with significant inhibition observed at 5 mg/L of Cr_2O_3 (73 %). MDA levels in snails' gills were highly affected by Al_2O_3 NPs, with increased rates ranging from 214 % to 41. Similarly, Cr_2O_3 exposure resulted in increased ($p < 0.05$) MDA levels at 2.5 mg/L (81 %) and 5 mg/L (227 %). There was a significant ($p < 0.05$) depletion of MT rates at 1 mg/L Cr_2O_3 (53 %) and a significant ($p < 0.05$) induction at 1 mg/L of Al_2O_3 . AChE activity was induced by both NPs, with a dose-dependent increase ranging from 5 % to 46 % due to Cr_2O_3 exposure. An increase ranging from 152 % to 203 % was also observed after Al_2O_3 exposure. Significant ($p < 0.05$) among exposure duration points was observed for GSH, CAT, MDA, MT, and AChE in the gills of snails treated with Cr_2O_3 , and GSH, GST, CAT, and MDA in individuals exposed to Al_2O_3 .




Fig. 10. IBR v2 values evolution through exposure periods (7, 14, and 28 days) and both Cr_2O_3 and Al_2O_3 NPs concentrations (control, 1, 2.5, and 5 mg/L) in *Stramonita haemastoma* gills.

3.3. IBR v2

Applying the multi-biomarkers approach is challenging for the comparative aspects among biomarkers or even pollutants effects. During this study, IBR v2 was used to analyze various biomarkers and present their responses in graphical summaries with standardized numeric values, allowing better identification of the impacts (Liu et al., 2017; Li et al., 2021). IBR positive values indicate biomarker induction, while negative ones are inhibition, following the recommendation of IBR v2 conception (Sanchez et al., 2013). IBR values have been plotted and organized into two categories: 1- enzymatic biomarkers, and 2- non-enzymatic biomarkers, providing a clearer understanding of the results and specific identification. IBR v2 results are represented in Fig. 7 for individuals exposed to Cr_2O_3 and Fig. 8 for individuals exposed to Al_2O_3 . IBR v2 global evolution according to exposure periods and concentrations for each and both NPs are represented in Fig. 10 and could serve as a summary tool to identify the potential biomarkers sensitivity trend and thus biomarkers behavior and response to both NPs exposure.

3.3.1. At 7 days

After 7 days of exposure to Cr_2O_3 NPs at 1 mg/L, the IBR values showed negative responses for SOD, CAT, MT, and positive responses for MDA. At 2.5 mg/L, CAT and SOD had a more negative impact along with MT. At 5 mg/L, IBR values highlighted negative AChE, GSH, and positive responses for GST and MDA. The IBR values highlighted positive values for MDA and GPx, and negative responses for SOD and MT responses in all groups exposed to Al_2O_3 .

3.3.2. At 14 days

On day 14, IBR levels at 1 mg/L (Cr_2O_3) were related to CAT, negative AChE, and MT response, and further highlighted negative GST, MDA, and GSH response. SOD and GPx were the most affected biomarkers at 5 mg/L. On the other hand, in individuals exposed to 1 mg/L of Al_2O_3 the IBR values reflected SOD inhibition and GSH induction. At 2.5 mg/L, GST and SOD were mainly inhibited and GSH was induced. The IBR index represented positive MT, MDA, and negative AChE response at 5 mg/L of Al_2O_3 . Overall, when setting

the mean values of the IBR index per NPs concentration, the trend was mostly dose-dependent at all periods.

3.3.3. At 28 days

After 28 days, the IBR showed the co-action of all biomarkers in a dose-dependent trend, showing high responses for all biomarkers for both NPs. $\text{CAT} < \text{GST} < \text{GPx} < \text{AChE} < \text{MDA}$ induction with SOD inhibition represented the main response for both NPs with the higher IBR values observed for Al_2O_3 .

3.4. Histological analysis

The respiratory system anatomy of *Muricidae* like *Stramonita haemastoma* is considered far-developed. However, there is limited information available in the literature. According to our observations and literature (Beesley et al., 1998), *S. haemastoma* gills are mainly composed of a left ctenidium as a primary respiratory organ suspended by membranes to the mantle cavity and a sensitive organ called ospharidium which primarily plays a water-taster role. Gills are composed mostly of leaflets (filaments) containing running blood vessels. Histological shots as shown in Fig. 9, revealed that the control gills exhibited a regular structure, normal filament organization, and a central cavity (Fig. 9. A, B). However, after 28 days of exposure to 1 mg/L of Cr_2O_3 . There was a pluristratification of the goblet cells in the gills filaments extremities. Exposure to 1 mg/L of Al_2O_3 and 2.5 mg/L of both NPs resulted in focal thickening of the gills epithelium and filaments, with a progressive loss in structure organization (Fig. 9. G, H, I). Lamella fusion was also observed (Fig. 9. F, I). On the other hand, exposure to 5 mg/L of Cr_2O_3 led to a high proliferation of goblet cells throughout the whole gill blades and filaments, along with partial regression of the filaments and signs of inflammation (Fig. 9. D, E). Exposure to 5 mg/L of Al_2O_3 induced focal thickening of the ctenidium epithelium and filaments (Fig. 9. H, F). At all NPs exposure concentrations, there was partial hypertrophy of the central cavity of the ctenidium and/or ospharidium, accompanied by an increase in mucus cells compared to the control. (See Fig. 9)




Fig. 11. The trend variation of biomarkers (GSH, GST, GPx, CAT, SOD, MDA, AChE, and MT) in the gills of *Stramonita haemastoma* exposed to Al_2O_3 and Cr_2O_3 NPs through exposure durations (* $p < 0.05$).

4. Discussion

In the present study, the toxicity of both NPs was assessed over time in the gills of the marine snail *Stramonita haemastoma* using various indicators. These indicators were employed to evaluate antioxidant responses, oxidative stress, cellular damage, disruption of neuronal activity, and structural changes at the tissue level, providing a comprehensive understanding of NPs toxicity in this marine snail. Additionally, the size, shape, zeta potential, and other characterization of Cr_2O_3 and Al_2O_3 NPs were investigated as they are important factors that could enhance or reduce NPs stability in the medium and thus their toxicity (Mezni et al., 2018; Arvidsson et al., 2020). Both NPs showed zeta values allowing them to be more likely stable in the NSW with a zeta average size ranging from 200 nm to 700 nm for Cr_2O_3 , and relatively lower at a maximum of 300 nm for Al_2O_3 which could suggest their higher bioavailability. NSW in this study was used to mimic natural conditions and thus take into account the impact of natural organic matter (NOM) (Canesi et al., 2017).

Enzymatic and non-enzymatic biomarkers are crucial cellular antioxidant defenses against oxidative damages caused by NPs, they play a key role in the detoxification process which could be by their direct interaction with xenobiotic in biotransformation pathway, or indirectly via maintaining the redox system homeostasis and elimination of ROS (Valko et al., 2007; Forman et al., 2009; Board and Menon, 2013). Non-enzymatic biomarkers like GSH and MT

play an important role in NPs detoxification by scavenging ROS as anion superoxide O_2^- and other radicals like HO, exhibiting metallic binding properties, and NPs modulation, especially in the case of MT. GSH serves as an important subtraction for antioxidants. Any imbalance in these biomarkers serves as an indicator of NPs toxicity (Saad et al., 2016; Gulcin, 2020; Roma et al., 2020). Enzymatic biomarkers that trigger or inhibit are considered indicators of NPs toxicity occurrence mainly through oxidative stress induction. The primary function of GST, GPx, CAT, and SOD is the removal of many types of ROS, with each antioxidant targeting specific types of ROS. Furthermore, AChE is crucial in neurotransmission by inhibiting ACh and has been used as a marker of neurotoxicity in several studies (Sedrati et al., 2024). In the same way, MDA is a molecule resulting from the lipidic peroxidation process triggered by oxidative stress, and it is also been widely applied as an effective marker of NPs toxicity (Hu and Gao, 2010; Shaw and Handy, 2011; Brigelius-Flohé and Maiorino, 2013; Vale et al., 2016).

The exposure of *S. haemastoma* to both types of NPs had varied effects on biomarkers in the gills, depending on the concentrations and duration of exposure. Overall, after 7 days, Cr_2O_3 NPs caused MT, AChE, and SOD activities to decrease with MDA levels to increase, on the other hand, Al_2O_3 NPs induced GST, CAT activities, GSH, MDA rates to increase, SOD activity and MT rates to decrease. Suggesting the response of marine snails to NPs exposure by activation of antioxidant defenses to counteract the mitigated oxidative stress indicated by MDA rates increase due to Al_2O_3 NPs (Fig. 5) as

additionally, IBR results confirmed the high induction of MDA rates (Fig. 8&9) and AChE inhibition due to Cr₂O₃ NPs (Fig. 6) exposure. MT levels reduction point also the detoxification process of NPs. Puerari et al. (2016) reported the induction of ROS-mediated toxicity by Cr₂O₃ in *Daphnia Magna*. Our results observed in the IBR analysis indicating the sensitivity of SOD, CAT activities and MDA rates to Cr₂O₃ NPs (Fig. 8) were similar to the findings of Kanwal et al. (2019) in fish *Labeo rohita* by CAT, SOD decrease, and MDA induction due to 25 mg/L of Cr₂O₃. Exposure to high concentrations of Al₂O₃ was found to decrease the levels of SOD while increasing the levels of CAT and MDA in *D. magna* (Nogueira et al., 2020a, 2020b). Additionally, the induction of GST, MDA, CAT, and SOD was reported in the fish *Carassius auratus*, at low concentrations (10 and 100 µg/L) after 7 days, while our results indicated that SOD activity was mostly decreased at all exposure concentrations for both NPs (Benavides et al., 2016). Furthermore, in zebrafish, *Danio rerio*, and clam *Ruditapes decussatus*, the levels of CAT activity, MDA rates, and ROS content increased due to exposure to Ag, ZnO, SiO₂, and TiO₂ NPs (Saidani et al., 2019; Rashidian et al., 2023). The study by Nunes et al. (2020) reported an increase in GST, CAT activities, and MDA contents accompanying the formation of NPs-mucus filaments in the gills of the mussel *Perna perna* following exposure to TiO₂ (48 h). In this study, similar phenomena were observed with microfilaments with NPs accumulation, which was clearer in the case of Cr₂O₃ potentially due to their green pigmentation. This observation is, to our best knowledge, the first report of such a case in gastropods (Fig. 5 in supplementary) and could indicate NPs bioavailability. Similar to the observed results in this study, De Oliveira et al. (2014) and Katuli et al. (2014) reported AChE activity inhibition in zebrafish *D. rerio* after acute exposure to high doses (20–200 mg/kg) of iron oxide NPs (IOs NPs) and sub-acute exposure to Ag NPs. Moreover, Nelson et al. (2018) reported neurotoxic effects on the gills functions of bivalves due to metallic traces.

On day 14, both NPs induced a decrease in GST, SOD activities (Fig. 4&5), and MT rates with AChE activity decrease due to Cr₂O₃ exposure (Fig. 6). GSH levels increased due to Al₂O₃ exposure while Cr₂O₃ caused GSH content depletion at 2.5 mg/L (Fig. 3), MDA rates showed lower rates (Figs. 5, 11). IBR results for the same identified the higher response of CAT activity, GSH, and MT rates to Cr₂O₃ exposure (Fig. 7), IBR results indicated also the sensitivity of MT rates and AChE activity to Al₂O₃ exposure (Fig. 8) which further could refer to the limitation of standard classical biomarkers evaluation. These results could suggest the onset of adaptive mechanisms in response to oxidative damage. Continuous decreases in AChE activity and MT levels, variations in GSH, rates, and decrease in GST activity could indicate enzymatic consumption and detoxification process of NPs, while the decrease in SOD further indicates the high production of O²⁻ resulting in cellular redox instability. The differential response of biomarkers for each NP exposure highlights the complexity of gastropods' interaction in response to pollutant contamination. The up and down-regulation of antioxidant system genes were reported, with similar findings in the digestive gland of *Mytilus galloprovincialis* in response to TiO₂ exposure (Gornati et al., 2016). Marine organisms possess a diversity of complex mechanisms and biochemical cascades as an adaptation to oxidative stress, from gene expression to catalytic activities that could vary in function of time and exposure doses (Regoli and Giuliani, 2014). Federici et al. (2007) reported GSH rates induction and NPs accumulation after 14-day exposure to 1 mg/L of TiO₂ NPs in the gills of Rainbow trout *Oncorhynchus mykiss*, consistent with the finding of Ray et al. (2020) which reported SOD and CAT activities reduction after 7 and 14 exposure days to CuO NPs in the gills of the mussel *Lamellidens marginalis* at 0.5 to 5 mg/L. Cordeiro et al. (2021) reported the higher impact of TiO₂ NPs (1 mg/L) on the digestive gland of the shrimp *Litopenaeus vannamei* compared to their gills, where GST activity

decreased and GSH content induction was reported. Ramsden et al. (2013) reported GSH rate induction in the gills of *D. rerio* following 1 mg/L exposure to TiO₂ NPs for 14 days.

Overall, at the end of the exposure period (28 days), both types of NPs induced various changes in biomarkers response, showing some similarities and differences. For instance, Cr₂O₃ NPs caused increased GSH content, GPx, CAT (Fig. 3&4), and AChE activities (Fig. 6) compared to the control, especially at higher exposure concentrations. The group exposed to Al₂O₃ NPs had increased activities of GSH, GST (Fig. 3), MT (Fig. 6), and MDA (Fig. 6) rates in some exposure concentrations, along with a decrease in CAT activity (Fig. 4). IBR results indicated the sensitivity and response of GST, CAT, GPx, AChE, MT and MDA contents with the highest response factor recorded at 2.5 mg/L for Cr₂O₃ (Fig. 7) and in a dose-dependent manner for Al₂O₃ (Fig. 8). These results suggest that oxidative damages accumulated over time, leading to instability in the redox system. The imbalance in antioxidants and MDA levels further confirms that the defensive systems were unable to preserve the cellular stability from the action of ROS and further indicates that gills are a potential target of NPs toxicity. The induction of GST activity and GSH rates suggests the continuous presence of NPs and their uptake in gills thus acting in the detoxification process which could potentially contribute to the release of NPs ions (Caixeta et al., 2020; Dube and Okuthe, 2023). Huang et al. (2018) found that exposure to TiO₂ exposure (2.5 mg/L–10 mg/L) in the mussel *Mytilus coruscus* caused CAT, GPx, GST activity induction, and an increase in MDA rates, along with SOD activity inhibition. Similarly, another study by Canesi et al. (2010b) reported that TiO₂ exposure (0.05–5 mg/L) caused CAT and GST activity induction in the gills of *M. galloprovincialis*. Consistent with our findings, other studies have reported MT level induction, depletion, and gene-up regulations in bivalves due to NPs exposure, indicating that MT imbalance is a key marker of NPs toxicity, as our results indicated the ability of both studied NPs to disturb MT rates (Buffet et al., 2011 and 2012; Gomes et al., 2011 and 2012). The exposure to TiO₂, CuO, and Al₂O₃ induced similar effects in biomarkers with lesser impact for Al₂O₃. This included CAT, SOD activity decrease, and GST activity induction in the digestive gland and gills of the mussel *Unio tigris* (Canli and Canli, 2021). Al₂O₃ accumulation into gills was also reported in *U. tigris* (Canli et al., 2022), while Al₂O₃ was reported to induce higher toxicity than other NPs, including SOD activity decrease and GPx activity induction in rats (Canli et al., 2019). Additionally, neurotoxic effects were reported in *D. rerio* due to Al₂O₃ NPs, including (ACh) inhibition, Al₂O₃ NPs accumulation, and memory capacity alteration resulting from oxidative damage (Chen et al., 2020). The different responses observed of some biomarkers to each type of NPs for instance SOD, GPx and, CAT could be interpreted as due potentially to the intracellular pathways and locations in which those NPs could induce oxidative stress and consequently unbalancing biomarkers activities and rates.

Histological assessment is a crucial and reliable tool that could provide evidence of NPs toxicity and impacts on the cellular to tissue level (Cid et al., 2015; Leite et al., 2020). In the present study, the histological evaluation results showed clear evidence of alterations in the gills of *S. haemastoma* for both NPs. Various histological damage was observed, ranging from lamella fusion to gills central cavities hypertrophy with signs of inflammation and tissue destruction. The gills of the marine snails were sensitive to both NPs concentrations and, after 28 days, showed strong evidence of progressive histological alteration. The findings further imply the failure of the redox system defenses in the preservation from oxidative damage and suggest that both Cr₂O₃ and Al₂O₃ NPs were uptake and translocated through the gills of *S. haemastoma*, thus, inducing harmful effects by direct interaction along cell membranes or by their internalization (Canesi et al., 2008; Bonany et al., 2022). The

present results are consistent with previously reported histological findings as several studies noted gills histological damage in bivalves (Cuevas et al., 2015; Canli et al., 2022), gastropods (Arrighetti et al., 2022), fish (Xiong et al., 2011), and *Daphnia magna* (Puerari et al., 2016) due to NPs contamination. Cr₂O₃ NPs were reported to induce inflammation and edema in the liver and kidneys of *Labeo rohita* after 21 days of exposure (Kanwal et al., 2019), and cellular dilatation and degeneration were also reported in rats (Fatima and Ahmad, 2019). Al₂O₃ NPs induced Lipofuscin aggregates, lamellar fusion, and hemolytic infiltration following 96 h exposure in the gills of *M. galloprovincialis* (Ertürk Gürkan and Gürkan, 2021). Similarly to our findings, Al₂O₃ was reported to induce lamella fusion and hyperplasia in the gills of *Carassius auratus*, at low concentrations (10 and 100 µg/L) (Benavides et al., 2016), necrosis, hyperplasia, and inflammation were also observed in freshwater fish *Oreochromis mossambicus* (Murali et al., 2018). The results obtained in the present studies pointed to the ability of both NPs to induce a large diversity of permanent histological damages and further prove that NPs have the potential to decrease the health status of aquatic ecosystems including marine ones, for instance, NPs concentrations will inevitably rise with their increased applications by time thus, may cause direct harmful effects to aquatic organisms or indirect one as in the case of trophic web translocations.

The usage of a multi-biomarkers approach to evaluate the pollutant's toxicity is crucial. However, interpreting various biomarkers responses to contamination independently can be challenging, and researchers face a dilemma in achieving proper results interpretation.

The integrated biomarkers response (IBR) is a valuable tool for evaluating trends in oxidative stress biomarkers (Beliaeff and Burgeot, 2002; Sanchez et al., 2013; Roma et al., 2024). The IBRv2 analysis provides an integrative assessment of multiple biomarkers, allowing us to evaluate the overall stress response of snail organisms to nanoparticle (NPs) exposure in a comprehensive manner. Unlike the individual biomarker analyses, which focus on specific biochemical pathways or enzymatic activities, the IBRv2 combines these responses into a single index, offering a broader perspective on the cumulative effects of stressors. This holistic approach adds significant value by facilitating comparisons across experimental groups and identifying patterns that may not be apparent from individual biomarker data alone. In this study, IBR was used to further clarify and validate results obtained from assessing numerous biomarkers. Briefly, the results of IBR showed clear responses of biomarkers to different concentrations for both NPs and exposure periods. Specifically at day 7 and day 14, IBRv2 indicated that both NPs have a significant impact on biomarkers, with higher response observed in 2 to 3 biomarkers, namely after 7 days SOD, AChE, and MDA due to Cr₂O₃ NPs exposure while the most sensitive biomarkers due to Al₂O₃ exposure were SOD, MT, and MDA. Following 14 exposure days, IBRv2 results indicated that AChE and MT were the most sensitive to Cr₂O₃ NPs exposure, and SOD, MT, MDA due to Al₂O₃ exposure. At 28 days, there was a higher global response in nearly all biomarkers, indicating the onset of oxidative stress and the activation of mechanisms to reduce damage. The results for both NPs confirm that effects triggered at 7 days were slightly reduced at day 14, suggesting the early response, with the activation of defensive mechanisms, and induced higher values at day 28, indicating cumulative oxidative damages causing coregulation of all biomarkers' action IBRv2 global evolution according to exposure periods and concentrations are shown in Fig. 10. Such results could not be detected when evaluating biomarkers individually, highlighting the usefulness of IBR in the evaluation NPs toxicity. To our best knowledge, the present study is the first to assess NPs toxicity using IBR in marine gastropods. While IBR remains poorly used in the literature, several related studies have reported

its successful application, primarily to bivalves (Xia et al., 2013; Devin et al., 2017; Liu et al., 2017; Li et al., 2021; Gonçalves et al., 2022; Roma et al., 2024). Previous studies have reported time and dose-dependent impacts on IBR of SOD, CAT, AChE, and MDA Xia et al. (2017) following exposure to 1 mg/L of TiO₂ NPs in the gills and digestive glands of the clam *C. farreri*. The exposure to high concentrations of ZnO₂ NPs (1 to 100 mg/L) showed high values after 5 days and then reduced IBR values in fluvial biofilm bacteria of GPx, CAT, SOD, and GR (Hou et al., 2016). The present study differs from previous ones as we use a variety of biomarkers, including AChE and MT responses.

Several studies have identified ion release as a primary mechanism for increasing ROS production (Yung et al., 2014; Peng et al., 2017). Previous studies indicate that the toxicity of Cr₂O₃ NPs is attributed to the liberation of Hexavalent Cr ions in A549 cells and other models, leading to constantly increased intra and extracellular ROS levels (Horie et al., 2013; Alarifi et al., 2016). In the same way, da Costa et al. (2015) reported higher ROS production due to Cr₂O₃ NPs exposure in the green alga *Chlamydomonas reinhardtii*. The results observed in this study indicate the uptake of both NPs in the gills of *S. haemastoma*, as gills are responsible for ensuring respiratory function in marine snails, the contamination of both NPs caused their direct contact with gills. The uptake of NPs agglomeration mainly through the gills of bivalves and then their translocation to the digestive gland was previously identified as a main pathway of NPs accumulation (Koehler et al., 2008; Ciacci et al., 2012; Volland et al., 2015). Internalization of NPs could occur and be influenced by several factors, including NPs size, zeta average diameter, shape, and concentration (Barmo et al., 2013). Furthermore, NPs toxicity could be amplified in the environment, as interaction with other pollutants and their absorption on NPs' surface, significantly increases their toxicity. Conversely, this aspect should also be considered in future studies. The present study addressed this divergence by using NSW, as potential interactions could be important factors (Canesi et al., 2017; Latchere et al., 2023a, 2023b).

5. Conclusion

In this present work, the toxicity of Cr₂O₃ and Al₂O₃ NPs was investigated over time in the gills of the marine gastropod *Stramonita haemastoma*. The findings provided important novel insights into the induction of oxidative stress by NPs and the adaptive mechanisms of marine snails. Both NPs were able to induce various changes in the antioxidant system, with the most marked effects observed in SOD, CAT, AChE, MT, and MDA. In the same way, histological evaluation showed clear evidence of both NPs' bioavailability in the gills of snails as dose-dependent alterations including inflammation, hypertrophy, and lamella fusion signs suggesting that Cr₂O₃ and Al₂O₃ NPs could translocate through gills in marine gastropods. The use of IBR in this study was crucial for identifying NPs toxicity patterns and further provided evidence of the time-dependent impact of exposure on biomarkers and induction of oxidative stress. IBR was applied to a battery of biomarkers and appeared to be a potentially ideal tool to assess NPs toxicity and compare impact trends in marine gastropods. The methodological contribution of this paper could be extended to future related studies. Additionally, the sensitivity of *S. haemastoma* to NPs exposure validates this gastropod as a promising bioindicator of NPs contamination in marine ecosystems. Future extended toxicological evaluations are required to elucidate NPs uptake, translocation, and toxicity mechanisms at the molecular level, especially for Cr₂O₃ which has received less attention compared to other NPs in ecotoxicological studies. We strongly suggest and recommend a more expanded assessment of the actual coastal NPs contamination especially regarding specific NPs as

Cr₂O₃ due to the lack of studies involving this type of NPs and others compared to their economic importance. Another important crucial point to consider in perspective is as discussed the combined impacts and interactions among several types of NPs and other pollutants and toxins.

CRediT authorship contribution statement

Fateh Sedrati: Writing – review & editing, Writing – original draft, Methodology, Investigation, Conceptualization. **Hana Bouzahouane:** Visualization, Validation, Supervision, Methodology. **Mohcen Menaa:** Software, Formal analysis, Data curation. **Fadila Khaldi:** Validation, Supervision. **Tayeb Bouarroudj:** Software, Investigation, Formal analysis. **Lassaad Gzara:** Visualization, Formal analysis. **Mounira Bensalem:** Resources, Investigation. **Omar Laouar:** Methodology, Investigation. **Noomene**

Appendix A. Supplementary data

Supplementary data to this article can be found online at <https://doi.org/10.1016/j.cbpc.2025.110159>.

Funding declaration

This research did not receive any specific grant from funding agencies in the public, commercial, or not-for-profit sectors. Carla O. Silva acknowledges the independent funding received from Fundação para a Ciência e a Tecnologia (FCT I.P., Portugal) within the framework of the UID/04292/MARE-Centro de Ciências do Mar e do Ambiente and the project LA/P/0069/2020 (<https://doi.org/10.54499/LA/P/0069/2020>) granted to the Associate Laboratory ARNET - Aquatic Research Network

References

- Abdel-Latif, H.M.R., Dawood, M.A.O., Menaiea-Le double, S., El-Matbouli, M., 2020. Environmental transformation of n-TiO₂ in the aquatic systems and their ecotoxicity in bivalve mollusks: a systematic review. *Ecotoxicol. Environ. Saf.* 200. <https://doi.org/10.1016/j.ecoenv.2020.110776>.
- Abraham, N., Mashkin, P., Volkov, S., Olshanskiy, V., Kustov, L., 2021. Fresh-water mollusks as biomonitors for ecotoxicity of nanomaterials. *Nanomaterials* 11 (4). <https://doi.org/10.3390/nano11040944>.
- Adnan, W.G., Mohammed, A.M., 2024. Green synthesis of chromium oxide nanoparticles for anticancer, antioxidant, and antibacterial activities. *Inorg. Chim. Commun.* 159, 111683. <https://doi.org/10.1016/j.inoche.2023.111683>.
- Alarifi, S., Ali, D., Alkahtani, S., 2016. Mechanistic investigation of toxicity of chromium oxide nanoparticles in murine fibroblast cells. *Int. J. Nano Medicine* 11, 1253–1259. <https://doi.org/10.2147/IJN.S99995>.
- Arrighetti, F., Landro, S.M., Lavarias, S.M.L., 2022. Sensitivity of histopathological and histoc hemical parameters in the digestive gland of the apple snail *Pomacea canaliculata* exposed to cypermethrin. *Aquat. Toxicol.* 252, 106292. <https://doi.org/10.1016/j.aquatox.2022.106292>.
- Arvidsson, R., Hansen, S.F., Baun, A., 2020. Influence of natural or organic matter on the aquatic ecotoxicity of engineered nanoparticles: a review. *Nano Impact* 20. <https://doi.org/10.1016/j.nimpac.2020.100263>.
- Ates, M., Demir, V., Arslan, Z., Daniels, J., Farah, I.O., Bogatu, C., 2015. Evaluation of alpha and gamma aluminum oxide nanoparticle accumulation, toxicity, and depuration in *Artemia salina* larvae: Al₂O₃ NP and *Artemia Salina* larvae. *Environ. Toxicol.* 30 (1), 109–118. <https://doi.org/10.1002/tox.21917>.
- Baker, T.J., Tyler, C.R., Galloway, T.S., 2014. Impacts of metal and metal oxide nanoparticles on marine organisms. *Environ. Pollut.* 186, 257–271. <https://doi.org/10.1016/j.envpol.2013.11.014>.
- Barm, C., Ciacci, C., Canonic, B., Fabbri, R., Cortese, K., Balbi, T., Marcomini, A., Pojana, G., Gallo, G., Canesi, L., 2013. In vivo effects of n-TiO₂ on the digestive gland and immune function of the marine bivalve *Mytilus galloprovincialis*. *Aquat. Toxicol.* 132–133, 9–18. <https://doi.org/10.1016/j.aquatox.2013.01.014>.
- Becker, L.C., Boyer, I., Bergfeld, W.F., Belsito, D.V., Hill, R.A., Klaassen, C.D., Liebler, D.C., Marks, J.G., Shank, R.C., Slaga, T.J., Snyder, P.W., Andersen, F.A., 2016. Safety assessment of aluminum and aluminum hydroxide as used in cosmetics. *Int. J. Toxicol.* 35 (3 suppl), 16S–16S33S. <https://doi.org/10.1177/1091581816677948>.
- Beer, R.F., Sizer, I.W., 1952. A spectrophotometric method for measuring the breakdown of hydrogen peroxide by catalase. *J. Biol. Chem.* 195 (1), 133–140.
- Sleimi, Resources. Hichem Nasri: Resources. Carla O. Silva: Writing – review & editing, Writing – original draft, Visualization. Kheirddine Ouali: Validation, Resources.
- ## Uncited references
- ## Declaration of competing interest
- The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.
- ## Data availability
- Data will be made available on request.

- Brigelius-Flohé, R., Maiorino, M., 2013. Glutathione peroxidases. *Biochim. Biophys. Acta Gen. Subj.* 1830 (5), 3289–3303. <https://doi.org/10.1016/j.bbagen.2012.11.020>.
- Buffet, P.E., Amiard-Tricket, C., Dybowska, A., Risso-de Favey, C., Guibolini, M., Valsami-Jones, E., Moueyrac, C., 2012. Fate of isotopically labeled zinc oxide nanoparticles in sediment and effects on two endobenthic species, the clam *Scrobicularia plana*, and the ragworm *Mediste diversicolor*. *Environ. Saf.* 84, 191–198. <https://doi.org/10.1016/j.ecoenv.2012.07.010>.
- Buffet, P.E., Tankou, O.F., Pan, J.F., Berhanu, D., Herrenknecht, C., Poirier, L., Amiard-Tricket, C., Amiard, J.C., Bérard, J.B., Risso, C., Guibolini, M., Roméo, M., Reip, P., Valsami-Jones, E., Moueyrac, C., 2011. Behavioral and biochemical responses of two marine invertebrates *Scrobicularia plana* and *Mediste diversicolor* to copper oxide nanoparticles. *Chemosphere* 84 (1), 166–174. <https://doi.org/10.1016/j.chemosphere.2011.02.003>.
- Caixeta, M.B., Araújo, P.S., Gonçalves, B.B., Silva, L.D., Granato-Maldonado, M.I., Rocha, T.L., 2020. Toxicity of engineered nanomaterials to aquatic and land snails: a scientometric and systematic review. *Chemosphere* 260. <https://doi.org/10.1016/j.chemosphere.2020.127654>.
- Canesi, L., Balbi, T., Fabbri, R., Salis, A., Damonte, G., Volland, M., Blasco, J., 2017. Biomolecular coronas in invertebrate species: implications in the environmental impact of nanoparticles. *Nano Impact* 8, 89–98. <https://doi.org/10.1016/j.nimp.2017.08.001>.
- Canesi, L., Ciacci, C., Balbi, T., 2015. Interactive effects of nanoparticles with other contaminants in aquatic organisms: friend or foe? *Mar. Environ. Res.* 111, 128–134. <https://doi.org/10.1016/j.marenvres.2015.03.010>.
- Canesi, L., Ciacci, C., Betti, M., Fabbri, R., Canonic, B., Fantinati, A., Marcomini, A., Pojana, G., 2008. Immunotoxicity of carbon black nanoparticles to blue mussel hemocytes. *Environ. Int.* 34 (8), 1114–1119. <https://doi.org/10.1016/j.envint.2008.04.002>.
- Canesi, L., Ciacci, C., Fabbri, R., Marcomini, A., Pojana, G., Gallo, G., 2012. Bivalve molluscs as a unique target group for nanoparticle toxicity. *Mar. Environ. Res.* 76, 16–21. <https://doi.org/10.1016/j.marenvres.2011.06.005>.
- Canesi, L., Ciacci, C., Vallotto, D., Gallo, G., Marcomini, A., Pojana, G., 2010a. *In vitro* effects of suspension of selected nanoparticles (C60 fullerene, TiO₂, SiO₂) on *Mytilus hemocytes*. *Aquat. Toxicol.* 96 (2), 151–158. <https://doi.org/10.1016/j.aquatox.2009.10.017>.
- Canesi, L., Fabbri, R., Gallo, G., Vallotto, D., Marcomini, A., Pojana, G., 2010b. Biomarkers in *Mytilus galloprovincialis* exposed to suspension of selected nanoparticles (Nano carbon black, C60 fullerene, Nano-TiO₂, Nano-SiO₂). *Aquat. Toxicol.* 100 (2), 168–177. <https://doi.org/10.1016/j.aquatox.2010.04.009>.
- Canli, E.G., Canli, M., 2021. Antioxidant system biomarkers of freshwater mussel (*Unio tigris*) respond to nanoparticle (Al₂O₃, CuO, TiO₂) exposure. *Biomarkers* 26 (5), 434–442. <https://doi.org/10.1080/1354750X.2021.1909655>.
- Canli, E.G., Celenk, A., Canli, M., 2022. Accumulation and distribution of nanoparticles (Al₂O₃, CuO, and TiO₂) in tissues of freshwater mussel (*Unio tigris*). *Bull. Environ. Contam. Toxicol.* 108 (4), 702–707. <https://doi.org/10.1007/s0128-021-03410-5>.
- Canli, E.G., Ila, H.B., Canli, M., 2019. Response of the antioxidant enzymes of rats following oral administration of metal-oxide nanoparticles (Al₂O₃, CuO, TiO₂). *Environ. Sci. Pollut. Res.* 26 (1), 938–945. <https://doi.org/10.1007/s1356-018-3592-8>.
- Chen, J., Fan, R., Wang, Y., Huang, T., Shang, N., He, K., Zhang, P., Zhang, L., Niu, Q., Zhang, Q., 2020. Progressive impairment of learning and memory in adult zebrafish treated by Al₂O₃ nanoparticles when in embryonic. *Chemosphere* 254. <https://doi.org/10.1016/j.chemosphere.2020.126608>.
- Chiu, D.T.Y., Stults, F.H., Tappell, A.L., 1976. Purification and properties of rat lung soluble glutathione peroxidase. *Biochimica et Biophysica Acta (BBA) - Enzymology* 445 (3), 558–566. [https://doi.org/10.1016/0005-2744\(76\)90110-8](https://doi.org/10.1016/0005-2744(76)90110-8).
- Ciacci, C., Canonic, B., Bilaničová, D., Fabbri, R., Cortese, K., Gallo, G., Marcomini, A., Pojana, G., Canesi, L., 2012. Immunomodulation by different types of N-oxides in the hemocytes of the marine bivalve *Mytilus galloprovincialis*. *PLoS One* 7 (5). <https://doi.org/10.1371/journal.pone.0036937>.
- Cid, A., Picaud, A., Correia, J.B., Chaves, R., Silva, H., Caldeira, J., de Matos, A.P.A., Diniz, M.S., 2015. Oxidative stress and histological changes following exposure to diamond nanoparticles in the freshwater Asian clam *Corbicula fluminea* (Müller, 1774). *J. Hazard. Mater.* 284, 27–34. <https://doi.org/10.1016/j.jhazmat.2014.10.055>.
- Coelho, L., Müller, L., Manske Nunes, S., Kist, L.W., Bogo, M.R., Ruas, C.P., Gulesky, M., Wasielesky, W., Fattorini, D., Regoli, F., Monserrat, J.M., Ventura-Lima, J., 2021. Co-exposure to nTiO₂ impairs arsenic metabolism and affects antioxidant capacity in the marine shrimp *Litopenaeus vannamei*. *Drug Chem. Toxicol.* 44 (1), 30–38. <https://doi.org/10.1080/01480545.2018.1563610>.
- Corsi, I., Bergami, E., Grassi, G., 2020. Behavior and bio-interactions of anthropogenic particles in marine environment for a more realistic ecological risk assessment. *Front. Environ. Sci.* 8, 60. <https://doi.org/10.3389/fenvs.2020.00060>.
- Cuervas, N., Zorita, I., Costa, P.M., Franco, J., Larreta, J., 2015. Development of histopathological indices in the digestive gland and gonad of mussels: integration with contamination levels and effects of confounding factor. *Aquat. Toxicol.* 162, 152–164. <https://doi.org/10.1016/j.aquatox.2015.03.011>.
- da Costa, C.H., Perrault, F., Ouakroum, A., Melegari, S.P., Popovic, R., Matias, W.G., 2015. Effect of chromium oxide (III) nanoparticles on the production of reactive oxygen species and photosystem II activity in the green alga *Chlamydomonas reinhardtii*. *Sci. Total Environ.* 565, 951–960. <https://doi.org/10.1016/j.scitotenv.2016.01.028>.
- De Felice, B., Parolini, M., 2020. Can proteomics be considered as a valuable tool to assess the toxicity of nanoparticles in marine bivalves? *Journal of Marine Science and Engineering* 8 (12), 1–16. <https://doi.org/10.3390/jmse812033>.
- De Oliveira, G.M.T., Kist, L.W., Pereira, T.C.B., Bortolotto, J.W., Paquete, F.L., De Oliveira, E.M.N., Leite, C.E., Bonan, C.D., De Souza Basso, N.R., Papaleo, R.M., Bogo, M.R., 2014. Transient modulation of acetylcholinesterase activity caused by exposure to dextran-coated iron oxide nanoparticles in brain of adult zebrafish. *Comp. Biochem. Physiol. Part C: Toxicol. Pharmacol.* 162, 77–84. <https://doi.org/10.1016/j.cbpc.2014.03.010>.
- Deng, Y., Yang, F., Cocco, E., Song, E., Zhang, J., Cui, J., Mohideen, M., Bellone, S., Santin, A.D., Saltzman, W.M., 2016. Improved i.p. drug delivery with bioadhesive nanoparticles. *Proc. Natl. Acad. Sci.* 113 (41), 11453–11458. <https://doi.org/10.1073/pnas.1523141113>.
- Devlin, S., Buffet, P.E., Châtel, A., Perrin-Etjani, H., Valsami-Jones, E., Moueyrac, C., 2017. The integrated biomarker response: a suitable tool to evaluate toxicity of metal-based nanoparticles. *Nanotoxicology* 11 (1), 1–6. <https://doi.org/10.1080/17435390.2016.1269374>.
- Devlin, S., Burgeot, T., Giamberini, L., Minguez, L., Païn-Devlin, S., 2014. The integrated biomarker response revisited: optimization to avoid misuse. *Environ. Sci. Pollut. Res.* 21 (4), 2448–2454. <https://doi.org/10.1007/s1356-013-2169-9>.
- Di Bella, G., Pizzullo, G., Bua, G.D., Portoli, A.G., Santini, A., Giacobbe, S., 2018. Mapping toxic mineral contamination: the southern oyster drill, *S. Haemastoma* (L., 1767), as evaluable sentinel species. *Environ. Monit. Assess.* 190 (1), 7. <https://doi.org/10.1007/s10661-017-6380-x>.
- Dube, E., Okuthe, G.E., 2023. Engineered nanoparticles in aquatic systems: toxicity and mechanism of toxicity in fish. *Emerging Contaminants* 9 (2), 100212. <https://doi.org/10.1016/j.emcon.2023.100212>.
- El Ayari, T., Bierne, N., El Menif, N.T., 2018. Imposex incidence in *Stramonita haemastoma* (Gastropoda: Muricidae) from the Mediterranean and Atlantic coast after tributyltin global ban. *J. Sea Res.* 134, 10–15. <https://doi.org/10.1016/j.seares.2017.12.004>.
- Ellman, G.L., Courtney, K.D., Andres, V., Featherstone, R.M. (1961). A NEW AND RAPID COLORIMETRIC DETERMINATION OF ACETYLCHOLINESTERASE ACTIVITY (Biochemical Pharmacology, Vol. 10. 1st ed., pp. 88–95). Pergamon Press Ltd.
- Ertekin, G., Gürk, M., Gürk, M., 2021. Toxicity of gamma aluminum oxide nanoparticles in the Mediterranean mussel (*Mytilus galloprovincialis*): histopathological alterations and antioxidant responses in the gill and digestive gland. *Biomarkers* 26 (3), 248–259. <https://doi.org/10.1080/1354750X.2021.1878558>.
- Fan, X., Wang, C., Wang, P., Hu, B., Wang, X., 2018. TiO₂ nanoparticles in sediments: effect on the bioavailability of heavy metals in the freshwater bivalve *Corbicula fluminea*. *J. Hazard. Mater.* 342, 41–50. <https://doi.org/10.1016/j.jhazmat.2017.07.041>.
- Fatima, M., Ahmad, I., Sayed, I., Athar, M., Raisuddin, S., 2000. Pollutants induce over-activation of phagocytes is concomitantly associated with peroxidative damage in fish tissues. *Aquatic Toxicology* 49, 243–250. www.elsevier.com/locate/aquatox.
- Fatima, R., Ahmad, R., 2019. Hepatotoxicity and chromosomal abnormalities evaluation due to single and repeated oral exposures of chromium oxide nanoparticles in Wistar rats. *Toxicol. Ind. Health* 35 (8), 548–557. <https://doi.org/10.1177/074823719863632>.
- Federici, G., Shaw, B., Handy, R., 2007. Toxicity of titanium dioxide nanoparticles to rainbow trout (*Oncorhynchus mykiss*): gill injury, oxidative stress, and other physiological effects. *Aquat. Toxicol.* 84 (4), 415–430. <https://doi.org/10.1016/j.aquatox.2007.07.009>.
- Metal nanoparticles global market to reach \$4.2 bn by 2026 at a CAGR of 11.5%. (2023). Focus on Catalysts, 2023(1), 4. doi:<https://doi.org/10.1016/j.focac.2022.12.013>.
- Forman, H.J., Zhang, H., Rinn, A., 2009. Glutathione: overview of its protective roles, measurement, and biosynthesis. *Mol. Asp. Med.* 30 (1–2), 1–12. <https://doi.org/10.1016/j.mam.2008.08.006>.
- García-Saucedo, C., Field, J.A., Otero-González, L., Sieral-Alvarez, R., 2011. Low toxicity of HfO₂, SiO₂, Al₂O₃ and CeO₂ nanoparticles to the yeast *Saccharomyces cerevisiae*. *Journal of Hazardous Materials* 192 (3), 1572–1579. <https://doi.org/10.1016/j.jhazmat.2011.06.081>.
- Gomes, T., Pereira, C.G., Cardoso, C., Pinheiro, J.P., Canicio, I., Bebianno, M.J., 2012. Accumulation and toxicity of copper oxide nanoparticles in the digestive gland of *Mytilus galloprovincialis*. *Aquat. Toxicol.* 118–119, 72–79. <https://doi.org/10.1016/j.aquatox.2012.03.017>.
- Gomes, T., Pinheiro, J.P., Canicio, I., Pereira, C.G., Cardoso, C., Bebianno, M.J., 2011. Effects of copper nanoparticles exposure in the mussel

- Mytilus galloprovincialis. *Environ. Sci. Technol.* 45 (21), 9356–9362. <https://doi.org/10.1021/es200955s>.
- Gonçalves, J.M., Sousa, V.S., Teixeira, M.R., Bebianno, M.J., 2022. Chronic toxicity of polystyrene nanoparticles in the marine mussel *Mytilus galloprovincialis*. *Chemosphere* 287, 132356. <https://doi.org/10.1016/j.chemosphere.2021.132356>.
- Gornati, R., Longo, A., Rossi, F., Maisano, M., Sabatino, G., Mauceri, A., Bernadini, G., Fasulo, S., 2016. Effects of titanium dioxide nanoparticles exposure in *Mytilus galloprovincialis* gills and digestive gland. *Nano toxicology* 10 (6), 807–817. <https://doi.org/10.3109/17435390.2015.1132348>.
- Gulcin, I., 2020. Antioxidants and antioxidant methods: an updated overview. *Arch. Toxicol.* 94 (3), 651–715. <https://doi.org/10.1007/s00204-020-02689-3>.
- Habig, W.H., Pabst, M.J., And, ~, & Jakob, W.B. (1974). Glutathione S-Transferase: THE FIRST ENZYMIC STEP IN MERCAPTURIC ACID FORMATION* (THE JOURNAL OF Biochemistry, Vol. 73, No. 1, 249, Issue 22, pp. 7130–7139). <http://www.jbc.org/>
- Handy, R.D., Henry, T.B., Scowen, T.M., Johnston, B.D., Tyler, C.R., 2008. Manufacture of nanoparticles: their uptake and effects on fish—a mechanistic analysis. *Ecotoxicology* 17 (5), 396–409. <https://doi.org/10.1007/s10646-008-0205-1>.
- Hao, L., Chen, L., 2012. Oxidative stress responses in different organs of carp (*Cyprinus carpio*) with exposure to ZnO nanoparticles. *Ecotoxicol. Environ. Saf.* 80, 103–110. <https://doi.org/10.1016/j.ecoenv.2012.02.017>.
- Hichem, N., Hadjer, Z., Fath, S., Feriel, L., Wang, Z., 2022. The potential exposure and hazards of zinc nanoparticles: a review. *Ecotoxicology and environmental contamination* 17 (1), 1–21. <https://doi.org/10.5132/ee.2022.01.01>.
- Horie, M., Nishio, K., Endoh, S., Kato, H., Fujita, K., Miyachi, A., Nakamura, A., Kinugasa, S., Yamamoto, K., Niki, E., Yoshida, Y., Iwahashi, H., 2013. Chromium (III) oxide nanoparticles induce remarkable oxidative stress and apoptosis in cultured cells. *Environ. Toxicol.* 28 (2), 61–75. <https://doi.org/10.1002/tox.20695>.
- Hou, J., Wang, L., Wang, C., Zhang, S., Liu, H., Li, S., Wang, X., 2019. Toxicity and mechanisms of action of titanium dioxide nanoparticles in living organisms. *J. Environ. Sci. (China)* 75, 40–53. <https://doi.org/10.1016/j.jes.2018.06.010>.
- Hou, J., You, G., Xu, Y., Wang, C., Wang, P., Miao, L., Dai, S., Lv, B., Yang, Y., 2016. Antioxidant enzyme activities as biomarkers of fluvial biofilm to ZnO NP's ecotoxicity and the integrated biomarker responses (IBR) assessment. *Ecotoxicol. Environ. Saf.* 133, 10–17. <https://doi.org/10.1016/j.ecoenv.2016.06.014>.
- René Houliard. (1984). *Techniques d'histopathologie et de cytopathologie* (1984 Montré: Décarie, Ed.; p. 400 p).
- Hu, Y.L., Gao, J.Q., 2010. Potential neurotoxicity of nanoparticles. *Int. J. Pharm.* 394 (1–2), 115–121. <https://doi.org/10.1016/j.ijpharm.2010.04.026>.
- Huang, X., Liu, Z., Xie, Z., DuPont, S., Huang, W., Wu, F., Kong, H., Liu, L., Su, Y., Lin, D., Lu, W., Hu, M., Wang, Y., 2018. Oxidative stress induced by titanium dioxide nanoparticles increases under seawater acidification in the thick shell mussel *Mytilus coruscus*. *Mar. Environ. Res.* 137, 49–59. <https://doi.org/10.1016/j.marenvres.2018.02.029>.
- Kafi-Ahmad, L., Khademir, S., Pourshari, Marjan, A., Nozad, E., 2022. Microwave-assisted preparation of polystyrene nanoparticles using Zingiber extract synthesized green Cr2O3 nanoparticles. *Sci. Rep.* 12 (1). <https://doi.org/10.1038/s41598-022-24364-6>.
- Kanwal, Z., Raza, M.A., Manzoor, F., Riaz, S., Jabeen, G., Fatima, S., Naseem, S., 2019. A comparative assessment of nano toxicity induced by metal (silver, nickel) and metal oxide (cobalt, chromium) nanoparticles in *Laeborhita*. *Nanomaterials* 9 (2). <https://doi.org/10.3390/nano902309>.
- Katuli, K.K., Massarsky, A., Hadadi, A., Pourmehr, A., 2014. Silver nanoparticles inhibit the gill Na⁺/K⁺-ATPase and erythrocyte AChE activities and induce the stress response in adult zebrafish (*Danio rerio*). *Ecotoxicol. Environ. Saf.* 106, 173–180. <https://doi.org/10.1016/j.ecoenv.2014.04.001>.
- Khan, I., Saeed, K., Khan, I., 2019. Nanoparticles: properties, applications and toxicities. *Arab. J. Chem.* 12 (7), 908–931. <https://doi.org/10.1016/j.arabjc.2017.05.011>.
- Köhler, A., Marx, U., Bröck, K., Bahn, S., Bressling, J., 2008. Effects of nanoparticles in *Mytilus edulis* gills and hepatopancreas—a new threat to marine life? *Mar. Environ. Res.* 66 (1), 12–14. <https://doi.org/10.1016/j.marenvres.2008.02.009>.
- La Tchere, O., Roman, C., Métais, I., Perrin-Ettajani, H., Moulou, D., M., Georges, D., Ferrer-Mazel, A., Gigault, J., Castrouillet, C., Baudrimont, M., Châtel, A., 2023a. Toxicity assessment of environmental MPs and NPs and polystyrene NPs on the bivalve *Corbicula fluminea* using a multi-marker approach. *Comparative Biochemistry and Physiology Part C: Toxicology & Pharmacology* 273, 109714. <https://doi.org/10.1016/j.cbpc.2023.109714>.
- La Tchere, O., Roman, C., Métais, I., Perrin-Ettajani, H., Moulou, D., M., Georges, D., Ferrer-Mazel, A., Gigault, J., Castrouillet, C., Baudrimont, M., Châtel, A., 2023b. Toxicity assessment of environmental MPs and NPs and polystyrene NPs on the bivalve *Corbicula fluminea* using a multi-marker approach. *Comparative Biochemistry and Physiology Part C: Toxicology & Pharmacology* 273, 109714. <https://doi.org/10.1016/j.cbpc.2023.109714>.
- Leite, C., Coppola, F., Monteiro, R., Russo, T., Poiese, G., Lourenço, M.A.O., Silva, M.R.F., Ferreira, P., Soares, A.M.V.M., Freitas, R., Pereira, E., 2020. Biochemical and histopathological impacts of rutile anatase (TiO₂ form) in *Mytilus galloprovincialis*. *Sci. Total Environ.* 719. <https://doi.org/10.1016/j.scitotenv.2019.134886>.
- Li, L., Gu, H., Chang, X., Huang, W., Sokolova, I.M., Wei, S., Sun, L., Li, S., Wang, X., Hu, M., Zeng, J., Wang, Y., 2021. Oxidative stress induced by nanoparticles in the liver of juvenile largemouth bass (*Micropterus dolomieu*). *Mar. Pollut. Bull.* 170, 112661. <https://doi.org/10.1016/j.marpolbul.2021.112661>.
- Li, H., Zhang, W., Fang, Y., Yang, H., Tian, L., Li, K., Lai, W., Bian, L., Lin, B., Liu, X., Xi, Z., 2020. Neurotoxicity of aluminum oxide nanoparticles and their mechanistic role in dopaminergic neuron injury involving p53-related pathways. *J. Hazard. Mater.* 392. <https://doi.org/10.1016/j.jhazmat.2020.122312>.
- Liu, Y., Yan, Z., Xia, J., Wang, K., Ling, X., Yan, B., 2017. Potential toxicity in crucian carp following exposure to metallic nanoparticles of copper, chromium, and their mixtures: a comparative study. *Pol. J. Environ. Stud.* 26 (5), 2085–2094. <https://doi.org/10.15244/pjoes/69251>.
- Madeira, C., Mendonça, V., Flores, A.A.V., Diniz, M.S., Vinagre, C., 2018. High thermal tolerance does not protect from chronic warming—a multiple end-point approach using a tropical gastropod, *Strombina haemastoma*. *Ecol. Indic.* 91, 626–635. <https://doi.org/10.1016/j.ecolind.2018.04.044>.
- Mari Gómez, I., Garmendia, L., Soto, M., Orbe, A., Izagirre, U., & Cajarville, M. P. (2013). Marine ecosystem health status assessment through integrative biomarker indices: A comparative study after the Prestige oil spill “Mussel Watch.” *Ecotoxicology*, 22 (3), 486–505. doi: <https://doi.org/10.1007/s10646-013-1042-4>.
- Marklund, S., Marklund, G., 1974. Involvement of the superoxide anion radical in the autoxidation of pyrogallol and a convenient assay for superoxide dismutase. *Eur. J. Biochem.* 47 (3), 469–474. <https://doi.org/10.1111/j.1432-1033.1974.tb03714.x>.
- Matrangola, V., Corsi, I., 2012. Toxic effects of engineered nanoparticles in the marine environment: model organisms and molecular approaches. *Mar. Environ. Res.* 76, 32–40. <https://doi.org/10.1016/j.marenvres.2012.01.006>.
- Messai, Y., Vileño, B., Martel, D., Turek, P., Mekki, D.E., 2018. Milling effect on the photo-activated properties of TiO₂ nanoparticles: electronic and structural investigations. *Bull. Mater. Sci.* 41 (2). <https://doi.org/10.1007/s12034-018-1572-8>.
- Mezri, A., Alghoo, L.S., Sellami, B., Ben Saber, N., Althali, T., 2018. Titanium dioxide nanoparticles: synthesis, characterization and aquatic ecotoxicity effects. *Chem. Ecol.* 34 (3), 288–299. <https://doi.org/10.1080/02757540.2017.1420178>.
- Mineoto, D., Libralato, G., Volpi Ghirardini, A., 2014. Ecotoxicity of engineered TiO₂ nanoparticles to saltwater organisms: an overview. *Environ. Int.* 66, 18–27. <https://doi.org/10.1016/j.envint.2014.01.012>.
- Mineoto, D., Volpi Ghirardini, A., Libralato, G., 2016. Saltwater ecotoxicology of Ag, Au, CuO, TiO₂, ZnO and CdO engineered nanoparticles: an overview. *Environ. Int.* 92–93, 189–201. <https://doi.org/10.1016/j.envint.2016.03.041>.
- Mouneyrac, C., Buffet, P.-E., Poirier, L., Zalouk-Vergoux, A., Guibolini, M., Favre, C.R., Gilliland, D., Berhan, U., Dybowska, A., Châtel, A., Perrin-Ettajani, H., Pan, J.-F., Thom as-Guyon, H., Reip, P., Valsamis-Jones, E., 2014. Fate and effects of metal-based nanoparticles in two marine invertebrates, the bivalve *Scrobicularia plana* and the annelid *Polychaete Hediste diversicolor*. *Environ. Sci. Pollut. Res.* 21 (13), 7899–7912. <https://doi.org/10.1007/s11356-014-2745-7>.
- Mui, J., Ngo, J., Kim, B., 2016. Aggregation and colloidal stability of commercially available Al₂O₃ nanoparticles in aqueous environments. *Nanomaterials* 6 (5), 90. <https://doi.org/10.3390/nano605090>.
- Murali, M., Athif, P., Suganthi, P., Sadiq Bukhari, A., Syed Mohamed, H.E., Basu, H., Singhal, R.K., 2018. Toxicological effect of Al₂O₃ nanoparticles on histomorphology of the freshwater fish *Oreochromis mossambicus*. *Environ. Toxicol. Pharmacol.* 59, 74–81. <https://doi.org/10.1016/j.etap.2018.03.004>.
- Nam, D.-H., Lee, B., Eom, I., Kim, P., Yeo, M.-K., 2014. Uptake and bioaccumulation of titanium- and silver-nanoparticles in aquatic ecosystems. *Mol. Cell. Toxicol.* 10 (1), 9–17. <https://doi.org/10.1007/s13273-014-0002-2>.
- Nelson, M., Adams, T., Ojo, C., Carroll, M.A., Capatne, E.J., 2018. Mangrove toxicity is targeting an early step in the dopamine signal transduction pathway that controls lateral cilia activity in the bivalve *Mytilus coruscus*. *Comparative Biochemistry and Physiology Part C: Toxicology and Pharmacology* 213, 1–6. <https://doi.org/10.1016/j.cbpc.2018.07.002>.
- Nogueira, D., Vaz, V.P., Neto, O.S., Silva, M.L.N., Simion, I.C., Ouriques, L.C., Vicentini, D.S., & Matias, W.G. (2020a). Crystalline phase-dependent toxicity of aluminum oxide nanoparticles toward *Daphnia magna* and ecological risk assessment. *Environ. Res.*, 182. doi: <https://doi.org/10.1016/j.envres.2019.108987>.

- Neto, O. S., Silva, M. L. N. da, Simioni, C., Ouriques, L. C., Vicentini, D. S., & Matias, W. G. (2020b). Crystalline phase-dependent toxicity of aluminum oxide nanoparticles toward *Daphnia magna* and ecological risk assessment. *Environ. Res.*, 182. doi: <https://doi.org/10.1016/j.envres.2019.108987>.
- Neto, O. S., Arl, M., Köerich, J. S., Simioni, C., Ouriques, L. C., Vicentini, D. S., Matias, W. G., 2019. Comparison of cytotoxicity of α -Al₂O₃ and η -Al₂O₃ nanoparticles toward neuronal and bronchial cells. *Toxicology in Vitro* 61, 104596. <https://doi.org/10.1016/j.tiv.2019.104596>.
- Nunes, S.M., Müller, L., Simioni, C., Ouriques, L.C., Gelesky, M.A., Fattorini, D., Regoli, F., Monserrat, J.M., Ventura-Lima, J., 2020. Impact of different crystalline forms of nTiO₂ on metabolism and arsenic toxicity in *Limnoperna fortunei*. *Sci. Total Environ.* 728. <https://doi.org/10.1016/j.scitotenv.2020.138318>.
- Peng, C., Zhang, W., Gao, H., Li, Y., Tong, X., Li, K., Zhu, X., Wang, Y., Chen, Y., 2017. Behavioral and potential impacts of metal-based engineered nanoparticles in aquatic environments. *Nanomaterials* 7 (1). <https://doi.org/10.3390/nano7010021>.
- Puerrari, R.C., da Costa, C.H., Vicentini, D.S., Fuzinato, C.F., Melegari, S.P., Schmidt, É.C., Bouzon, Z.L., Matias, W.G., 2016. Synthesis, characterization and toxicological evaluation of Cr₂O₃ nanoparticles using *Daphnia magna* and *Alivibacter fischeri*. *Ecotoxicol. Environ. Saf.* 128, 36–43. <https://doi.org/10.1016/j.ecoenv.2016.02.011>.
- R Core Team, 2023. *R: A Language and Environment for Statistical Computing*. R Foundation for Statistical Computing, Vienna, Austria. <https://www.R-project.org/>.
- Ramsden, C.S., Henry, T.B., Handy, R.D., 2013. Sub-lethal effects of titanium dioxide nanoparticles on the physiology and reproduction of zebrafish. *Aquat. Toxicol.* 126, 404–413. <https://doi.org/10.1016/j.aquatox.2012.08.021>.
- Rashidian, G., Mohammadi-Aloucheh, R., Hosseini-adeh-Otaghvari, F., Chupani, L., Stejskal, V., Samadikhah, H., Zamanli, S., Multisanti, C.R., Faggio, C., 2023. Long-term exposure to small-sized silica nanoparticles (SiO₂-NPs) induces oxidative stress and impair reproductive performance in adult zebrafish (*Danio rerio*). *Comparative Biochemistry and Physiology Part C: Toxicology & Pharmacology* 273, 109715. <https://doi.org/10.1016/j.cbpc.2023.109715>.
- Ray, A., Gautam, A., Das, S., Pal, K., Das, S., Karmakar, P., Ray, M., Ray, S., 2020. Effects of copper oxide nanoparticle on gill filtration rate, respiration rate, hemocyte associated immune parameters and oxidative status of an Indian freshwater mussel. *Comparative Biochemistry and Physiology Part C: Toxicology & Pharmacology* 237, <https://doi.org/10.1016/j.cbpc.2020.108855>.
- Regoli, F., Giuliani, M.E., 2014. Oxidative pathways of chemical toxicity and oxidative stress biomarkers in marine organisms. *Mar. Environ. Res.* 93, 106–117. <https://doi.org/10.1016/j.marenvres.2013.07.006>.
- Roche, T.L., Gomes, T., Sousa, V.S., Mestre, N.C., Bebianno, M.J., 2015. Ecotoxicological impact of engineered nanoparticles in bivalve molluscs: an overview. *Mar. Environ. Res.* 111, 74–88. <https://doi.org/10.1016/j.marenvres.2015.06.013>.
- Roma, J., Matos, A.R., Vinagre, C., Duarte, B., 2020. Engineered metal nanoparticles in the marine environment: a review of the effects on marine fauna. *Mar. Environ. Res.* 161. <https://doi.org/10.1016/j.marenvres.2020.105110>.
- Roma, J., Missiouri, M., Madeira, C., Matos, A.R., Vinagre, C., Costa, P.M., Duarte, B., 2024. Comparative responses and effects of exposure to metallic and nanoparticle zinc in the mussel *Mytilus galloprovincialis*. *Estuar. Coast. Shelf Sci.* 297, 108616. <https://doi.org/10.1016/j.ecss.2024.108616>.
- Saad, A.A., El-Sikaily, A., Kassem, H., 2016. Metallothionein and glutathione contents as biomarkers of metal pollution in mussels and local fisherman in Abu Qir Bay. *Egypt. J. Health and Pollution* 6 (12), 50–60. <https://doi.org/10.5696/2156-9614-6-12-50>.
- Saidani, W., Sellami, B., Khazri, A., Mezni, A., Dellali, M., Joubert, O., Sheehan, D., Beyr, H., 2019. Metal accumulation, biochemical and behavioral responses on the Mediterranean clams *Ruditapes decussatus* exposed to two photocatalytic nanocomposites (TiO₂ NPs and AuTiO₂ NPs). *Aquat. Toxicol.* 208, 71–79. <https://doi.org/10.1016/j.aquatox.2019.01.003>.
- Sánchez, A., Recillas, S., Font, X., Cals, E., González, E., Puentes, V., 2011. Ecotoxicity of, and remediation with, engineered inorganic nanoparticles in the environment. *TrAC Trends Anal. Chem.* 30 (3), 507–516. <https://doi.org/10.1016/j.trac.2010.11.011>.
- Sánchez, W., Burgeot, T., Porcher, J.-M., 2013. A novel "integrated biomarker response" calculation based on reference deviation concept. *Environ. Sci. Pollut. Res.* 20 (5), 2721–2725. <https://doi.org/10.1007/s11356-012-1359-1>.
- Sedrati, F., Bouzouane, H., Khaldi, F., Mena, M., Bouarrudj, T., Gzar, L., Zaidi, H., Bensalem, M., Laouar, O., Sleimi, N., Nasri, H., Ouali, K., 2024. In vivo assessment of oxidative stress, neurotoxicity and histological alterations induction in the marine gastropod *Stramonita haemastoma* exposed to Cr₂O₃ and Al₂O₃ nanoparticles. *Chemosphere* 366, 143434. <https://doi.org/10.1016/j.chemosphere.2024.143434>.
- Sharma, A., Vishwakarma, K., Singh, N.K., Prakash, V., Ramawat, N., Prasad, R., Sahi, S., Singh, V.P., Tripathi, D.K., Sharma, S., 2022. Synergistic action of silicon nanoparticles and dodecyl acetic acid in alleviation of chromium (CrVI) toxicity in *Oryza sativa* seedlings. *J. Biotechnol.* 343, 71–82. <https://doi.org/10.1016/j.jbiotec.2021.09.005>.
- Shaw, B.J., Handy, R.D., 2011. Physiological effects of nanoparticles on fish: a comparison of nano metals versus metal ions. *Environ. Int.* 37 (6), 1083–1097. <https://doi.org/10.1016/j.envint.2011.03.009>.
- Singh, S.P., Chandra, S., Kamal, S.S.K., Rahman, M.F., Mahboob, M., Grover, P., 2016. Genotoxic effects of chromium oxide nanoparticles and micro particles in Wistar rats after 28 days of repeated oral exposure. *Environ. Sci. Pollut. Res.* 23 (4), 3914–3924. <https://doi.org/10.1007/s11356-015-5622-0>.
- Thiruprasanth, C.J., Padmini, R., Devraj, S.N., 2009. Effect of berberine on the antioxidant status, ultrastructural modifications and protein bound carbohydrate in azoxymethane-induced colon cancer in rats. *Chem. Biol. Interact.* 177 (3), 190–195. <https://doi.org/10.1016/j.cb.2008.09.027>.
- Tou, F., Wu, J., Fu, J., Niu, Z., Liu, M., Yang, Y., 2021. Titanium and zinc-containing nanoparticles in estuarine sediments: occurrence and their environmental implications. *Sci. Total Environ.* 754, 142388. <https://doi.org/10.1016/j.scitotenv.2020.142388>.
- Turan, N.B., Erkan, H.S., Engin, G.O., Bilgili, M.S., 2019. Nanoparticles in the aquatic environment: usage, properties, transformation and toxicity—a review. *Process. Saf. Environ. Prot.* 130, 238–249. <https://doi.org/10.1016/j.psep.2019.08.014>.
- Vale, G., Mehnado, K., Cambier, S., Libalato, G., Jomini, S., Domingos, R.F., 2016. Manufactured nanoparticles in the aquatic environment: biochemical responses on freshwater organisms: a critical overview. *Aquat. Toxicol.* 170, 162–174. <https://doi.org/10.1016/j.aquatox.2015.11.019>.
- Valko, M., Leibfriz, D., Moncol, J., Cronin, M.T.D., Mazur, M., Telser, J., 2007. Free radicals and antioxidants in normal physiological functions and human disease. *Int. J. Biochem. Cell Biol.* 39 (1), 44–84. <https://doi.org/10.1016/j.biocel.2006.07.001>.
- Viarango, A., Pozzano, E., Dondor, F., Fabbrì, R., 1997. A simple spectrophotometric method for metallothionein evaluation in marine organisms: an application to Mediterranean and Antarctic molluscs. *Mar. Environ. Res.* 44 (1), 69–84. [https://doi.org/10.1016/S0141-1136\(96\)00103-1](https://doi.org/10.1016/S0141-1136(96)00103-1).
- Volland, M., Hampel, M., Martos-Sitcha, J.A., Trombini, C., Martínez-Rodríguez, G., Blasco, J., 2015. Cytotoxic gold nanoparticle exposure in the marine bivalve *Ruditapes philippinarum*: uptake, elimination and oxidative stress response. *Environ. Sci. Pollut. Res.* 22 (22), 17414–17424. <https://doi.org/10.1007/s11356-015-4718-x>.
- Weckbecker, G., Cory, J.G., 1988. Ribonucleotide reductase activity and growth of glutathione-depleted mouse leukemia L1210 cells in vitro. *Cancer Lett.* 40 (3), 257–264. [https://doi.org/10.1016/0304-3835\(88\)90084-5](https://doi.org/10.1016/0304-3835(88)90084-5).
- Wu, H., Zhang, R., Liu, J., Guo, Y., Ma, E., 2011. Effects of malathion and cholinergic acetylcholinesterase and antioxidant defense system in *Oxyechinus* (Thunberg) (Orthoptera: Acrididae). *Chemosphere* 83 (4), 599–604. <https://doi.org/10.1016/j.chemosphere.2010.12.004>.
- Xia, B., Zhu, L., Han, Q., Sun, X., Chen, B., Qu, K., 2017. Effects of TiO₂ nanoparticles at predicted environmental relevant concentration on the marine scallop *Chlamys farreri*: an integrated biomarker approach. *Environ. Sci. Pollut. Res.* 50, 128–135. <https://doi.org/10.1016/j.etap.2017.01.016>.
- Xia, J., Zhao, H.Z., Lu, G.H., 2013. Effects of selected metal oxide nanoparticles on multiple biomarkers in *Carassius auratus*. *Biomol. Biotechnol.* 26 (9), 742–749. <https://doi.org/10.1007/s10575-013-0900-5>.
- Xiong, D., Fang, T., Yu, L., Sima, X., Zhu, W., 2011. Effects of nano-scale TiO₂, ZnO and their bulk counterparts on zebrafish: acute toxicity, oxidative stress and oxidative damage. *Sci. Total Environ.* 409 (8), 1444–1452. <https://doi.org/10.1016/j.scitotenv.2011.01.015>.
- Yung, M.M.N., Moueyrac, C., Leung, K.M.Y., 2014. Ecotoxicity of zinc oxide nanoparticles in the marine environment. In: *Encyclopedia of Nanotechnology*. Springer, Netherlands, pp. 1–17. https://doi.org/10.1007/978-94-007-6178-0_10970-1.
- Zaidi, H., Amrani, A., Sedrati, F., Maaruf, H., Leghrib, F., Benamara, M., Amar, H., Wang, Z., Nasri, H., 2021. Histological and chemical damage induced by microcystin-LR and microcystin-RR on land snail *Helix aspersa* tissues after acute exposure. *Comparative Biochemistry and Physiology Part C: Toxicology and Pharmacology* 245. <https://doi.org/10.1016/j.cbpc.2021.109031>.
- Zaleśka-Radziwiłł, M., Doskocz, N., Affek, K., Muszyński, A., 2020. Effect of aluminum oxide nanoparticles on aquatic organisms – a microcosm study. *Desalination Water Treat.* 195, 286–296. <https://doi.org/10.5004/dwt.2020.25882>.
- Zhao, Y., Xie, P., Zhang, X., 2009. Oxidative stress response after prolonged exposure of domestic rabbit to a lower dosage of extracted microcystins. *Environ. Sci. Pollut. Res.* 16 (2), 195–199. <https://doi.org/10.1016/j.etap.2008.10.005>.